

10-22-2011

## Prescribing exercise in primary care: Ten practical steps on how to do it

Karim M. Khan

Richard Weller

Steven N. Blair

*University of South Carolina - Columbia*, [sblair@mailbox.sc.edu](mailto:sblair@mailbox.sc.edu)

Follow this and additional works at: [https://scholarcommons.sc.edu/sph\\_epidemiology\\_biostatistics\\_facpub](https://scholarcommons.sc.edu/sph_epidemiology_biostatistics_facpub)



Part of the [Public Health Commons](#)

---

### Publication Info

Published in *British Medical Journal*, Volume 343, Issue d4141, 2011, pages 806-807.

This Article is brought to you by the Epidemiology and Biostatistics at Scholar Commons. It has been accepted for inclusion in Faculty Publications by an authorized administrator of Scholar Commons. For more information, please contact [digres@mailbox.sc.edu](mailto:digres@mailbox.sc.edu).

bmj.com

▶ Read David Pencheon's blogs at [bmj.com/blogs](http://bmj.com/blogs)

Secondly, agreement is needed on how to cost and value the immediate and longer term health benefits of mitigating climate change. The multiple health benefits of reducing carbon emissions and their effect on the economy are yet to be systematically captured and valued. The importance of doing this was recognised in the Stern review.<sup>12</sup>

Designing and agreeing systematic measures of the health benefits of taking action requires interdisciplinary collaboration between health professionals, economists, and climate change scientists. It also involves interagency collaboration between the Department of Energy and Climate Change; the Department for Environment, Food and Rural Affairs; the Department of Health; and the National Institute for Health and Clinical Excellence to ensure that this is done in rigorous ways that benefit patients, the public, and future generations. These methods need to quantify the multiple benefits in ways that stimulate action from all parts of the global health system, from local nurses and doctors to global drug companies.

- 1 Costello A, Abbas M, Allen A, Ball S, Bell S, Bellamy R, et al. Managing the health effects of climate change. *Lancet* 2009;373:1693-733.
- 2 Jarvis L, Montgomery H, Morisetti N, Gilmore I. Climate change, ill health, and conflict. *BMJ* 2011;342: d1819.
- 3 Gilding P. The great disruption: how the climate crisis will transform the global economy. Bloomsbury, 2011.
- 4 Griffiths J, Rao M. Public health benefits of strategies to reduce greenhouse gas emissions. *BMJ* 2009;339:b4952.
- 5 Roberts I, Edwards P. The energy glut: the politics of fatness in an overheating world. Zen Books, 2010.
- 6 Zander A, Niggebrugge A, Pencheon D, Lyratzopoulos G. Changes in travel-related carbon emissions associated with modernization of services for patients with acute myocardial infarction: a case study. *J Public Health* 2011;33:272-9.
- 7 Connor A, Mortimer F, Higgins R. The follow-up of renal transplant recipients by telephone consultation: three years experience from a single UK renal unit. *Clin Med* 2011;11:242-6.
- 8 NHS Sustainable Development Unit, Forum for the Future. Fit for the Future. Scenarios for low-carbon healthcare 2030. Cambridge: 2009.
- 9 HM Treasury. The green book. 2011. [www.hm-treasury.gov.uk/data\\_greenbook\\_index.htm](http://www.hm-treasury.gov.uk/data_greenbook_index.htm).
- 10 Atkinson S, Ingham J, Cheshire M, Went S. Defining quality and quality improvement. *Clin Med* 2010;10:537-9.
- 11 World Bank. Data. 2009. <http://data.worldbank.org/indicator/>.
- 12 Stern N. Stern review on the economics of climate change. 2006. [http://webarchive.nationalarchives.gov.uk/+http://www.hm-treasury.gov.uk/sternreview\\_index.htm](http://webarchive.nationalarchives.gov.uk/+http://www.hm-treasury.gov.uk/sternreview_index.htm).

## Prescribing exercise in primary care

Ten practical steps on how to do it

Chronic diseases—the leading causes of morbidity and mortality—are strongly linked to unhealthy lifestyles. The World Health Organization recently published a report on global health risks.<sup>1</sup> Leading causes of global mortality are high blood pressure (13% of total deaths), tobacco use (9%), high blood glucose (6%), physical inactivity (6%), and obesity (5%). Physical inactivity is fourth on this list, but it influences most of the other causes. The importance of physical activity in preventing and treating many diseases and conditions is indisputable, as documented by the authoritative, accessible, and practical guide of the Swedish Professional Associations for Physical Activity.<sup>2</sup> Reports from the United States,<sup>3</sup> Canada, and the United Kingdom concur.<sup>4</sup>

Physical activity is a complex behaviour, and modern physical and social environments discourage it. Countering these influences will require a coordinated approach involving multiple societal, institutional, and departmental collaborations.<sup>5</sup> Clinicians need to contribute. Healthcare is one of eight sectors listed in the US national physical activity plan that aims to influence activity ([www.physicalactivity-plan.org/theplan.php](http://www.physicalactivity-plan.org/theplan.php)).<sup>6</sup> Advice on physical activity in primary care is a pillar of WHO's global physical activity plan (2010).<sup>7</sup> Physical activity counselling in clinical settings provides "exceptional value for money."<sup>4</sup>

Based on the recommendation of the Swedish physical activity book,<sup>2</sup> key principles of the US physical activity plan, and the UK's Department of Health's *Let's Get Moving*,<sup>8</sup> the following steps can help general practitioners encourage patients to initiate and maintain a physically active lifestyle (box).

As with any clinical consultation, diagnosis is key. Physical inactivity is a modifiable vital sign. Thus, GPs should begin by administering the UK's general practice physical activity questionnaire (GPPAQ)<sup>9</sup> or asking the exercise vital sign (EVS) questions.<sup>10</sup> The GPPAQ takes 30 seconds

to administer plus two to three minutes to categorise patients into one of four levels of activity. The "exercise is medicine" initiative ([www.exerciseismedicine.org](http://www.exerciseismedicine.org)) recommends asking two EVS questions,<sup>10</sup> so that each patient's weekly activity can be compared with the recommended guideline. The results should be incorporated into the electronic medical record.<sup>10</sup> From this diagnosis, management begins.

General practitioners should use their counselling method of choice and apply it to physical activity. Most general practitioners are familiar with the "6As" to guide counselling—assess, advise, agree, assist, arrange, and assess again. This approach is discussed using a case study in the Swedish book. Motivational interviewing is an increasingly popular method for GPs to counsel patients about their lifestyles.<sup>2</sup> This, and other behavioural therapy techniques, are valuable skills and provide a worthy CME (continuing medical education) activity.

### Practical steps for immediate exercise prescription in general practice

- Ask about physical activity at every consultation; consider it a vital sign
- Apply the "6As" to guide counselling—assess, advise, agree, assist, arrange, and assess again
- A written ("green") prescription is crucial—it takes just 30 seconds
- Display a poster with the physical activity guidelines prominently in the waiting room
- Consider categorising patients into frailty levels. There is no need to medicalise physical activity for most people
- Refer on—consider appropriate physicians, physiotherapists, clinical exercise physiologists, and certified fitness instructors
- Know your local resources for activity—the people and the places
- Remember that walking is free; find tips at: [www.everybodywalk.org](http://www.everybodywalk.org)
- Follow up the patient to chart progress, set goals, solve problems, and identify and use social support
- Lobby to make low cost, evidence based, cognitive and behavioural interventions widely available for referral by healthcare providers

**Karim M Khan** professor, Department of Family Practice, University of British Columbia, Vancouver, BC, Canada V6T 1Z4 [karim.khan@ubc.ca](mailto:karim.khan@ubc.ca)

**Richard Weiler** specialist in sport and exercise medicine, Sport and Musculoskeletal Medicine Clinic, Homerton University Hospital NHS Foundation Trust, London, UK

**Steven N Blair** professor, Arnold School of Public Health, Public Health Research Center, University of South Carolina, Columbia, SC, USA

**Competing interests:** All authors have completed the ICMJE uniform disclosure form at [www.icmje.org/coi\\_disclosure.pdf](http://www.icmje.org/coi_disclosure.pdf) (available on request from the corresponding author) and declare: no support from any organisation for the submitted work; no financial relationships with any organisations that might have an interest in the submitted work in the previous three years; no other relationships or activities that could appear to have influenced the submitted work.

**Provenance and peer review:** Commissioned; not externally peer reviewed.

Cite this as: *BMJ* 2011;343:d4141 doi: 10.1136/bmj.d4141

doc2doc

▶ Legacy of the Olympics—will more people take up exercise? <http://bit.ly/nrVfBU>



The written “green” prescription (which comprises exercise and lifestyle goals) is a crucial element to signal that exercise is medicine. As outlined in chapter 3 of the Swedish book, the written prescription may vary from simple advice on activity, including written information on health benefits and national guidelines, to highlighting local physical activity opportunities. The national physical activity guidelines should be prominent in the waiting room.

As one prescription does not fit all, a useful tip in the Swedish book is to categorise the prescription according to four levels of patient frailty.<sup>2</sup> The frailest are referred to appropriate health professionals (such as physiotherapists). For patients with stable conditions, general practitioners can provide customised “exercise on prescription,” (outlined by condition in the book). Healthier patients are encouraged to join community support groups and to manage their own physical activity programme.

Whether a doctor should manage a patient alone or refer the patient depends on local skills, experience, services, and patient care pathways. Doctors not skilled and trained in exercise prescription, probably a majority in most countries, should feel confident to refer on. In many countries including the UK, referral might be to a sports and exercise medicine specialist, in others such as the US, referral may be to a primary care sports medicine physician.

Referral will generally reach beyond health professionals alone. It is useful to link patients with appropriate community based resources. Only a few people have the resources to hire a health professional or a fitness instructor to design and supervise a workout. Communities, however, often have low cost public facilities. Walking is free and now being promoted on a very practical, user friendly, and unbranded website ([www.everybodywalk.org](http://www.everybodywalk.org)).

Follow-up is crucial to signal the clinician’s conviction, determine the patient’s progress, solve problems, help identify social support, fine tune the dose, and reset goals. The exercise vital sign should be obtained at every follow-up visit.<sup>9</sup> <sup>10</sup>Doctors do not prescribe an antihypertensive drug at one dose and then ignore the patient for ever.

At the medical society level, general practitioner organisations should take time to vote on whether they support exercise as medicine and are prepared to commit resources to appropriate political action. If they do, it would be logical

to lobby to make low cost, evidence based, cognitive and behavioural interventions widely available for referral by healthcare providers. There is also evidence for high tech patient support applications, such as email and text message reminders, website support, and smart phone apps.<sup>11</sup> Such innovations align with the current UK government’s approach to “nudge” people towards healthier behaviour by subtly changing environments.

In addition to these steps, medical education curriculums need to include the promotion of physical activity. A new generation of doctors must avoid the knee jerk reaction of prescribing “preventive” drugs as a first response to diseases of inactivity. There is strong economic evidence that to manage lifestyle diseases, doctors should first encourage patients to adopt a healthy lifestyle and then help them to maintain it.<sup>8</sup> This is particularly relevant given the budget based debates on health system reform in the NHS and in the US right now.

It is obvious that clinicians must be active role models. As well as improving their own health, 22 minutes of physical activity daily provides credibility when prescribing and highlights the problems faced by patients.<sup>3</sup>

- 1 WHO. Global health risks: mortality and burden of disease attributable to selected major risks. 2009. [www.who.int/healthinfo/global\\_burden\\_disease/GlobalHealthRisks\\_report\\_full.pdf](http://www.who.int/healthinfo/global_burden_disease/GlobalHealthRisks_report_full.pdf).
- 2 Swedish Professional Associations for Physical Activity. Physical activity in the prevention and treatment of disease. 2011. [www.fhi.se/PageFiles/10682/Physical-Activity-Prevention-Treatment-Disease-webb.pdf](http://www.fhi.se/PageFiles/10682/Physical-Activity-Prevention-Treatment-Disease-webb.pdf).
- 3 Office of Disease Prevention and Health Promotion. 2008 Physical Activity Guidelines for Americans. 2008. [www.health.gov/paguidelines/pdf/paguide.pdf](http://www.health.gov/paguidelines/pdf/paguide.pdf).
- 4 National Institute for Health and Clinical Excellence. Four commonly used methods to increase physical activity. 2006. [www.nice.org.uk/nicemedia/pdf/PHYSICAL-ALS2\\_FINAL.pdf](http://www.nice.org.uk/nicemedia/pdf/PHYSICAL-ALS2_FINAL.pdf).
- 5 Waxman A. World Health Assembly: WHO global strategy on diet, physical activity and health. *Food Nutr Bull* 2004;25:292-302.
- 6 Patrick K, Pratt M, Sallis RE. The healthcare sector’s role in the U.S. national physical activity plan. *J Phys Act Health* 2009;6(suppl 2):S211-9.
- 7 WHO. Global recommendations on physical activity for health. 2010. [http://whqlibdoc.who.int/publications/2010/9789241599979\\_eng.pdf](http://whqlibdoc.who.int/publications/2010/9789241599979_eng.pdf).
- 8 NHS Department of Health. Let’s get moving. Commissioning guidance. 2009. [www.paha.org.uk/Resource/lets-get-moving-commissioning-guidance](http://www.paha.org.uk/Resource/lets-get-moving-commissioning-guidance).
- 9 Physical Activity Policy and Health Improvement Directorate. The general practice physical activity questionnaire (GPPAQ): a screening tool to assess adult physical activity levels, within primary care. 2009. [www.dh.gov.uk/prod\\_consum\\_dh/groups/dh\\_digitalassets/documents/digitalasset/dh\\_101579.pdf](http://www.dh.gov.uk/prod_consum_dh/groups/dh_digitalassets/documents/digitalasset/dh_101579.pdf).
- 10 Sallis R. Developing healthcare systems to support exercise: exercise as the fifth vital sign. *Br J Sports Med* 2010;45:473-4.
- 11 Marcus BH, Ciccolo JT, Sciamanna CN. Using electronic/computer interventions to promote physical activity. *Br J Sports Med* 2009;43:102-5.

## The future of the primary medical workforce

Is in jeopardy unless GPs have a more central role in healthcare and develop special skills in areas of unmet need

Yet another study has reported increasing difficulty in recruiting medical students to careers in general internal medicine. This study from the United States showed that although 24% of 1177 students who graduated in 2007 planned careers in internal medicine, they perceived this specialty to be associated with a higher workload and more stress than other specialties and were less interested in a career in general internal medicine than students in a similar study in 1990.<sup>1</sup> The trends in these data are not

dissimilar to those reported in relation to general practice and family medicine in Australia,<sup>2</sup> New Zealand and Canada,<sup>3</sup> and the United Kingdom.<sup>4</sup>

Recruitment to general practice has improved a little in Australia in recent years since the creation of rural clinical schools that provide undergraduates with at least half of their clinical training in rural environments. But the fact remains that around the world primary medical care is not a favoured career choice for specialty training and practice.

Cite this as: *BMJ* 2011;343:d5006  
doi: 10.1136/bmj.d5006