

12-17-2010

Parity violation in the $n + 3\text{He} \rightarrow 3\text{H} + p$ reaction: Resonance approach

Vladimir Gudkov

University of South Carolina - Columbia, gudkov@sc.edu

Follow this and additional works at: http://scholarcommons.sc.edu/phys_facpub

 Part of the [Physics Commons](#)

Publication Info

Published in *Physical Review C*, Volume 82, Issue 6, 2010, pages 065502-1-065502-5.

Gudkov, V. (2010). Parity violation in the $n + 3\text{He} \rightarrow 3\text{H} + p$ reaction: Resonance approach. *Physical Review C*, 82(6), 065502-1 - 065502-5. doi: 10.1103/PhysRevC.82.065502

© 2010 The American Physical Society.

This Article is brought to you for free and open access by the Physics and Astronomy, Department of at Scholar Commons. It has been accepted for inclusion in Faculty Publications by an authorized administrator of Scholar Commons. For more information, please contact SCHOLARC@mailbox.sc.edu.

Parity violation in the $n + {}^3\text{He} \rightarrow {}^3\text{H} + p$ reaction: Resonance approach

Vladimir Gudkov*

Department of Physics and Astronomy, University of South Carolina, Columbia, South Carolina 29208, USA

(Received 18 August 2010; published 17 December 2010)

The method based on microscopic theory of nuclear reactions has been applied for the analysis of parity-violating effects in few-body systems. Different parity-violating and parity-conserving asymmetries and their dependence on neutron energy have been estimated for the $n + {}^3\text{He} \rightarrow {}^3\text{H} + p$ reaction. The estimated effects are in a good agreement with available exact calculations.

DOI: [10.1103/PhysRevC.82.065502](https://doi.org/10.1103/PhysRevC.82.065502)

PACS number(s): 24.80.+y, 25.10.+s, 25.40.Ny

I. INTRODUCTION

The study of parity-violating (PV) effects in low-energy physics is important for the understanding of the main features of the standard model and for the possible search for manifestations of new physics. During the past decades, many calculations of different experimentally observed PV effects in nuclear physics have been done. However, in the past years it became clear (see, for example, Refs. [1–4] and references therein) that the traditional Desplanques, Donoghue, and Holstein (DDH) [5] method for the calculation of PV effects cannot reliably describe the available experimental data. This could be blamed on “wrong” experimental data; however, it might be that the DDH approach is not adequate for the description of the set of precise experimental data because it is based on a number of models and assumptions. To resolve this discrepancy, it is desirable to increase the number of experimental data for different PV parameters in few-body systems. The calculations of nuclear related effects for these systems can be done with high precision, eliminating as many nuclear-model-dependent factors involved in PV effects as possible. Unfortunately, currently available data of experimentally measured PV effects in these systems are not enough to constrain all parameters required for calculations; therefore, any new potentially possible measurement is very important. Because PV effects in few-body systems are usually very small and precise calculations of them are rather difficult, it is desirable to have a method for a reliable estimate of possible observable parameters using available experimental data. This will give the opportunity to choose the right system and right PV observables for new experiments.

Recently, it has been proposed to measure PV asymmetry of protons in the $n + {}^3\text{He} \rightarrow {}^3\text{H} + p$ reaction with polarized neutrons at the Spallation Neutron Source at the Oak Ridge National Laboratory. The ${}^3\text{He}$ and ${}^4\text{He}$ systems were subjects of intensive investigation for a long time, and as a result, many parameters related to reactions with neutrons and protons, as well as to excitation energy levels of these nuclei, have been measured and evaluated by a number of different groups. These rather comprehensive data provide the opportunity to estimate values of possible PV effects and their dependence on neutron

energy in the $n + {}^3\text{He} \rightarrow {}^3\text{H} + p$ reaction using a microscopic nuclear reaction theory approach.

II. DESCRIPTION OF PARITY-VIOLATING EFFECTS

Let us consider the $n + {}^3\text{He} \rightarrow {}^3\text{H} + p$ reaction with low-energy neutrons. For neutron energy $E_n \sim 0.01$ eV, which corresponds to a wave vector $k_n \sim 2.19 \times 10^{-5}$ fm $^{-1}$, the energy of outgoing protons and proton wave vector are $E_p = 0.764$ MeV and $k_p = 0.19$ fm $^{-1}$, correspondingly. Taking a characteristic ${}^3\text{He}$ radius as $R = 1.97$ fm, one obtains $(k_n R) \sim 4 \times 10^{-4}$ and $(k_p R) \sim 0.4$. Therefore, for the initial channel, contributions from p -wave neutrons to a reaction matrix (amplitude) are highly suppressed, whereas for the final channel, the amplitude with orbital momenta of protons $l = 0$ and $l = 1$ have the same order of magnitude. The contribution from d -wave protons is suppressed by a factor of ~ 0.025 ; therefore, one can ignore d waves within the accuracy of our estimates. Assuming that neutrons, as well as ${}^3\text{He}$ nuclei, may have a polarization, one shall consider four PV and four parity-conserving (PC) angular correlations, shown in Table I, where $\vec{\sigma}$ and \vec{I} are neutron and nuclear spins. (It is important to know the values of PC correlations because they usually are one of the main sources of experimental errors for PV effects and also because they can be rather easily measured and, as a consequence, can serve as an indirect proof of the correctness of calculations of PV effects.) We focus on PV correlation $(\vec{\sigma} \cdot \vec{k}_p)$, which leads to PV asymmetry α_{PV} of outgoing protons in a direction along the neutron polarization and opposite to it. It can be seen that the asymmetry related to the $(\vec{I} \cdot \vec{k}_p)$ correlation has exactly the same value for this reaction. For the completeness of the consideration, we also calculate the PV effect related to differences of total cross sections $\sigma_{\pm}^{\text{tot}}$ for neutrons with opposite helicities when they propagate through a ${}^3\text{He}$ target, $P = (\sigma_{+}^{\text{tot}} - \sigma_{-}^{\text{tot}})/(\sigma_{+}^{\text{tot}} + \sigma_{-}^{\text{tot}})$, which is related to the $(\vec{\sigma} \cdot \vec{k}_n)$ correlation. The corresponding difference of total cross sections for a propagation of unpolarized neutrons through polarized target, the $(\vec{I} \cdot \vec{k}_n)$ correlation, has the same value P in our case. It should be noted that the $(\vec{\sigma} \cdot \vec{k}_n)$ correlation leads also to PV neutron spin precession around the direction of neutron momentum. However, the angle of precession is very small (about 10^{-9} – 10^{-10} rad for thermal neutrons, and it can reach a value up to about 10^{-5} rad for $E_n \sim 0.5$ MeV for the target of the size of neutron mean free path); therefore, we do not consider it here. In addition, we

*gudkov@sc.edu

TABLE I. Possible parity-violating and four parity-conserving angular correlations.

PV	PC
$(\vec{\sigma} \cdot \vec{k}_p)$	$(\vec{k}_n \cdot \vec{k}_p)$
$(\vec{\sigma} \cdot \vec{k}_n)$	$(\vec{\sigma} \cdot [\vec{k}_n \times \vec{k}_p])$
$(\vec{I} \cdot \vec{k}_p)$	$(\vec{\sigma} \cdot \vec{I})$
$(\vec{I} \cdot \vec{k}_n)$	$(\vec{I} \cdot [\vec{k}_n \times \vec{k}_p])$

calculate the left-right asymmetry α_{LR} which corresponds to the PC correlation $(\vec{\sigma} \cdot [\vec{k}_p \times \vec{k}_n])$ because it could be a source of systematic effects in the measurement of α_{PV} .

Using standard techniques (see, for example, Ref. [6]), one can represent these asymmetries in terms of matrix \hat{R} which is related to reaction matrix \hat{T} and to S matrix as

$$\hat{R} = 2\pi i \hat{T} = \hat{1} - \hat{S}. \quad (1)$$

Then, for our case

$$\alpha_{PV} = \frac{2}{r} \text{Re}[-3\sqrt{2}\langle 01|R^1|10\rangle\langle 00|R^0|00\rangle^* + (\sqrt{6}\langle 11|R^0|00\rangle + 6\langle 11|R^1|10\rangle)\langle 10|R^1|10\rangle^*] \quad (2)$$

and

$$\alpha_{LR} = \frac{1}{r} \text{Im}[6\sqrt{3}\langle 10|R^1|11\rangle\langle 00|R^0|00\rangle^* + 6\sqrt{3}\langle 11|R^1|01\rangle\langle 10|R^1|10\rangle^* + 3(2\langle 11|R^0|00\rangle + \sqrt{6}\langle 11|R^1|10\rangle)\langle 10|R^1|11\rangle^* + 6\sqrt{2}\langle 00|R^0|11\rangle\langle 01|R^1|10\rangle^* + 6\sqrt{3}\langle 10|R^1|01\rangle\langle 11|R^1|10\rangle^* + \sqrt{6}\langle 10|R^1|10\rangle(2\langle 11|R^0|11\rangle^* + 3\langle 11|R^1|11\rangle^*) + 5\sqrt{6}\langle 11|R^2|11\rangle\langle 10|R^1|10\rangle^*], \quad (3)$$

where

$$r = (|\langle 00|R^0|00\rangle|^2 + 3|\langle 10|R^1|10\rangle|^2). \quad (4)$$

We use spin-channel representation, where for the matrix element $\langle s'l'|R^J|sl\rangle$, l and l' are orbital momenta of initial and final channels with corresponding spin channels s and s' , respectively, and J is the total spin of the system. For a transmission type of observable P , one obtains

$$P = -\frac{\text{Re}[\langle 00|R^0|11\rangle + \langle 11|R^0|00\rangle]}{\text{Re}[\langle 00|R^0|00\rangle + 3\langle 10|R^1|10\rangle]} + \sqrt{3}\frac{\text{Re}[\langle 10|R^1|01\rangle + \langle 01|R^1|10\rangle]}{\text{Re}[\langle 00|R^0|00\rangle + 3\langle 10|R^1|10\rangle]} + \sqrt{6}\frac{\text{Re}[\langle 10|R^1|11\rangle + \langle 11|R^1|10\rangle]}{\text{Re}[\langle 00|R^0|00\rangle + 3\langle 10|R^1|10\rangle]}.$$

Calculations of matrix elements $\langle s'l'|R^J|sl\rangle$ for PV effects in nuclear reactions have been done [6] using distorted-wave Born approximation in microscopic theory of nuclear reactions [7]. They lead to the symmetry-violating amplitudes induced by PV potential W ,

$$R_{PV}^{fi} = 2\pi i \langle \Psi_f^- | W | \Psi_i^+ \rangle, \quad (5)$$

where $\Psi_{i,f}^\pm$ are the eigenfunctions of the nuclear P -invariant Hamiltonian with the appropriate boundary conditions [7]:

$$\Psi_{i,f}^\pm = \sum_k a_{k(i,f)}^\pm(E) \phi_k + \sum_m \int b_{m(i,f)}^\pm(E, E') \chi_m^\pm(E') dE'. \quad (6)$$

Here ϕ_k is the wave function of the k th resonance and $\chi_m^\pm(E)$ is the potential scattering wave function in the channel m . The coefficient

$$a_{k(i,f)}^\pm(E) = \frac{\exp(\pm i\delta_{i,f})}{(2\pi)^{\frac{1}{2}}} \frac{(\Gamma_k^{i,f})^{\frac{1}{2}}}{E - E_k \pm \frac{i}{2}\Gamma_k} \quad (7)$$

describes nuclear resonances contributions, and the coefficient $b_{m(i,f)}^\pm(E, E')$ describes potential scattering and interactions between the continuous spectrum and resonances. [Here E_k , Γ_k , and Γ_k^i are the energy, the total width, and the partial width in the channel i of the k th resonance, respectively, E is the neutron energy, and δ_i is the potential scattering phase in the channel i ; $(\Gamma_k^i)^{\frac{1}{2}} = (2\pi)^{\frac{1}{2}} \langle \chi_i(E) | V | \phi_k \rangle$, where V is a residual interaction operator.] As was shown in [6] for nuclei with rather large atomic numbers, the resonance contribution is dominant. Then, for the simplest case with only two resonances with opposite parities, the expression for matrix element \hat{R} for a neutron-proton reaction with parity violation is

$$\langle s'l'|R^J|sl\rangle = -\frac{iw[\Gamma_l^n(s)\Gamma_{l'}^p(s')]^{\frac{1}{2}}}{(E - E_l + i\Gamma_l/2)(E - E_{l'} + i\Gamma_{l'}/2)} e^{i(\delta_l^n + \delta_{l'}^p)}, \quad (8)$$

and with conservation of parity (one resonance contribution) is

$$\langle s'l'|R^J|sl\rangle = \frac{i[\Gamma_l^n(s)\Gamma_{l'}^p(s')]^{\frac{1}{2}}}{(E - E_l + i\Gamma_l/2)} e^{i(\delta_l^n + \delta_{l'}^p)}, \quad (9)$$

where $w = -\int \phi_l W \phi_{l'} d\tau$ is PV nuclear matrix element mixing parities of two resonances. The preceding \hat{R} matrix elements could be represented by the diagrams shown on Fig. 1. Thus, PV asymmetry α_{PV} is proportional to the real part the product of \hat{R} matrices presented by diagrams (a) and (c), and PC asymmetry α_{LR} is proportional to the imaginary part of the product of \hat{R} matrices presented by diagrams (a) and (b) (for details, see [6]).




FIG. 1. Parity-conserving [diagrams (a) and (b)] and parity-violating [diagrams (c) and (d)] matrix elements of \hat{R} . Symbols s and p correspond to s - and p -wave neutrons for the initial (neutron) channel and indicate the parity of the final channel.

TABLE II. Resonance parameters (Set 1). Here E_r is a resonance energy; T and J^π are resonance isospin and the total resonance spin with parity, respectively; Γ and Γ_p are total and proton widths; Γ_n , Γ_n^0 , and l are neutron width, reduced width, and angular momentum, correspondingly; and α_{PV} (%) and P (%) are normalized contribution of the resonance to α_{PV} and P , correspondingly.

E_r (MeV)	J^π	l	T	Γ_n (MeV)	Γ_n^0 (eV)	Γ_p (MeV)	Γ (MeV)	α_{PV} (%)	P (%)
-0.211	0+	0	0		954.4	1.153	1.153		
0.430	0-	1	0	0.48		0.05	0.53	3.1	100
3.062	1-	1	1	2.76		3.44	6.20	100 ± 26	2 ± 1
3.672	1-	1	0	2.87		3.08	6.10	75 ± 24	1 ± 1
4.702	0-	1	1	3.85		4.12	7.97	20	3
5.372	1-	1	1	6.14		6.52	12.66	79 ± 18	1
7.732	1+	0	0	4.66		4.725	9.89		
7.792	1-	1	0	0.08		0.07	3.92	2 ± 1	0
8.062	0-	1	0	0.01		0.01	4.89	14	0

This technique has been proven to work very well for calculation of nuclear PV effects for intermediate and heavy nuclei. Assuming the dominant resonance contribution to PV effects for the $n + {}^3\text{He} \rightarrow {}^3\text{H} + p$ reaction, we apply this approach to estimate characteristic values of PV effects using parametrization of PV effects in terms of known resonance structure of the system. Fortunately, the detailed structure of resonances (${}^4\text{He}$ levels) [8] and low-energy neutron scattering parameters [9] are well known for this reaction from numerous experiments.

To estimate PV and PC asymmetries in $n + {}^3\text{He} \rightarrow {}^3\text{H} + p$ reactions using the previously described formalism, we take into account all known resonances [8,9] which result in multiresonance representation for \hat{R} matrix elements. From the selection rules for angular momenta (see Eqs. (2), (3), (5) and general expressions in [6]), one can see that for low-energy neutrons only resonances with the total spin of $J = 0, 1$ contribute to PV asymmetries of the interest. However, for the left-right PC asymmetry, we have to consider $J = 0, 1, 2$. Thus, for PV effects we consider contributions from nine low-energy resonances [8,9] (see Table II): one resonance with total angular momentum and parity $J^\pi = 0^+$, three with $J^\pi = 0^-$, four with $J^\pi = 1^-$, and one with $J^\pi = 1^+$. For further calculations, we assume that all weak matrix elements, which mix resonances with opposite parities, have the same values and are described by the phenomenological formula [6]

$w = 2 \times 10^{-4} \text{ eV} \sqrt{\bar{D}} \text{ (eV)}$ (where \bar{D} is an average energy level spacing). This formula is in good agreement with other statistical nuclear model estimates [10–12] of nuclear weak matrix elements for medium and heavy nuclei. The extrapolation of this formula to the region of one-particle nuclear excitation leads to the correct value for weak nucleon-nucleon interaction. Therefore, one can use this approximation for rough estimates of average values of weak matrix elements in few-body systems. This leads to the value of weak matrix element $w = 0.5 \text{ eV}$ (with $\bar{D} \simeq 6 \text{ MeV}$), which is rather close to the typical value of one-particle weak matrix element. One can see from Eqs. (8) and (9) that the expressions for PV and PC \hat{R} matrices depend not on neutron and proton partial widths but on their amplitudes, the values of which depend on particular spin channels. Because we know only partial widths, we have to make assumptions about values of amplitudes of partial widths for a specific spin channel and about their signs (phases). This leads to another uncertainty in our estimation in addition to the previously given assumption about weak matrix elements. To treat the spin-channel dependence of partial width amplitudes, we assume that partial widths for each spin channel are equal to each other. This gives us an average factor of uncertainty of about 2. The signs of width amplitudes, as well as the signs of weak matrix elements w , are left undetermined (random). This also can lead to a factor of uncertainty of 2 or 3. Therefore, one can see that the

TABLE III. Resonance parameters (Set 2). Here E_r is a resonance energy; T and J^π are resonance isospin and the total resonance spin with parity, respectively; Γ and Γ_p are total and proton widths; Γ_n , Γ_n^0 , and l are neutron width, reduced width, and angular momentum, correspondingly; and α_{PV} (%) and P (%) are normalized contribution of the resonance to α_{PV} and P , correspondingly.

E_r (MeV)	J^π	l	T	Γ_n (MeV)	Γ_n^0 (eV)	Γ_p (MeV)	Γ (MeV)	α_{PV} (%)	P (%)
-0.211	0+	0	0		954.4	1.153	1.153		
0.430	0-	1	0	0.20		0.640	0.84	100	100
3.062	1-	1	1	2.76		3.44	6.20	82 ± 27	4 ± 3
3.672	1-	1	0	2.87		3.08	6.10	62 ± 20	3 ± 2
4.702	0-	1	1	3.85		4.12	7.97	16	8
5.372	1-	1	1	6.14		6.52	12.66	65 ± 15	2 ± 1.5
7.732	1+	0	0	4.66		4.725	9.89		
7.792	1-	1	0	0.08		0.07	3.92	2 ± 1	0
8.062	0-	1	0	0.01		0.01	4.89	1	0.5




FIG. 2. (Color online) Resonance enhancement of the P parameter.

uncertainty of our multiresonance calculations is about of one order of magnitude.

III. DISCUSSION OF RESULT OF CALCULATIONS

Taking into account the considerations given earlier and using resonance parameters [8,9] of the Table II, one can estimate the PV asymmetry for thermal neutrons as

$$\alpha_{PV} = -(1-4) \times 10^{-7}. \quad (10)$$

The set of resonance parameters of the Table III results in a slightly larger PV asymmetry:

$$\alpha_{PV} = -(4-8) \times 10^{-7}. \quad (11)$$

The difference of these two sets is related to the discrepancy between [9] and [8] for resonance parameters for the first positive resonance ($E_n = 0.430$ MeV).

The left-right asymmetry at thermal energy is less sensitive to the parameters of this first resonance and has about the same value for these two sets of resonance parameters:

$$\alpha_{LR} = -(2-8) \times 10^{-4}. \quad (12)$$

It should be noted that for the calculation of the left-right asymmetry, we add first three 2^- resonances [8], which do not contribute to PV effects for low-energy neutrons.

The value of PV in neutron transmission for the first choice of parameters is

$$P = -(2-4) \times 10^{-10} \quad (13)$$

and for the second choice is

$$P = -(0.8-1.6) \times 10^{-10}. \quad (14)$$

One can see that the parameter P is very small for neutrons with thermal energy, but it is essentially enhanced in a few-MeV region (see Fig. 2). The asymmetry α_{PV} also shows




FIG. 3. (Color online) Resonance enhancement of the α_{PV} asymmetry (for the first set of parameters).

a resonance behavior, but its enhancement is not very large (see Fig. 3).

To show contributions of each resonance to PV asymmetry α_{PV} and to transmission parameter P , we normalized contributions from each resonance in terms of relative intensity to the strongest one, which is taken as 100% (see last two columns in Tables II and III). Some resonances contribute through two different spin channels: $s = 0$ and $s = 1$. In those cases, the contributions from two spin channels can be either with the same sign or with the opposite sign, depending on unknown phases of amplitudes of partial widths and weak matrix elements (see, for example, resonance at 3.062 MeV in Table II). As can be seen from these tables, different resonances contribute essentially differently to the value of PV violating effects (this is also correct for PC asymmetries). Moreover, different sets of resonance parameters can change weights of the resonances for a particular asymmetry. For example, the lowest 0^- -resonance contribution to the asymmetry α_{PV} appears to be 3% using the parameters of Table II, while it would be the dominant one using the parameters of Table III. This is related to the fact that for the set of Table II the contribution of the 0^- -resonance to the α_{PV} is suppressed by a factor of about 40 owing to destructive interference between PC and PV amplitudes. Therefore, the readability of this method can be essentially improved by increasing the accuracy in measurements of parameters of the most “important” resonances.

It should be noted that the estimated value of the PV asymmetry α_{PV} at thermal energy [see Eqs. (10) and (11)] is surprisingly in very good agreement with exact calculations for zero-energy neutrons [13]. This could be considered as an additional argument for the reliability of the suggested resonance approach. Also, matching the estimated value of the observable parameter with exact calculations at low energy gives us the opportunity to predict PV effects in a wide range of neutron energies.

ACKNOWLEDGMENTS

I am grateful J. D. Bowman for fruitful discussions. This work was supported by DOE Grants No. DE-FG02-09ER41621.

- [1] S.-L. Zhu, C. M. Maekawa, B. R. Holstein, M. J. Ramsey-Musolf, and U. van Kolck, *Nucl. Phys. A* **748**, 435 (2005).
 [2] B. Holstein, in *Proceedings of the International Workshop on Theoretical Problems in Fundamental Neutron Physics, October 14–15, 2005, Columbia, SC* [<http://www.physics.sc.edu/TPFNP/Talks/Program.html>].

- [3] B. Desplanque, *Proceedings of the International Workshop on Theoretical Problems in Fundamental Neutron Physics, October 14–15, 2005, Columbia, SC* [<http://www.physics.sc.edu/TPFNP/Talks/Program.html>].
 [4] M. J. Ramsey-Musolf and S. A. Page, *Annu. Rev. Nucl. Part. Sci.* **56**, 1 (2006).

- [5] B. Desplanques, J. F. Donoghue, and B. R. Holstein, *Ann. Phys.* **124**, 449 (1980).
- [6] V. E. Bunakov and V. P. Gudkov, *Nucl. Phys. A* **401**, 93 (1983).
- [7] C. Mahaux and H. A. Weidenmuller, *Shell-model Approach to Nuclear Reactions* (North-Holland, Amsterdam/London, 1969).
- [8] D. R. Tilley, H. R. Weller, and G. M. Hale, *Nucl. Phys. A* **541**, 1 (1992).
- [9] S. F. Mughabghab, *Atlas of Neutron Resonances: Resonance Parameters and Thermal Cross Sections: Z = 1–100*, 5th ed. (Elsevier Science, 2006).
- [10] S. G. Kadmsky, V. P. Markushev, and V. I. Furman, *Sov. J. Nucl. Phys.* **37**, 345 (1983).
- [11] V. E. Bunakov, V. P. Gudkov, S. G. Kadmsky, I. A. Lomachenkov, and V. I. Furman, *Sov. J. Nucl. Phys.* **49**, 613 (1989).
- [12] M. B. Johnson, J. D. Bowman, and S. H. Yoo, *Phys. Rev. Lett.* **67**, 310 (1991).
- [13] M. Viviani, R. Schiavilla, L. Girlanda, A. Kievsky, and L. E. Marcucci, *Phys. Rev. C* **82**, 044001 (2010).