

4-3-2019

## Impact of Medicaid Coverage Expansion Under the Affordable Care Act on Mammography and Pap Tests Utilization Among Low-Income Women

Abeer G. Alharbi

M. Mahmud Khan

Ronnie Horner

Heather M. Brandt  
hmbrand@mailbox.sc.edu

Cole Garrett Chapman  
chapmac8@mailbox.sc.edu

Follow this and additional works at: [https://scholarcommons.sc.edu/sph\\_health\\_promotion\\_education\\_behavior\\_facpub](https://scholarcommons.sc.edu/sph_health_promotion_education_behavior_facpub)



Part of the [Public Health Education and Promotion Commons](#)

---

### Publication Info

Published in *Plos One*, Volume 14, Issue 4, 2019.

© 2019 Alharbi et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

This Article is brought to you by the Health Promotion, Education, and Behavior at Scholar Commons. It has been accepted for inclusion in Faculty Publications by an authorized administrator of Scholar Commons. For more information, please contact [digres@mailbox.sc.edu](mailto:digres@mailbox.sc.edu).


## RESEARCH ARTICLE

# Impact of Medicaid coverage expansion under the Affordable Care Act on mammography and pap tests utilization among low-income women

Abeer G. Alharbi<sup>1\*</sup>, M. Mahmud Khan<sup>1</sup>, Ronnie Horner<sup>1</sup>, Heather Brandt<sup>2</sup>, Cole Chapman<sup>1</sup>

**1** Health Services Policy and Management Department, Arnold School of Public Health, University of South Carolina, Columbia, South Carolina, United States of America, **2** Health Promotion, Education, and Behavior Department, Arnold School of Public Health, University of South Carolina, Columbia, South Carolina, United States of America

\* [abeer.g.alharbi@gmail.com](mailto:abeer.g.alharbi@gmail.com)



## OPEN ACCESS

**Citation:** Alharbi AG, Khan MM, Horner R, Brandt H, Chapman C (2019) Impact of Medicaid coverage expansion under the Affordable Care Act on mammography and pap tests utilization among low-income women. PLoS ONE 14(4): e0214886. <https://doi.org/10.1371/journal.pone.0214886>

**Editor:** Eugenio Paci, Centro per lo Studio e la Prevenzione Oncologica, ITALY

**Received:** January 16, 2019

**Accepted:** March 21, 2019

**Published:** April 3, 2019

**Copyright:** © 2019 Alharbi et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** Data cannot be shared publicly because only researchers and users with approved research projects can access restricted data files that have not been publicly released for reasons of confidentiality at the AHRQ Data Center in Rockville, Maryland. I have an approved research project and I have access to the restricted data. Upon your request, I can add your name to my project so you would have access to the data. Or you can ask for access directly from AHRQ. For more information contact AHRQ Data Center Coordinator: Ray F. Kuntz CFACT Data

## Abstract

### Introduction

The Affordable Care Act (ACA) expanded the coverage of Medicaid to include entire population with income below 138% of federal poverty line. It remains unclear whether this policy change has improved access to and utilization of health care, particularly use of mammography and Pap tests among poor women.

### Methods

We used a difference-in-difference (DID) design to estimate the impact of Medicaid expansion on mammography and Pap tests utilization among low-income women. Expansion states are the treatment group and non-expansion states are the control group. The years 2012–13 are the pre-expansion period and 2015–16 are the post-expansion period for the purpose of estimating the DID parameters.

### Results

The difference-in-difference estimate show that likelihood of utilizing mammograms did not change significantly among low-income women after the implementation of Medicaid expansion (DID coefficient -0.0476 with t-statistics at -1.26), Pap test decreased (coefficient -0.0615, t-statistics -2.76), and Medicaid enrollment has increased significantly among low-income women living in expansion states (coefficient 0.0889 with t-value of 3.68).

### Conclusion

Expansion of Medicaid was associated with increased Medicaid enrollment but did not yield near-term improvement in use of mammography and Pap tests among low-income women. Factors beyond health insurance coverage may be important in determining the likelihood of

Center Coordinator Contractor 5600 Fishers Lane  
MAILSTOP 07W41A Rockville, MD 20857 Room  
06N160 301-427-1654 E-mail: [ray.kuntz@ahrq.hhs.gov](mailto:ray.kuntz@ahrq.hhs.gov)

**Funding:** The author(s) received no specific funding for this work.

**Competing interests:** The authors have declared that no competing interests exist.

obtaining these screenings. Policy makers should try to identify other barriers to cancer screenings among low-income women in the USA.

## Introduction

Breast cancer is the most commonly diagnosed cancer among American women, and the second most common cause of death from cancer besides lung cancer [1,2]. Cervical cancer incidence rates declined by half between 1975 and 2014 due to the widespread uptake of the Pap test, but declines have slowed down in recent years [1,2]. Evidence show that women who appropriately screen for breast and cervical cancer are likely to receive more timely diagnosis and treatment [3–10] and yet, rates of mammography and Pap test screenings remained suboptimal in the United States [11]. Low-income women utilize less screenings than middle or high income women. In 2015, 54.9% of low-income women received mammography while 60% received Pap test [11]. Goals of Healthy People 2020 include increasing the proportion of women who get mammograms to 81%, and Pap tests to 93%, based on the most recent guidelines [12]. There are several possible reasons for the suboptimal screening rates, among which lack of health insurance coverage is considered an important one. There is evidence that health insurance is associated with uptake of mammogram and Pap test use [13–17].

The Patient Protection and Affordable Care Act of 2010 (ACA) expanded the coverage of Medicaid to include the entire population aged 18–64 with income below 138% of the federal poverty line [18]. Since uninsured adults were more likely to be low-income, Medicaid expansion has the potential for improving access to health care among this poor segment of the population [19–21]. One important role of health insurance is reducing the cost of receiving preventive care and in turn, reduce costly events from poorly or unmanaged chronic conditions. Under ACA, participation of States in Medicaid expansion became optional after a supreme court ruling in 2012 [22] but many states decided to participate in Medicaid expansion immediately after the policy change and by September 2015, majority of the states have expanded Medicaid. As of September 2018, 34 states have adopted Medicaid expansion, 3 states are considering expanding, and 14 states did not expand. [S1 Table](#) lists the states with Medicaid expansion status as of September 2018 [23].

Previous evidence gave mixed results regarding the impact of Medicaid expansion on utilization of mammography and Pap tests [24–29]. Expansion experiences from pre-ACA anticipated positive effect on screening rates [26–28], while post-ACA either did not find significant impact [24,25] or found a favorable results when analyzing utilization in a community health center [29]. In any case, these studies had a short follow-up time after the introduction of ACA which perhaps made it difficult to detect significant change. Also, the individuals' lack of knowledge/awareness of preventive care benefits provided by the ACA [30] may explain the limited impact of the coverage expansion on their use of these services.

Since no conclusive evidence is available, this study made an attempt to understand the effect of Medicaid expansion on probability of obtaining screening tests like mammography and Pap tests among low-income women. Since the study is using nationally representative data set, the results would indicate the effects of policy change for the country as a whole. Nationally representative data will also allow identification of factor affecting utilization rates.

## Materials and methods

### Data source

Data for this study was obtained from the Medical Expenditure Panel Survey—Household Component MEPS-HC [31]. The MEPS is a set of large-scale surveys which collects data from

a sample of families and individuals in selected communities across the United States, drawn from a nationally representative subsample of households. The MEPS contains the data on health care utilization, health insurance status, and coverage source that are required to answer the research question of the study. The combined average response rate for the years 2012–2016 was 50.7% [32]. The study was reviewed and approved by the Office of Research Compliance, an administrative office that supports the University of South Carolina Institutional Review Board (USC IRB). All data used was fully anonymized before it was accessed.

## Sample

From the 2012–16 MEPS datasets, the sample extracted consists of nonelderly low-income women living in the U.S. The sample size was 6,427 in expansion states (3,459 pre-ACA and 2,968 post-ACA) and 6,831 in non-expansion states (3,729 pre-ACA and 3,102 post-ACA). Women living in states that already provided Medicaid or similar coverage to low-income adults before ACA's Medicaid expansion in 2014 were excluded from the analysis (District of Columbia, Delaware, Massachusetts, New York, and Vermont). Women aged 65 years or older were excluded because they are eligible for Medicare. Women belonging to low-income households, as defined by the ACA, were selected for the analysis as this group is eligible for participation in Medicaid after the policy change, if they were not enrolled in Medicaid at the time of expansion. In accordance with screening guidelines, the mammography cohort will include women aged 40–64 and the Pap test cohort will include women aged 21–64. Women with concurrent or past diagnoses with breast or cervical cancer were excluded from the analysis to focus on utilization of screening services for preventive or early diagnosis purposes.

## Design

This study used a difference-in-difference (DID) design in a regression framework. This analytic design tests a comparison of the change in trends of outcomes before and after Medicaid expansion across expansion states vs non-expansion states, controlling for other covariates representing risk attitudes and preference structure. The treatment group includes women living in Medicaid expansion states and control group includes women living in non-expansion states. Only the states that expanded Medicaid between January 2014 and January 2016 were included in the treatment group (S1 Table). The states that already provided Medicaid or similar coverage to low-income adults before 2014 were excluded (District of Columbia, Delaware, Massachusetts, New York, and Vermont) (S1 Table). For estimating the DID parameters, pre-ACA period is defined as the years 2012–13 and the post period is defined as the years 2015–16.

The following multivariate linear regression was estimated to find the effect of the policy change on the outcome variables, the likelihood of receiving mammography and pap smears:

$$Y_{ist} = \beta_0 + \beta_1 Treatment + \beta_2 Post + \beta_3(Treatment * Post) + \beta_x Covariates + \epsilon$$

Where “ $Y_{ist}$ ” represents outcome for individual “ $i$ ” living in state “ $s$ ” at time “ $t$ ”.  $\beta_0$  is the baseline average. The term “Treatment” is a dummy variable equal to 1 if the individual resides in a treatment group (expansion state).  $\beta_1$  is the difference between the two groups pre-intervention. The term “Post” is a dummy variable equal to 1 if the time is after the Medicaid expansion.  $\beta_2$  is the time trend in control group. The term “Treatment\*Post” is an interaction term of intervention and time,  $\beta_3$  represents the difference-in-differences estimator capturing the effect of Medicaid expansion. Covariates are added to the model to control for preference structure and risk attitudes.

This regression model, in theory, will be able to indicate the effect of treatment if the intervention and control groups are identical at the baseline or show similar pattern of change over

the years. In the real world, the intervention and control groups in pre-intervention period are never identical and therefore differences between the groups need to be explicitly considered and incorporated in the analysis. The effect of program change can be estimated if the assumption of similar pattern of changes over the years in pre-intervention years may be assumed in post-intervention periods as well. This is known as the “parallel assumption” in DID analysis. Since the parallel assumption must hold for an unbiased DID estimator, we can test the parallel movements or trend in the outcomes prior to policy change in treatment and control groups over a number of years. To assess the validity of this assumption, we regressed each outcome for the years 2005 to 2013 on variables indicating years, state expansion status and an interaction term of year and state expansion status. If the coefficient of the interaction term is not statistically different from zero, it implies that the rate of change of the dependent variables is not different between the intervention and control areas confirming the parallel movement of the outcome over the years prior to the implementation of the intervention.

### Outcome

The outcomes for this study are the self-reported receipt of mammogram, Pap test, and Medicaid enrollment status. For the preventive services, respondents were asked “About how long has it been since you had this mammogram/Pap test?” with possible responses being “within past year,” “within past 2 years,” etc. In accordance with screening guidelines, a dummy variable was created for mammogram utilization equal to 1 if the test was taken within 1 to 2 years, and a dummy variable for pap test utilization equals to 1 if the test was taken within 1 to 3 years.

### Covariates

We controlled for variables that we believe may modify the preference structure and risk attitude of women in the sample. According to the Demand Theory, demand for health services is a function of prices of the services, household income, preference structure, and risk-attitude. We chose covariates that may modify the preference structure and risk attitude, making individuals more risk averse and therefore more likely to undergo screening tests. The covariates chosen are: age, race, marital status, education, health insurance status, comorbidity, physical activity, smoking status, and metropolitan area.

### Statistical analysis

First, univariate analysis was done to produce baseline descriptive statistics of the low-income women living in treatment (expansion states) and control groups (non-expansion states). Second, we tested the parallel trends assumption across expansion and non-expansion states by regressing each outcome for the years 2005 to 2013 on variables indicating years, state expansion status and an interaction term of year and state expansion status. Third, a difference-in-differences regression model was estimated by linear ordinary least squares. A linear model was chosen to allow a direct interpretation of the coefficients and avoid interpretive issues inherent to interaction terms in nonlinear models [33,34]. The key parameter of interest from the DID model was the parameter associated with the interaction between treatment and time. This parameter represents the estimated difference in outcome rates between pre- and post-policy change, across states that were and were not affected by the policy change. Differences were considered statistically significant if P-value of t-statistics  $<0.05$ . Finally, a sub-group analyses was done using linear regression and univariate analysis to explain the effect of different demographics, socioeconomics, and geographic determinants on screening use. All analyses were carried out using STATA software version 14 (2015; Stata 14.0 Statistical Software,

**Table 1. Baseline characteristics of low-income women aged 18–64, living in expansion and non-expansion states, pre-ACA (2012–13), MEPS dataset.**

Characteristic	Non-expansion states N = 3,729	Expansion states N = 3,459	P value
Age	37 (mean)	37 (mean)	0.0839
Race			0.000
White	57.39	67.07	
Black	37.38	23.94	
Other minorities	5.23	8.99	
Education			0.000
Some school	34.51	38.90	
High school	34.54	31.17	
College	30.95	29.94	
Health insurance			0.000
Private	18.10	16.48	
Public	41.67	53.22	
Uninsured	40.23	30.30	
Metropolitan area			0.000
Metro	82.38	87.80	
Non-metro	17.62	12.20	

<https://doi.org/10.1371/journal.pone.0214886.t001>

College Station, TX, USA). The analyses accounted for probability weighting in the MEPS [35,36] to obtain national estimates of effects of the policy change.

## Results

Table 1 shows the baseline characteristics of the nonelderly low-income women living in expansion and non-expansion states. Majority of the low-income women were white in both expansion and non-expansion states, however, more black women lived in non-expansion states (37.38%) compared to expansion states (23.94%). In both treatment and control groups, majority of low-income women did not have a college degree (Table 1). In expansion and non-expansion states, majority of low-income women had public health insurance, however, more women had public health insurance in expansion states (53.22%) as compared to non-expansion states (41.67%). Majority of the low-income women lived in metropolitan areas in both treatment and control groups (Table 1). Women in treatment and control groups had a similar average age (37). Therefore, states deciding to expand Medicaid were different from the states deciding not to expand in terms of percent of low income population not white, level of coverage of public insurance program and percent of poor women living in metro areas.

Fig 1 shows trends in mammogram and Pap tests rates across expansion and non-expansion state for the years 2005 to 2013. Results from the regression that tested the parallel assumption of the time trend of outcome variable show that the slope of the trend functions were similar for these two groups of states prior to the implementation of the ACA policy on Medicaid expansion (S2 Table).

Results from the univariate analysis that examined screening rates among women living in expansion states post-ACA by different sub-groups can be found in S3 Table. The results show majority of low-income women who used mammograms and Pap tests were high-income (76.20%, 85.73%) high-education attainment (72.93%, 83.88%), Black (73.61%, 87.91%), with private insurance (74.20%, 84.13%), living in metropolitan areas (71.36%, 83.13%), and reported having a usual source of care (74.52%, 83.92%), for mammograms and Pap tests respectively.




Fig 1. Trends in mammogram and Pap test uptakes in expansion and non-expansion states among low-income women, MEPS 2005–13.

<https://doi.org/10.1371/journal.pone.0214886.g001>

**Table 2. Results from the difference-in-differences adjusted regression model, nonelderly low-income women (2012–16), MEPS dataset.**

Outcome	Expansion states		Non-expansion states		Difference-in-differences
	Pre-ACA rate	Post-ACA rate	Pre-ACA rate	Post-ACA rate	
Mammogram	62.66%	64.69%	58.87%	61.77%	-0.0476 (-1.26)
Pap tests	81.90%	80.19%	78.80%	79.36%	-0.0615** (-2.76)
Medicaid enrollment	38.10%	52.31%	21.12%	25.03%	0.0889*** (3.68)

t statistics in parentheses

\* p<0.05

\*\* p<0.01

\*\*\* p<0.001

<https://doi.org/10.1371/journal.pone.0214886.t002>

Table 2 reports the results from the difference-in-difference adjusted regression model. The DID estimates indicate that the probability of enrolling to Medicaid has increased significantly among the low-income women after the implementation of Medicaid expansion (estimated coefficient 0.0889 with t-value 3.68). The DID estimates indicate that for the Medicaid expansion states the probability of utilizing mammograms did not change significantly (estimated coefficient -0.0476 with t-value -1.26) while for the Pap tests the probability of utilizing the test has decreased significantly among low-income women after the implementation of Medicaid expansion compared to non-expansion states (estimated coefficient -0.0615, t-value -2.76).

Table 3 shows the results on the likelihood of receiving mammograms and Pap tests among low-income women using a number of possible determinants of utilization of the screening tests. The sub-group analysis shows that poor women with higher age were more likely to receive mammograms (estimated coefficient 0.0102, t-value 4.16) and less likely to receive Pap tests (estimated coefficient -0.174, t-value -5.08). Black women were more likely to receive mammograms (estimated coefficient 0.0812, t-value 3.87) and Pap tests (estimated coefficient 0.0686, t-value 5.75) as compared to white women. The table also indicates that women from other minority population groups were less likely to receive Pap tests as compared to white women (estimated coefficient -0.0646, t-value -2.90). Women with a college degree were more likely to receive mammograms (estimated coefficient 0.0605, t-value 2.56) than those who had less than high school education. Women who were divorced were less likely to receive mammograms (coefficient -0.0875, t-value -3.47) and Pap tests (coefficient -0.0385, t-value -2.00) compared to married women. Women with public health insurance or no insurance (uninsured) were less likely than those with private insurance to receive mammograms (estimated coefficients -0.0654 and -0.250). No statistically significant difference was detected for Pap test use between women with public and private insurance (estimated coefficient -0.00391, t-value -0.26) although uninsured women were less likely to receive and Pap tests (estimated coefficient -0.110, t-value -6.61).

Women living in non-metropolitan areas were less likely than those in metropolitan areas to receive Pap tests (estimated coefficient -0.0615, t-value -3.41) but no difference was detected for mammogram use (estimated coefficient -0.0489, t-value -1.62). Women who reported not having a usual source of care were less likely to receive mammograms (estimated coefficient -0.141, t-value -4.03) and Pap tests (estimated coefficient -0.0562, t-value -3.02) compared to those who have a usual source of care. Non-smokers were more likely to receive mammograms



**Table 3. Likelihoods of receiving mammograms and Pap tests using a number of determinants, results from adjusted linear regression model (2012–16), MEPS dataset.**

Determinants	Mammogram	Pap test
Age (continuous)	0.0102*** (4.16)	0.000137 (0.11)
21–39	NA	Reference group
40–49	Reference group	-0.0789*** (-3.59)
50–64	-0.00818 (-0.23)	-0.174*** (-5.08)
Black(Reference group: White)	0.0812*** (3.87)	0.0686*** (5.75)
Other minorities(Reference group: White)	-0.0588 (-1.59)	-0.0646** (-2.90)
High school(Reference group: Some school)	0.0419 (1.82)	-0.00437 (-0.33)
College(Reference group: Some school)	0.0605* (2.56)	-0.0120 (-0.85)
Widowed(Reference group: Married)	-0.103* (-2.56)	-0.0883* (-2.47)
Divorced(Reference group: Married)	-0.0875*** (-3.47)	-0.0385* (-2.00)
Separated from spouse(Reference group: Married)	-0.0799* (-2.37)	-0.00593 (-0.27)
Never Married(Reference group: Married)	-0.0749** (-2.86)	-0.0277* (-2.03)
Public insurance(Reference group: Private insurance)	-0.0654* (-2.55)	-0.00391 (-0.26)
Uninsured(Reference group: Private insurance)	-0.250*** (-8.97)	-0.110*** (-6.61)
Non-metropolitan area(Reference group: Metropolitan area)	-0.0489 (-1.62)	-0.0615*** (-3.41)
Usual source of care not available(Reference group: Usual source of care not available)	-0.141*** (-4.03)	-0.0562** (-3.02)
Do not exercise regularly(Reference group: Exercise regularly)	-0.00171 (-0.09)	-0.0102 (-0.92)
Non-smoker(Reference group: Smoker)	0.0706** (3.15)	0.0446*** (3.30)
1 chronic disease(Reference group: No chronic disease)	0.0479 (1.68)	0.0223 (1.63)
+2 chronic diseases(Reference group: No chronic disease)	0.133*** (5.22)	0.0251 (1.58)

t statistics in parentheses

\* p<0.05

\*\* p<0.01

\*\*\* p<0.001

<https://doi.org/10.1371/journal.pone.0214886.t003>

(estimated coefficient 0.0706, t-value 3.15) and Pap tests (estimated coefficient 0.0446, t-value 3.30) as compared to smokers. Women with two or more chronic diseases were more likely to receive mammograms (estimated coefficient 0.133, t-value 5.22) but no differences were detected for Pap test use (estimated coefficient 0.0251, t-value 1.58).

## Discussion

The affordable care act (ACA) expanded Medicaid eligibility coverage to the entire low-income population in order to improve access and utilization among this disadvantage section of the population. In the years before the ACA, rates of mammograms and Pap tests showed declining trends among women and more so among poor women [11]. This study examined the impact of expanding health coverage through Medicaid on the rates of mammograms and Pap tests among poor women. The difference-in-difference (DID) estimates indicate that Medicaid enrollment has increased significantly among low-income women after the implementation of the Medicaid expansion (Table 2). This is a proximate measure of success of ACA in terms of providing coverage to poor women through Medicaid. Other studies also found increased Medicaid enrollment in expansion states compared to non-expansion states [37]. However, the increase in Medicaid enrollment among low-income women did not translate into increased rates of mammograms or Pap test utilization compared to poor women in non-expansion states. Other studies also found little impact of Medicaid expansion on mammography and Pap tests rates [37–39].

Although the difference-in-differences estimate did not show increase in mammograms and Pap tests rates, low-income women living in expansion states used more screenings than their counterparts in non-expansion states (Table 2). Historically, mortality rates of breast and cervical cancer were lower in the states that elected to expand Medicaid compared to those who elected not to expand [40]. We compared cancer burden in expansion vs non-expansion states in pre and post-ACA using data from the National Cancer Institute (NCI) and found that women in expansion states had lower mortality rates compared to women in non-expansion states (breast cancer: 20.13 vs 20.50; and cervical cancer: 1.97 vs 2.41) per 100,000 resident [40]. A previous study found that Southeastern states without Medicaid expansion tended to have higher cancer and lower screening rates and therefore disparities in cancer screening that already disfavor states with high cancer rates may widen in states that have chosen not to expand Medicaid [41].

A number of possible explanations can be advanced for this lack of improvements in mammograms and Pap tests rates among the low-income women despite gaining insurance coverage through Medicaid. First, poor knowledge/awareness about the availability of preventive benefits through ACA may result in low access to these services. There is evidence showing that newly insured individuals are not aware of the preventive services benefits of the ACA and thus do not use them [30]. Another study found that only 36.4% of adults reported knowing that the ACA requires coverage of certain preventive services without cost sharing [42]. Therefore, strategies aimed at improving public awareness about the availability of preventive services may be a vital tool in improving screening rates.

Second, in the U.S. health system, acute care takes priority over prevention. Studies have found that time constraints limit the ability of primary physicians to comply with preventive services recommendations [43]. A study has found that in states with higher Medicaid payments for office visits, Medicaid beneficiaries were more likely to be screened for breast and cervical cancer [44]. This indicate that increasing screenings among vulnerable population may be achieved through enhanced Medicaid reimbursements for physician consultations.

Third, our analysis show that low-income women with private insurance were more likely to receive mammograms than those with public insurance (Table 3). Also, in expansion states, the proportion of low-income women with private insurance used more mammograms and Pap tests than those with public insurance (S3 Table). Another study found that women with employer-based insurance/Medicare were more likely to get breast and cervical cancer screenings [45]. According to a survey conducted by the National Center for Health Statistics, only

two out of three primary care physicians surveyed in 2011 were willing to accept new Medicaid patients [46]. Our analysis showed that women who reported having a usual source of care were more likely to receive mammograms and Pap tests than those without a usual source of care (Table 3). In expansion states, the proportion of low-income women who have a usual source of care used more mammograms and Pap tests than their counterparts in non-expansion states (S3 Table). Therefore, insurance-type and having a usual source of care appear to be more important in improving access to care and receiving these screenings.

A number of studies did find that Medicaid expansion was effective in improving utilization rates of certain preventive services such as glucose testing, cholesterol testing, and annual check-up, but not for cancer screenings [24,47]. The U.S. Preventive Services Task Force (USPSTF) guidelines for mammograms and Pap tests were updated around the time ACA was being implemented. The guidelines for mammograms were updated in 2009 to recommend mammograms for women aged 50–75 years every 2 years from the previous guidelines that recommended screening every 1–2 years for women aged 40 or older [48]. The guidelines for cervical cancer screenings was updated in 2012 to recommend the Pap test for women aged 21–65 every three years from the previous guidelines that recommended screening annually for women who are sexually active. This may explain the overall decline in cancer screenings in recent years. However, the effect of revised guidelines should be universal applicable to both expansion and non-expansion states and therefore may not explain the lack of effect on mammograms and Pap tests in expansion states in our difference-in-difference design.

It is also possible that the results from the difference-in-difference model are biased or washed-out because of the possibility that a significant number of poor women received screenings through other national programs such as the National Breast and Cervical Cancer Early Detection Program (NBCCEDP) through the Centers for Disease Control and Prevention (CDC). The program was established in 1990 to provide free and/or reduced cost mammograms and Pap tests to women with limited incomes and those who lack health insurance. Although number of women receiving those services through the NBCCEDP has decreased in 2015–16 compared to 2013 [49], low-income women still benefit from this program which may have negated the impact of ACA on mammograms and Pap tests rates in our difference-differences analysis. Between 2012 and 2017 the NBCCEDP program provided 740,108 Pap tests and 902,751 mammograms to low-income women [49]. During 2012 to 2017, this program provided 441,206 Pap tests (452 screenings per 100,000 woman) and 498,659 mammograms (511 screenings per 100,000 woman) in expansion states. During this same period, 398 and 573 per 100,000 woman received Pap tests and mammography respectively in non-expansion states through this program [49]. Mammograms provided to women in non-expansion states was about 12% higher than the rate in expansion states and the opposite is true for Pap tests (12% lower in non-expansion states). Such differences in coverage rates of an external program can potentially bias the estimation of the effect of ACA on mammograms and Pap tests uptakes.

Our empirical analysis showed that screenings occurred at much higher rates among the high-income high-educated women while the low-income women had the lowest utilization rates (S3 Table). Before the ACA, a study found that the low-income women were less likely to receive possibly lifesaving recommended cancer screenings [50]. A post-ACA study examined the impact of Medicaid expansion on disparities in cancer screenings and found that large gaps remain in access, particularly for low-income adults [51]. Our analysis showed that disparities in terms of using mammograms and Pap tests remained and may have actually become worse. This possibly implies that other factors beyond insurance coverage (in this case, provided through Medicaid) should be examined in order to better understand the reasons for the persistence of socioeconomic disparities.

Finally, our sub-group analyses helped shed light on the likelihood of receiving mammograms and Pap tests among low-income women using a number of possible determinants of utilization. As women get older they are more likely to receive mammograms and less likely to receive pap tests (Table 3). This is expected as evidence suggests that the benefits from mammograms are more evident for older women while benefits from Pap tests are more evident for younger women [52,53]. Low-income women living in metropolitan areas were more likely to receive mammograms and Pap tests in both expansion and non-expansion states (Table 3). This is expected as access to care is better in metropolitan areas than in non-metro areas. Black women were more likely than white women to receive a Pap test (Table 3) and in expansion states, the proportion of black women receiving mammograms and Pap tests were higher than white women (S3 Table). African American women in the U.S. are more likely to be diagnosed and die from breast and cervical cancer than white women, which may explain the increased use of the screenings [54].

### Limitations

We acknowledge some important limitations of this study. First, information about outcomes relied on self-reported survey responses which might be subject to recall error. However, the MEPS follow up with health providers to reduce the reporting bias but some errors may still remain, especially for procedures and tests requiring longer recall time frame. Second, the data used in the analysis are cross-sectional and comparison of cross-sectional data at different years is not same as observing changes in the outcomes for a cohort with the implementation of ACA. The study design made an attempt to tease-out the effect of policy changes through DID and in most cases DID approach can identify the effect of policy change even when the starting characteristics of the control and intervention groups are significantly different. Third, this study examined the initial 3-year period after the ACA Medicaid expansion provision and a longer time frame may be needed to be able to see the effects of policy changes on outcomes. Finally, there were changes in the USPSTF guidelines for breast cancer and cervical screening that occurred around the same time as the ACA provisions but it should not affect the results of DID. One of the important sources of bias that could not be corrected for in the empirical analysis is the provision of these screenings to poor women by a national program free of charge. If this national program in post-policy change years provided more emphasis on offering screenings in non-expansion states, it can potentially offset any positive effects of Medicaid expansion in DID modeling. In any case, this lack of relative improvements in cancer screenings in the Medicaid expansion states (compared to non-expansion states) is perplexing and would require supplementing the national data with the effects of alternative programs and other structural differences between these two groups of states.

### Conclusion

Our study shows that expansion of Medicaid under the ACA was associated with increased Medicaid enrollment but did not yield near-term improvements in the use of mammography and Pap tests among low-income women. Although the difference-in-differences did not show improvements in mammograms and Pap tests due to Medicaid expansion under ACA, low-income women living in expansion states used higher level of screenings than their counterparts in non-expansion states. Since Medicaid expansion did not affect these screening tests, policy makers need to examine other factors that may act as barriers in improving access and utilization. Some possible explanations for this lack of impact of the Medicaid expansion on mammograms and Pap tests are presented in the discussion section but we have no concrete evidence to conclusively say which factors have affected access to screening tests adversely in

the expansion states compared to non-expansion states. It is also possible that a longer time-frame will be needed for a change to be manifested itself rather than the three-year time frame used here. Future research on provider availability and characteristics, insurance types, and geographical variations is warranted for a better understanding of the use of cancer screening procedures by the poor women in the USA.

## Supporting information

**S1 Table. State Medicaid expansion status as of September 2018 with the start date of expansion in the states (Source: Kaiser foundation website).**

(DOCX)

**S2 Table. Test of parallel trend assumption in expansion and non-expansion states prior to introduction of ACA Medicaid expansion.** Interaction term of year and state expansion status.

(DOCX)

**S3 Table. Rates of mammograms and Pap tests use and change in the rates in expansion states in post-ACA years by individual characteristics: Results from univariate analysis (2012–16) of MEPS data.**

(DOCX)

## Author Contributions

**Conceptualization:** Abeer G. Alharbi, M. Mahmud Khan, Ronnie Horner, Heather Brandt, Cole Chapman.

**Data curation:** Abeer G. Alharbi.

**Formal analysis:** Abeer G. Alharbi.

**Methodology:** Abeer G. Alharbi, M. Mahmud Khan, Ronnie Horner.

**Supervision:** M. Mahmud Khan.

**Validation:** M. Mahmud Khan.

**Writing – original draft:** Abeer G. Alharbi.

**Writing – review & editing:** M. Mahmud Khan, Ronnie Horner, Heather Brandt, Cole Chapman.

## References

1. American Cancer Society. Cancer Facts & Figures 2018 [Internet]. 2018 [cited 2018 Feb 19]. Available from: <https://www.cancer.org/content/dam/cancer-org/research/cancer-facts-and-statistics/annual-cancer-facts-and-figures/2018/cancer-facts-and-figures-2018.pdf>
2. Centers for Disease Control and Prevention CDC. United States Cancer Statistics [Internet]. [cited 2018 Nov 10]. Available from: <https://gis.cdc.gov/grasp/USCS/DataViz.html>
3. American Cancer Society. Cancer Facts & Figures 2018 [Internet]. 2018 [cited 2018 Feb 18]. Available from: <https://www.cancer.org/content/dam/cancer-org/research/cancer-facts-and-statistics/annual-cancer-facts-and-figures/2018/cancer-facts-and-figures-2018.pdf>
4. American Cancer Society. Key Statistics for Cervical Cancer [Internet]. 2018 [cited 2018 Feb 18]. Available from: <https://www.cancer.org/cancer/cervical-cancer/about/key-statistics.html>
5. Burd EM. Human Papillomavirus and Cervical Cancer. *Clin Microbiol Rev.* 2003 Jan; 16(1):1–17. <https://doi.org/10.1128/CMR.16.1.1-17.2003> PMID: 12525422
6. Institute of Medicine (US) and National Research Council (US) National Cancer Policy Board. Fulfilling the Potential of Cancer Prevention and Early Detection [Internet]. Curry SJ, Byers T, Hewitt M, editors.

- Washington (DC): National Academies Press (US); 2003. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK223926/>
7. Maciosek MV, Coffield AB, Flottemesch TJ, Edwards NM, Solberg LI. Greater Use Of Preventive Services In U.S. Health Care Could Save Lives At Little Or No Cost. *Health Aff.* 2010 Sep 1; 29(9):1656–60.
  8. Coldman A, Phillips N, Wilson C, Decker K, Chiarelli AM, Brisson J, et al. Pan-Canadian Study of Mammography Screening and Mortality from Breast Cancer. *J Natl Cancer Inst* [Internet]. 2014 Nov 1 [cited 2017 Jul 20]; 106(11). Available from: <https://academic-oup-com.pallas2.tcl.sc.edu/jnci/article/106/11/dju261/1496367/Pan-Canadian-Study-of-Mammography-Screening-and>
  9. Hofvind S, Ursin G, Tretli S, Sebuødegård S, Møller B. Breast cancer mortality in participants of the Norwegian Breast Cancer Screening Program. *Cancer.* 2013 Sep 1; 119(17):3106–12. <https://doi.org/10.1002/cncr.28174> PMID: 23720226
  10. Shapiro S, Strax P, Venet L. Evaluation of periodic breast cancer screening with mammography. Methodology and early observations. *JAMA.* 1966 Feb 28; 195(9):731–8. PMID: 5951878
  11. National Center for Health Statistics. Health, United States, 2016: With Chartbook on Long-term Trends in Health. Hyattsville, MD. [Internet]. 2017 [cited 2017 Jul 20]. Available from: [https://www.cdc.gov/nchs/data/16.pdf#070](https://www.cdc.gov/nchs/data/hus/16.pdf#070)
  12. Cancer | Healthy People 2020 [Internet]. [cited 2019 Feb 26]. Available from: <https://www.healthypeople.gov/2020/topics-objectives/topic/cancer/objectives>
  13. Carrasquillo O, Pati S. The role of health insurance on Pap smear and mammography utilization by immigrants living in the United States. *Preventive Medicine.* 2004 Nov 1; 39(5):943–50. <https://doi.org/10.1016/j.ypmed.2004.03.033> PMID: 15475028
  14. Sudano JJ, Baker DW. Intermittent Lack of Health Insurance Coverage and Use of Preventive Services. *Am J Public Health.* 2003 Jan 1; 93(1):130–7. PMID: 12511402
  15. Meer J, Rosen HS. Insurance and the utilization of medical services. *Social Science & Medicine.* 2004 May 1; 58(9):1623–32.
  16. Busch SH, Duchovny N. Family coverage expansions: Impact on insurance coverage and health care utilization of parents. *Journal of Health Economics.* 2005 Sep; 24(5):876–90. <https://doi.org/10.1016/j.jhealeco.2005.03.007> PMID: 15998548
  17. Rauscher GH, Allgood KL, Whitman S, Conant E. Disparities in Screening Mammography Services by Race/Ethnicity and Health Insurance. *J Womens Health (Larchmt).* 2012 Feb; 21(2):154–60.
  18. Patient Protection and Affordable Care Act [Internet]. 2010 p. 393. Available from: <http://housedocs.house.gov/energycommerce/ppacacon.pdf>
  19. Garfield R, Licata R, Feb 06 KYP, 2014. The Uninsured at the Starting Line: Findings from the 2013 Kaiser Survey of Low-Income Americans and the ACA [Internet]. The Henry J. Kaiser Family Foundation. 2014 [cited 2018 Nov 19]. Available from: <https://www.kff.org/uninsured/report/the-uninsured-at-the-starting-line-findings-from-the-2013-kaiser-survey-of-low-income-americans-and-the-aca/>
  20. Smith J, Medala C. Health Insurance Coverage in the United States: 2014. Current Population Reports [Internet]. 2015 [cited 2017 Jul 19]. Available from: <https://www.census.gov/content/dam/Census/library/publications/2015/demo/p60-253.pdf>
  21. Rowland D. Profiles in coverage: Medicaid, CHIP, and the uninsured. Presented at “On the Road to 2014: Medicaid and CHIP Eligibility and the Uninsured” [Internet]. 2011 [cited 2017 Jul 19]. Available from: [https://www.medicaid.gov/State-Resource-Center/downloads/0907\\_CB\\_1215\\_1\\_Profiles\\_in\\_Coverage\\_508.pdf](https://www.medicaid.gov/State-Resource-Center/downloads/0907_CB_1215_1_Profiles_in_Coverage_508.pdf)
  22. National Federation of Independent Business v. Sebelius 567 US \_\_\_\_ (2012) [Internet]. Justia Law. [cited 2017 Jul 30]. Available from: <https://www.supremecourt.gov/opinions/11pdf/11-393c3a2.pdf>
  23. Kaiser Family Foundation. Current Status of State Medicaid Expansion Decisions [Internet]. The Henry J. Kaiser Family Foundation. 2018 [cited 2018 Jan 22]. Available from: <https://www.kff.org/health-reform/slide/current-status-of-the-medicaid-expansion-decision/>
  24. Simon K, Soni A, Cawley J. The Impact of Health Insurance on Preventive Care and Health Behaviors: Evidence from the 2014 ACA Medicaid Expansions [Internet]. National Bureau of Economic Research; 2016 May. Available from: <http://www.nber.org/papers/w22265>
  25. Miller S, Wherry LR. Health and Access to Care during the First 2 Years of the ACA Medicaid Expansions. *N Engl J Med.* 2017 09; 376(10):947–56. <https://doi.org/10.1056/NEJMsa1612890> PMID: 28273021
  26. Sabik LM, Tarazi WW, Bradley CJ. State Medicaid Expansion Decisions and Disparities in Women’s Cancer Screening. *American Journal of Preventive Medicine.* 2015 Jan 1; 48(1):98–103. <https://doi.org/10.1016/j.amepre.2014.08.015> PMID: 25441234

27. Busch SH, Duchovny N. Family coverage expansions: Impact on insurance coverage and health care utilization of parents. *Journal of Health Economics*. 2005 Sep; 24(5):876–90. <https://doi.org/10.1016/j.jhealeco.2005.03.007> PMID: 15998548
28. Finkelstein A, Taubman S, Wright B, Bernstein M, Gruber J, Newhouse JP, et al. The Oregon Health Insurance Experiment: Evidence from the First Year\*. *Q J Econ*. 2012 Aug 1; 127(3):1057–106. PMID: 23293397
29. Cole MB, Galárraga O, Wilson IB, Wright B, Trivedi AN. At Federally Funded Health Centers, Medicaid Expansion Was Associated With Improved Quality Of Care. *Health Aff*. 2017 Jan 1; 36(1):40–8.
30. Williams JAR, Ortiz SE. Examining public knowledge and preferences for adult preventive services coverage. *PLoS One* [Internet]. 2017 Dec 20 [cited 2019 Feb 26]; 12(12). Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5738055/>
31. Medical Expenditure Panel Survey, Agency for Healthcare Research and Quality [Internet]. 2018 [cited 2018 Feb 19]. Available from: <https://meps.ahrq.gov/mepsweb/>
32. MEPS-HC response rates by panel [Internet]. Agency for Healthcare Research and Quality; 2017. Available from: [https://meps.ahrq.gov/mepsweb/survey\\_comp/hc\\_response\\_rate.jsp](https://meps.ahrq.gov/mepsweb/survey_comp/hc_response_rate.jsp)
33. Ai C, Norton EC. Interaction terms in logit and probit models. *Economics Letters*. 2003 Jul 1; 80(1):123–9.
34. Puhani PA. The treatment effect, the cross difference, and the interaction term in nonlinear “difference-in-differences” models. *Economics Letters*. 2012 Apr 1; 115(1):85–7.
35. Using Statistical Software Packages to Produce Estimates from MEPS Data Files [Internet]. [cited 2018 Sep 23]. Available from: [https://meps.ahrq.gov/survey\\_comp/hc\\_samplecodes\\_se.shtml](https://meps.ahrq.gov/survey_comp/hc_samplecodes_se.shtml)
36. Accounting for Clustering in the Analysis of MEPS Data: Frequently Asked Questions (FAQ):2.
37. Simon K, Soni A, Cawley J. The Impact of Health Insurance on Preventive Care and Health Behaviors: Evidence from the First Two Years of the ACA Medicaid Expansions. *J Policy Anal Manage*. 2017; 36(2):390–417. PMID: 28378959
38. Miller S, Wherry LR. Health and Access to Care during the First 2 Years of the ACA Medicaid Expansions. *N Engl J Med*. 2017 09; 376(10):947–56. <https://doi.org/10.1056/NEJMsa1612890> PMID: 28273021
39. Cawley J, Soni A, Simon K. Third Year of Survey Data Shows Continuing Benefits of Medicaid Expansions for Low-Income Childless Adults in the U.S. *J Gen Intern Med*. 2018 Sep; 33(9):1495–7. <https://doi.org/10.1007/s11606-018-4537-0> PMID: 29943107
40. State Cancer Profiles Historical Trends [Internet]. [cited 2018 Nov 19]. Available from: <https://statecancerprofiles.cancer.gov/historicaltrend/index.php>
41. Choi SK, Adams SA, Eberth JM, Brandt HM, Friedman DB, Tucker-Seeley RD, et al. Medicaid Coverage Expansion and Implications for Cancer Disparities. *Am J Public Health*. 2015 Oct 8; 105(S5):S706–12.
42. Lantz PM, Evans WD, Mead H, Alvarez C, Stewart L. Knowledge of and Attitudes Toward Evidence-Based Guidelines for and Against Clinical Preventive Services: Results from a National Survey. *Milbank Q*. 2016 Mar; 94(1):51–76. <https://doi.org/10.1111/1468-0009.12181> PMID: 26994709
43. Yarnall KSH, Pollak KI, Østbye T, Krause KM, Michener JL. Primary Care: Is There Enough Time for Prevention? *Am J Public Health*. 2003 Apr; 93(4):635–41. PMID: 12660210
44. Halpern MT, Romaine MA, Haber SG, Tangka FK, Sabatino SA, Howard DH. Impact of state-specific Medicaid reimbursement and eligibility policies on receipt of cancer screening. *Cancer*. 2014; 120(19):3016–24. <https://doi.org/10.1002/cncr.28704> PMID: 25154930
45. Zhao G, Okoro CA, Li J, Town M. Health Insurance Status and Clinical Cancer Screenings Among U.S. Adults. *Am J Prev Med*. 2018 Jan; 54(1):e11–9. <https://doi.org/10.1016/j.amepre.2017.08.024> PMID: 29102459
46. In 2011 nearly one-third of physicians said they would not accept new Medicaid patients, but rising fees may help.—PubMed—NCBI [Internet]. [cited 2018 Nov 20]. Available from: <https://www.ncbi.nlm.nih.gov/pallas2.tcl.sc.edu/pubmed/22869644>
47. Cawley J, Soni A, Simon K. Third Year of Survey Data Shows Continuing Benefits of Medicaid Expansions for Low-Income Childless Adults in the U.S. *J Gen Intern Med*. 2018 Sep; 33(9):1495–7. <https://doi.org/10.1007/s11606-018-4537-0> PMID: 29943107
48. USPSTF A and B Recommendations. U.S. Preventive Services Task Force [Internet]. 2017 Apr. Available from: <https://www.uspreventiveservicestaskforce.org/Page/Name/uspstf-a-and-b-recommendations/>

49. CDC—Cancer—NBCCEDP Screening Program Summaries—National Aggregate [Internet]. 2018 [cited 2018 Nov 20]. Available from: [https://www.cdc.gov/cancer/nbccedp/data/summaries/national\\_aggregate.htm](https://www.cdc.gov/cancer/nbccedp/data/summaries/national_aggregate.htm)
50. Swan J, Breen N, Graubard BI, McNeel TS, Blackman D, Tangka FK, et al. Data and trends in cancer screening in the United States. *Cancer*. 2010; 116(20):4872–4881. <https://doi.org/10.1002/cncr.25215> PMID: 20597133
51. Shartz A, Long SK, Anderson N. Access To Care And Affordability Have Improved Following Affordable Care Act Implementation; Problems Remain. *Health Aff*. 2015 Dec 16; <https://doi.org/10.1377/hlthaff.2015.0755> PMID: 26674536
52. Moss SM, Cuckle H, Evans A, Johns L, Waller M, Bobrow L, et al. Effect of mammographic screening from age 40 years on breast cancer mortality at 10 years' follow-up: a randomised controlled trial. *Lancet*. 2006 Dec 9; 368(9552):2053–60. [https://doi.org/10.1016/S0140-6736\(06\)69834-6](https://doi.org/10.1016/S0140-6736(06)69834-6) PMID: 17161727
53. Little benefit of cervical-cancer screening after age 55 with HPV-negative test [Internet]. *Journal Of Clinical Pathways*. [cited 2018 Nov 11]. Available from: <https://www.journalofclinicalpathways.com/news/little-benefit-cervical-cancer-screening-after-age-55-hpv-negative-test>
54. Viens LJ. Human Papillomavirus—Associated Cancers—United States, 2008–2012. *MMWR Morb Mortal Wkly Rep* [Internet]. 2016 [cited 2018 Nov 11];65. Available from: <https://www.cdc.gov/mmwr/volumes/65/wr/mm6526a1.htm>