University of South Carolina [Scholar Commons](https://scholarcommons.sc.edu/)

[Faculty Publications](https://scholarcommons.sc.edu/biol_facpub) **Biological Sciences**, Department of

2010

Evolution of an MHC Class La Gene Fragment in Four North American Morone Species

J. X. Liu

Bert Ely University of South Carolina - Columbia, ely@sc.edu

Follow this and additional works at: [https://scholarcommons.sc.edu/biol_facpub](https://scholarcommons.sc.edu/biol_facpub?utm_source=scholarcommons.sc.edu%2Fbiol_facpub%2F282&utm_medium=PDF&utm_campaign=PDFCoverPages)

Part of the [Biology Commons](https://network.bepress.com/hgg/discipline/41?utm_source=scholarcommons.sc.edu%2Fbiol_facpub%2F282&utm_medium=PDF&utm_campaign=PDFCoverPages)

Publication Info

Postprint version. Published in Journal of Fish Biology, Volume 76, 2010, pages 1984-1994.

This Article is brought to you by the Biological Sciences, Department of at Scholar Commons. It has been accepted for inclusion in Faculty Publications by an authorized administrator of Scholar Commons. For more information, please contact digres@mailbox.sc.edu.

Abstract

 The deep evolutionary history of MHC allelic lineages provides an opportunity to study the accumulation of mutations over a long time scale. In the present study, a sequence analysis of a fragment of an MHC class Ia gene was performed in four North American *Morone* species and 39 flanking region alleles was identified among 45 individuals. Furthermore, an analysis of the evolutionary dynamics of these alleles along with two microsatellite loci (SB83 and SB84) in this region provided evidence for gene duplications and extensive trans-species polymorphism. Although high levels of polymorphism were detected in *M. saxatilis* (striped bass), *M. americana* (white perch), and *M. mississippiensis* (yellow bass), extremely low levels of MHC diversity were detected in *M. chrysops* (white bass), suggesting the possibility of a severe population bottleneck for this species. The perfect microsatellite, SB83, showed extensive length variability among alleles. Polarity of base substitutions within repeats of SB83 was also detected. The structure of the short interrupted compound locus, SB84, also has evolved primarily by repeat duplications. Unlike SB83, however, the alleles of SB84 correlate with the microsatellite flanking region allelic lineages, indicating that SB84 has evolved more slowly than SB83. Key words: MHC, *Morone*, microsatellite, trans-species polymorphism, gene duplication, population bottleneck **Introduction** The major histocompatibility complex (MHC) of jawed vertebrates is a multigene family

involved in antigen presentation to immune system (Klein, 1986). A striking feature of MHC is

 the extensive allelic polymorphism found in many of its genes that is thought to be maintained by pathogen-driven balancing selection (Hughes & Yeager, 1998). Allelic lineages of MHC can persist over long periods of evolutionary time, even across multiple speciation events, resulting in trans-species polymorphisms (Klein *et al*., 1998a). The deep evolutionary history of the MHC allelic lineages provides an opportunity to study the historical accumulation of mutations over a long time scale.

 Microsatellites are short tandemly repeated sequence motifs (1-6 nucleotides) found throughout the genomes of many higher organisms (Tautz, 1993; Schlötterer, 2000; Ellegren, 2004). Due to their hypervariablity and ubiquitous occurrence, microsatellites are widely used in an impressive number of biological applications such as parentage analysis, gene mapping, and assessments of population structure (Schlötterer & Pemberton, 1994; Goldstein & Schlötterer, 1999). In recent years, much has been learned about the complex mutational process of microsatellites. DNA replication slippage has been described as the primary mechanism causing variation in the repeat number of microsatellites (Schlötterer & Tautz, 1992; Ellegren, 2000). Base substitution has played an essential role in microsatellite contractions, by breaking long repeat arrays into smaller units (Kruglyak *et al*., 1998), but few studies have elucidated the nature of base substitutions (interruptions) in microsatellite evolution (Brohede & Ellegren, 1999; Varela *et al*., 2008). Moreover, the genesis of microsatellites still remains a matter of debate (Schlötterer, 2000; Buschiazzo & Gemmell, 2006). Microsatellites are assumed to arise via the creation of a proto-microsatellite, i.e. a short intermediate stage with as few as 3 or 4 repeat units, 56 which is thought to be the substrate for further expansion (Schlötterer, 2000; Buschiazzo $\&$ Gemmell, 2006). The mechanism underlying the genesis and length expansion of short proto-microsatellites, however, is still not fully understood. Results of early studies indicated that short

 proto-microsatellites are highly stable and do not mutate by DNA replication slippage (Messier *et al*., 1996; Rose & Falush, 1998). Further studies, however, concluded that microsatellites with few repeats could gain and lose repeat units (Pupko & Graur, 1999; Zhu *et al*., 2000; Dieringer & Schlötterer, 2003).

 The genus *Morone* is comprised of four species found in North America, including a pair of closely related species, *Morone americana* (Gmelin) and *Morone mississippiensis* Jordan & Eigenmann, and their relatives *Morone saxatilis* (Walbaum) and *Morone chrysops* (Rafinesque). The *Morone* species supported valuable commercial and recreational fisheries. The diversity of MHC is important in the immune diversity of populations. However, no analysis of MHC diversity has been conducted in *Morone* species. In this study, a sequence analysis of a fragment of an MHC class Ia gene was performed in the four *Morone* species. The fragment was from a genomic clone of striped bass, initially developed as a marker for population genetic analysis (Leclerc *et al*., 1996). When the sequence was subjected to a homology search against the five available ENSEMBL teleost whole genome database, hits on 3' downstream sequences of multiple MHC class Ia genes on groupX of the three-spined stickleback, *Gasterosteus aculeatus* L. genome were detected. The sequence was then blasted against the NCBI EST database, it was found that the whole fragment was transcripted, and the ESTs were from European seabass *Dicentrarchus labrax* L. tissues infected with *V. anguillarum* or Nodavius. All evidence demonstrated that the fragment was in the 3' untranslated region of an *Morone* MHC class Ia gene. Two microsatellite loci were identified in this region. One, designated SB83, is a perfect microsatellite and the other, designated SB84, is a short interrupted compound repeat (Fig. 1). The primary objectives of this study were: 1) to analyze the allelic variation of the MHC class Ia fragment in the four *Morone* species; 2) to test whether there is trans-species evolution

products produced by PCR-mediated recombination were identified by repeating the cloning and

 teleost fish species (Clark *et al*., 2001). For example, in East African cichlid fishes, the number of MHC class II B loci per haplotype varies from individual to individual, ranging from 1 to 13 (Málaga-Trillo *et al*., 1998). Similarly, more than two alleles per individual were detected in some individuals of *M. saxatilis*, *M. americana*, and *M. mississippiensis*, indicating that this locus was duplicated either in some individuals or in all individuals in these species.

 Both striped bass and white bass have low levels of polymorphisms compared to other species of fish (Rogier *et al*., 1985; Leclerc *et al*., 1996; Diaz *et al*., 1998; White, 2000; Han & Ely, 2002). However, the level of polymorphism is five times lower in white bass than in striped bass (Han & Ely, 2002). Similar results were obtained in this study where white bass had extremely lower levels of polymorphism than any of the other *Morone* species. In most species, a high level of genetic diversity has been maintained at the MHC locus by balancing selection (Hughes & Yeager, 1998). The lower levels of polymorphism suggest that a dramatic population bottleneck may have reduced the MHC diversity in white bass. Low levels of MHC polymorphism have been reported among other species known to have gone through population bottlenecks (Mikko & Andersson, 1995; Mikko *et al*., 1999; Miller & Lambert, 2004; Wan *et al*., 2006; Mainguy *et al*., 2007).

 Identical numbers of SB83 repeats were observed in different MFR allelic lineages both in the same species and in different *Morone* species (Fig. 2). Thus, contemporary SB83 allele sizes cannot be used to reconstruct the evolutionary history of this complex locus. Polarity of base substitutions within the repeats of SB83 also was observed. Two substitutions occurred at a terminal repeat and the other three substitutions occurred in the fourth or fifth repeat from an end.

 Similar results were reported in previous studies focusing on base substitution within microsatellite repeat arrays. Through sequence analysis of 22 orthologous bovine and ovine 244 (CA)n loci, Brohede & Ellegren (1999) found that the substitution rate in the end of microsatellites was significantly higher than that in the middle of repeat regions. These authors proposed three models that could account for the relatively high mutation rate at the ends of the arrays, all of which were associated with the inefficiency of the mismatch repair system during either replication or recombination. Varela *et al*. (2008) explored the distribution of interruptions in dinucleotide repeats from the human genome and found that the interruptions tended to be towards the ends of microsatellites as well. Another feature of SB83 is that five of the six observed base substitutions occurred at a G in the SB83 microsatellite. A similar result was observed in dinucleotide repeats of human genome. Varela *et al*. (2008) found that mutations involving the substitution of C were approximately 3 times more frequent than those involving substitutions of A in (AC)10.

 In contrast to SB83, SB84 is a compound system of repeats with a complex mutation pattern and a slower rate of evolution. Several alleles of SB84 were found in more than one species suggesting that the complexity at this locus predates speciation. Also, three SB84 alleles were shared by distantly-related flanking region alleles suggesting that intragenic recombination may play a role in generating diversity in the SB83/SB84 region as well.

 In conclusion, high levels of allelic diversity and gene duplication were detected in the MHC class Ia fragment of *M. saxatilis*, *M. americana*, and *M. mississippiensis*. Extremely low levels of diversity were detected in *M. chrysops*, which could be the result of severe population

- Málaga-Trillo, E., Zaleska-Rutczynska, Z., McAndrew, B., Vincek, V., Figueroa, F., Sültmann,
- H. & Klein, J. (1998) Linkage relationships and haplotype polymorphism among cichlid MHC class II B loci. *Genetics* **149**, 1527-1537.
- Messier, W., Li, S.H. & Stewart, C.B. 1996. The birth of microsatellites. *Nature* **381**, 483.
- Mikko, S. & Andersson, L. (1995) Low major histocompatibility complex class II diversity in
- European and North American moose. *Proceedings of the National Academy of Sciences of the United States of America* **92**, 4259-4263.
- Mikko, S., Røed, K., Schmutz, S. & Andersson, L. (1999) Monomorphism and polymorphism at
- Mhc DRB loci in domestic and wild ruminants. *Immunological Reviews* **167**, 169-178.
- doi: 10.1111/j.1600-065X.1999.tb01390.x
- Miller, H.C. & Lambert, D.M. (2004) Genetic drift outweighs balancing selection in shaping post-bottleneck major histocompatibility complex variation in New Zealand robins
- (Petroicidae). *Molecular Ecology* **13**, 3709-3721. doi: 10.1111/j.1365-
- 294X.2004.02368.x
- Ottová, E., Šimková, A., Martin, J-F., Bellocq, J.G.de, Gelnar, M., Allienne, J.F. & Morand, S.
- (2005) Evolution and trans-species polymorphism of MHC class IIβ genes in cyprinid
- fish. *Fish & Shellfish Immunology* **18**, 199-222. doi: 10.1016/j.fsi.2004.07.004
- role of length and number of repeated units. *Journal of Molecular Evolution* **48**, 313-316. doi: 10.1007/PL00006474

Pupko, T. & Graur, D. (1999) Evolution of microsatellites in the yeast *Sacharomyces cerevisiae*:

- Rogier, C.G., Ney, J.J. & Turner, B.J. (1985) Electrophoretic analysis of genetic variability in a
- landlocked striped bass population. *Transactions of the American Fisheries Society* **114**,
- 244-249.
- Rose, O. & Falush, D. (1998) A threshold size for microsatellite expansion. *Molecular Biology and Evolution* **15**,613-615.
- Schlötterer, C. & Tautz, D. (1992) Slippage synthesis of simple sequence DNA. *Nucleic Acids Research* **20**, 211-215.
- Schlötterer, C. & Pemberton, J. (1994) The use of microsatellites for genetic analysis of natural populations. In: Schierwater, B., Streit, B., Wagner, G., DeSalle, R., (Eds), Molecular ecology and evolution: approaches and applications. Birkhäuser Verlag, Basel,
- Switzerland, pp. 203-214.
- Schlötterer, C. (2000) Evolutionary dynamics of microsatellite DNA. *Chromosoma* **109**, 365-371. doi: 10.1007/s004120000089
- Tautz, D. (1993) Notes on the definition and nomenclature of tandemly repetitive DNA
- sequences. In: Pena, S.D.J., Chakraborty, R., Epplen, J.T., Jeffreys, A.J., (Eds), DNA
- fingerprinting: State of the Science. Birkhäuser Verlag, Basel, Switzerland, pp. 21-28.
- Tamura, K., Dudley, J., Nei, M. & Kumar, S. (2007) MEGA4: Molecular Evolutionary Genetic
- Analysis (MEGA) software version 4.0. *Molecular Biology and Evolution* **24**, 1596-1599.
- Varela, M.A., Sanmiguel, R., Gonzalez-Tizon, A. & Martinez-Lage, A. (2008) Heterogeneous
- nature and distribution of interruptions in dinucleotides may indicate the existence of
- biased substitutions underlying microsatellite evolution. *Journal of Molecular Evolution*
- **66**, 575-580. doi: 10.1007/s00239-008-9107-3
- Wan, Q.H., Zhu, L., Wu, H. & Fang, S.G. (2006) Major histocompatibility complex class II
- variation in the giant panda (*Ailuropoda melanoleuca*). *Molecular Ecology* **15**, 2441-
- 2450. doi: 10.1111/j.1365-294X.2006.02966.x
- White, M.M. (2000) Genetic variation in white bass. *Transactions of the American Fisheries Society* **129**, 879-885.
- Zhu, Y., Strassmann, J.E. & Queller, D.C. (2000) Insertions, substitutions, and the origin of
- microsatellites. *Genetical Research* **76**, 227-236.

 Table I, List of samples analyzed in the present study. The number of clones sequenced, alleles, and name of Microsatellite flanking region (MFR) alleles were also shown. The MFR sequence designations were defined by their organism origins. Species are designated by four-letter abbreviations of the Latin names (one from the first letter of genus name and three from the first three letters of species name, e.g. *Msax* for *Morone saxatilis*).

	SB83	SB ₈₄	# observed
		(GT)4GCCT(GT)4AT	
		(GT)4GCCT	З
		(GT)4GCCT	
		(GT)6GCCT	
		(GT)5GCCT	5(5/0)
		(GT)6AT	6(5/1)
		(GT)4GCCT	
		(GT)5GCCT	5(4/1)
		(GT)4GCCT(GT)2CT(GT)3(AT)2	
		(GT)5GCCT	
		(GT)5GCCT	
		(GT)5GCCT	
		(GT)4GCCT	
		(GT)4GCCTGTAT	
		(GT)4GCCT(GT)4AT	
		(GT)4GCCT(GT)4AT	
98 82		(GT)4GCCT(GT)4AT	
	— <i>Мате03-------------</i> -(GT)10	(GT)4GCCT(GT)6(AT)2	
		(GT)4GCCT(GT)2CT(GT)3(AT)2	8
		(GT)4GCCT(GT)2CT(GT)3(AT)2	8(6/2)
99		(GT)4GCCT(GT)2CT(GT)3(AT)2	
		(GT)3GCCT(GT)2CT(GT)3(AT)2	
		(GT)3GCCT(GT)2CT(GT)3(AT)2	5(2/3)
100		(GT)3GCCT(GT)2CT(GT)3(AT)2	
		(GT)3GCCT(GT)2CT(GT)3(AT)2	
82		(GT)4GCCT(GT)4AT	
		(GT)4GCCT(GT)4AT	
		(GT)4GCCT(GT)4AT	
		(GT)4GCCT(GT)4AT	3(2/1)
98		(GT)4GCCT(GT)6(AT)2	1(1/0)
		(GT)4GCCT(GT)4AT	9
		(GT)4GCCT(GT)4AT	2
		(GT)4GCCT(GT)4AT	2
		(GT)4GCCT(GT)6(AT)2	
		(GT)4GCCT(GT)6AT	
99 88		(GT)4GCCT(GT)3(AT)2	
		(GT)4GCTT(GT)6(AT)3	
99		(GT)4GCTT(GT)6(AT)2	
		(GT)4GCCT(GT)9(AT)2	
		(GT)4GCCC(GT)5TT(GT)4TT(AT)2	
97		(GT)4GCCC(GT)5TT(GT)4TT(AT)2	
	- Dlab01-------------------TT(GT)14GCGTAT	(GT)3GCGCCT(GT)3TT(GT)2	1

 $\frac{}{0.005}$

 Figure 2. Neighbor-joining tree based on 39 MFR sequences with both microsatellite sequences superimposed. Observed instances of each flanking region sequence are shown. Nomenclature of the flanking region alleles were designated as in Table I. Numbers in brackets indicates observed numbers of MFR sequence in each species that shared that sequence.