

3-29-2022

A Resilience-Based Intervention to Mitigate the Effect of HIV-Related Stigma: Protocol for a Stepped Wedge Cluster Randomized Trial

Xiaoming Li

University of South Carolina, xiaoming@mailbox.sc.edu

Shan Qiao

University of South Carolina, shanqiao@mailbox.sc.edu

Xueying Yang

University of South Carolina, xueyyang@mailbox.sc.edu

Sayward Harrison

University of South Carolina, HARRI764@mailbox.sc.edu

Cheuk Chi Tam

University of South Carolina, ctam@mailbox.sc.edu

See next page for additional authors

Follow this and additional works at: https://scholarcommons.sc.edu/psyc_facpub



Part of the [Psychology Commons](#)

Publication Info

Published in *Frontiers in Public Health*, Volume 10, 2022.

This Article is brought to you by the Psychology, Department of at Scholar Commons. It has been accepted for inclusion in Faculty Publications by an authorized administrator of Scholar Commons. For more information, please contact digres@mailbox.sc.edu.

Author(s)

Xiaoming Li, Shan Qiao, Xueying Yang, Sayward Harrison, Cheuk Chi Tam, Zhiyong Shen, and Yuejiao Zhou



A Resilience-Based Intervention to Mitigate the Effect of HIV-Related Stigma: Protocol for a Stepped Wedge Cluster Randomized Trial

Xiaoming Li^{1,2*}, Shan Qiao^{1,2}, Xueying Yang^{1,2}, Sayward E. Harrison^{1,3}, Cheuk Chi Tam^{1,2}, Zhiyong Shen⁴ and Yuejiao Zhou⁴

¹ South Carolina SmartState Center for Healthcare Quality, Arnold School of Public Health, University of South Carolina, Columbia, SC, United States, ² Department of Health Promotion, Education, and Behavior, Arnold School of Public Health, University of South Carolina, Columbia, SC, United States, ³ Department of Psychology, College of Arts and Sciences, University of South Carolina, Columbia, SC, United States, ⁴ Guangxi Zhuang Autonomous Region Center for Disease Prevention and Control, Guangxi, China

OPEN ACCESS

Edited by:

Zixin Wang,
The Chinese University of Hong
Kong, China

Reviewed by:

Ranjit Kumar Dehury,
University of Hyderabad, India
Weiming Tang,
University of North Carolina at Chapel
Hill, United States

*Correspondence:

Xiaoming Li
xiaoming@mailbox.sc.edu

Specialty section:

This article was submitted to
Public Health Education and
Promotion,
a section of the journal
Frontiers in Public Health

Received: 18 January 2022

Accepted: 10 March 2022

Published: 29 March 2022

Citation:

Li X, Qiao S, Yang X, Harrison SE, Tam CC, Shen Z and Zhou Y (2022) A Resilience-Based Intervention to Mitigate the Effect of HIV-Related Stigma: Protocol for a Stepped Wedge Cluster Randomized Trial. *Front. Public Health* 10:857635. doi: 10.3389/fpubh.2022.857635

Background: Despite decades of global efforts to tackle HIV-related stigma, previous interventions designed to reduce stigma have had limited effects that were typically in the small- to-moderate range. The knowledge gaps and challenges for combating HIV-related stigma are rooted both in the complexity of the stigma and in the limitations of current conceptualizations of stigma reduction efforts. Recent research has shown the promise of resilience-based approaches that focus on the development of strengths, competencies, resources, and capacities of people living with HIV (PLWH) and their key supporting systems (e.g., family members and healthcare providers) to prevent, reduce, and mitigate the negative effects of stigma. However, the resilience-based approach, while hypothesized, has rarely been empirically tested in large intervention trials, especially in resource-limited settings.

Methods: In this study, we propose to develop, implement, and evaluate a theory-guided, multilevel, multimodal resilience-based intervention *via* a stepped wedge cluster randomized trial among 800 PLWH and their biological or surrogate family members, as well as 320 healthcare providers in Guangxi, China with a longitudinal follow-up period of 36 months at 6-month intervals. The primary outcome will be viral suppression and the intermediate outcomes will include perceived stress and medication adherence of PLWH as well as resilience measures at the level of the individual, the family, and the healthcare system.

Discussion: The proposed study will be one of the first large scale efforts to examine whether resilience among PLWH can be fostered and sustained through a multilevel and multi-component HIV-related stigma intervention and whether a resilience-based intervention can improve clinical outcomes and quality of HIV care among PLWH in a low-resource setting. If efficacious, the intervention components could be tailored to other groups of PLWH and adapted for other low- and middle-income countries.

Trial Registration: This trial is registered at ClinicalTrials.gov, registration number NCT05174936, registered 13 December 2021. <https://register.clinicaltrials.gov/prs/app/action/LoginUser?ts=3&cx=-jg9qo2>.

Keywords: HIV, stigma, resilience, intervention, stepped wedge cluster randomized trial, viral suppression

INTRODUCTION

The finding from the HPTN 052 trial of a 96% reduction in HIV incidence among discordant couples when the HIV-positive partner receives antiretroviral therapy (ART) has led to the emergence of “treatment as prevention” as the dominant strategy to end the global HIV epidemic (1–3). However, numerous obstacles continue to prevent appropriate treatment and optimal clinical outcomes, including stigma against people living with HIV (PLWH). This is a significant public health problem worldwide, particularly in low- and middle-income countries (LMICs), including China. Stigma and discrimination related to HIV and AIDS (i.e., “HIV-related stigma”) can be multifaceted (e.g., individual, community, and institutional) and lead to detrimental impacts, such as preventing PLWH from seeking and receiving appropriate treatment and care, contributing to psychiatric disorders (e.g., depression) and low quality of life, and producing poor clinical outcomes among PLWH (4–6).

Despite substantial global efforts to reduce HIV-related stigma, stigma and discrimination remain persistent and widespread and are among the most poorly understood aspects of the epidemic (7). Two earlier systematic reviews of HIV-related stigma reduction interventions conducted in almost 30 countries across a time span of 20 years (8, 9) identified numerous gaps in evidence-based interventions to reduce HIV-related stigma. These gaps included insufficient engagement of PLWH, only targeting a single socioecological level or a single domain of stigma, inadequate measures to evaluate HIV-related stigma reduction, limited public health relevance of the findings (e.g., lack of clinical endpoints), and a lack of rigor in research methodology. Likewise, Rao et al. in their 2019 review (10) of 24 multilevel stigma interventions around the globe confirmed that the effects of these interventions varied widely in magnitude and were typically in the small-to-moderate range. The authors acknowledged that while there has been progress over the past decade in developing and evaluating multilevel stigma interventions, much work remains to strengthen and expand this approach. A collection of studies in the 2020 *AIDS* special issue titled “Reducing stigma and discrimination: innovation in measurement and practice” also suggested that HIV-related stigma remains highly relevant and

persistent in multiple cultural settings and the gains from HIV-related stigma reduction initiatives have often been modest and rarely implemented at scale (7).

Many of the existing knowledge gaps and challenges for combating HIV-related stigma are rooted in the complexity of the stigma and in limitations in current conceptualizations of stigma reduction efforts (11–13). The focus of most existing intervention research on HIV-related stigma has primarily targeted HIV-negative populations with a goal of eradicating or reducing HIV-related stigma against PLWH and their families. However, HIV-related stigma, like other types of social stigma, has proven difficult to eradicate at the population level. This suggests that, in addition to continued efforts to decrease HIV-related stigma globally, interventions are also needed to help individuals targeted by stigma and discrimination, such as PLWH, to cope with stigma-related challenges (e.g., negative interpersonal experiences, stress) (1, 12). As such, based on their HIV Disparities Model and evidence from the global literature, Earnshaw and colleagues proposed a resilience agenda and suggested “enhancing resilience to societal stigma at the individual (e.g., PLWH, family members) and structural levels (e.g., healthcare settings) as a critical strategy to reduce HIV disparities” (page 231) (3). Recent multi-country analyses of data from the PLWH Stigma Index 2.0 also suggest that multilevel interventions are a promising approach to promote resilience among PLWH and to support PLWH engaging in services, adhering to ART, achieving and maintaining viral suppression, and achieving high quality of life (14, 15).

Resilience in the face of adversity or stress is a necessary capacity of human development (16, 17). Over the past three decades, the definition of resilience has evolved from the view of resilience as an individual personality trait (18), to an ecological view that resilience is a dynamic, multidimensional construct that incorporates bidirectional interaction between individuals and their environment (e.g., family, peers, community, society) (6, 18). Resilience can be cultivated or enhanced in an individual's life by the presence of one or more protective factors, such as close relationships with competent and caring partners in family and healthcare settings, a range of problem-solving skills, and a variety of psychological dispositions (e.g., self-esteem, self-confidence, positive future orientation) (9–20). Available data suggest the promises of these protective factors to foster resilience and mitigate the negative effect of stigma on HIV clinical outcomes (21, 22). The global literature has suggested the critical role of family members [either of origin (e.g., parents, spouse, siblings) or of choice (e.g., non-kin individuals who serve a family-like role by providing emotional or other support)] (23, 24) as a “network of mutual commitment” in assisting PLWH in achieving viral suppression and coping with HIV-related

Abbreviations: AIDS, Acquired Immunodeficiency Syndrome; ART, Antiretroviral Therapy; ARV, Antiretroviral; AZT, Zidovudine; CDC, Centers for Disease Control and Prevention; EFV, Efavirenz; HCP, HealthCare Providers; LPV, Lopinavir; LMICs, Low- and Middle-Income Countries; MSM, Men Who Have Sex With Men; NVP, Nevirapine; PLWH, People Living With HIV; POL, Popular Opinion Leaders; RTV, Ritonavir; SEM, Structural equation modeling; TFV, Tenofovir; 3TC, Lamivudine.

stigma (25, 26). The role of supportive family members in coping with HIV-related stigma appears especially salient in collectivistic societies such as China, where socio-cultural norms emphasizing the individual's obligation and responsibility to the family and community may indirectly facilitate the manifestation of HIV-related stigma. These findings indicate that interventions for HIV-related stigma can benefit from taking into account culturally appropriate intra- and interpersonal protective factors and seeking to actively enhance these resilience factors.

Healthcare providers (HCPs) can also play an essential role in promoting resilience against HIV-related stigma. While healthcare systems in China and other LMICs are facing increasing demands to provide quality care to a growing population of PLWH, healthcare settings are often the places where PLWH experience stigma and discrimination (27, 28). Because of the critical importance of engaging in HIV care for PLWH's quality of life and clinical outcomes, global literature has suggested the value of supportive healthcare systems, including HCPs, across various HIV epidemics and socio-cultural settings (28, 29). Previous work in both China (30–32) and other countries (33–38) has shown that interventions among HCPs can reduce HIV-related stigma by improving their knowledge about the prevention and treatment of HIV and AIDS, as well as their willingness to treat and support PLWH, which can lead to better quality of care for PLWH. Recent research, including our own preliminary data (39–43), has suggested the potential utility of adopting a resilience-based approach that focuses on the development of strengths, competencies, resources, and capacities in PLWH, as well as their families and healthcare systems to reduce and actively mitigate the negative effects of stigma. However, this approach, while hypothesized, has not been sufficiently tested in longitudinal studies or large intervention trials, especially in low-resource settings.

CONCEPTUAL FRAMEWORK

Figure 1 presents a multi-component, multi-level conceptual framework we developed that depicts the process of HIV stigma manifestations and role of resilience development in mitigating the negative effects of stigma on psychological, behavioral, and clinical outcomes. This framework is culturally adapted from an existing resilience-based intervention conceptual framework (20) by (1) specifically targeting PLWH and two key supporting sources in their lives, and (2) using HIV-related clinical outcomes as endpoint measures. Guided by the core principles of the socio-ecological model of human development (44), positive psychology, and theories of resilience (45), this model focuses on the dynamic process of resilience within the cultural context of multi-layered HIV stigma and emphasizes the centrality of individual assets and supportive social contexts in resilience development (46). Support from family members and HCPs, two critical entities in the context of HIV care and HIV-related stigma, can foster the development of PLWH's internal assets (e.g., self-esteem, self-efficacy, positive future orientation). A positive impact of the proposed intervention on supportive social contexts and PLWH's internal assets will in turn develop

resilience and produce improvement in intermediate outcomes (e.g., stress, medical adherence) and endpoint clinical outcomes among PLWH. The framework emphasizes individual and contextual factors (including diverse family structures and risk profiles) that may mediate or moderate the effects of the proposed intervention (13, 27). The framework also provides a “blueprint” that guides our measures for each of these critical domains and our analytic plan to test intervention effects and potential individual and contextual mediators and moderators of such effects.

AIMS AND OBJECTIVES


This study proposes to develop, implement, and evaluate a theory-guided, multilevel, multimodal resilience-based intervention *via* a stepped wedge cluster randomized trial among 800 PLWH and their biological or surrogate family members, as well as among 320 healthcare providers in Guangxi Zhuang Autonomous Region (“Guangxi”), China with a longitudinal follow-up over a period of 36 months with 6-month follow-up intervals. The specific aims are to:

- (1) Develop a multilevel resilience-based HIV stigma reduction and mitigation intervention program (“RISE-UP”) engaging PLWH, their biological or surrogate family members, and healthcare providers (HCPs). The RISE-UP intervention will target individual factors (e.g., resilience, self-efficacy), family factors (e.g., supportive family members), and structural factors (e.g., supportive HCPs and care facilities) by adapting existing intervention components that have shown preliminary efficacy in China or elsewhere.
- (2) Test the short-, medium-, and long-term efficacy of the RISE-UP intervention through a stepped wedge cluster randomized trial among 800 PLWH-family member dyads and 320 HCPs from ~40 HIV clinics in Guangxi with a longitudinal follow-up over a period of 36 months at 6-month intervals.
- (3) Identify individual and contextual factors that may mediate or moderate the effect of RISE-UP intervention on viral suppression, other clinical outcomes (e.g., CD4), and the intermediate psychosocial and behavioral outcomes (e.g., resilience, stress, medication adherence, quality of life).

METHODS/DESIGN

Population and Setting Research Setting

This proposed study will be conducted in Guangxi, one of the regions in China that is experiencing the fastest growth of the HIV epidemic. A total of 10,060 new HIV cases were reported in Guangxi in 2020 with a male to female ratio of 2.66:1. The HIV epidemic in Guangxi is characterized by: (1) sexual transmission (97.8% of all cases, including 90.5% heterosexual transmission and 7.3% male-to-male sexual transmission); (2) rural residents (72.6%); (3) low education levels (84.9% ≤ middle school education, including 7.3% with no formal education, 44.6% with elementary school education only, and 33.1% with



middle school education only); and (4) late diagnosis (52.0% of newly reported cases were AIDS patients). More than half (54.1%) of the cases were married, followed by divorced or widowed (21.4%). Individuals ≥ 50 years of age accounted for 51.0% of all the cases living with HIV.

Selection of Participating Sites

In Guangxi, geographic units important for our study design include cities that are composed of urban districts, as well as rural counties that are composed of rural townships. There is one designated primary public hospital with an HIV clinic in each urban district and rural township that works under the direction of the city or rural county Center for Disease Control and Prevention (CDC) to conduct clinical management and semi-annual follow-ups for all PLWH in the district or township. In collaboration with Guangxi CDC, we will select two cities and eight rural counties that have the largest number of reported HIV cases to participate in our proposed study. Guangxi CDC will identify all urban districts in the two cities and all rural townships in the eight rural counties that have at least 200 HIV cases. We will randomly select 42 of them (stratified by urban vs. rural) as our project sites (2 for pilot-testing; 40 for the actual intervention trial).

Intervention Components

Our proposed multilevel and multimodal resilience-based intervention program includes: (1) a PLWH component; (2) a family member component; and (3) an HCP component.

PLWH Intervention

The primary goal of the PLWH intervention is to assist PLWH in identifying and developing internal and external resilience resources to aid in coping with HIV stigma (including layered stigma), to mitigate the negative impacts of HIV stigma, and to improve their clinical health outcomes. The intervention curriculum will consist of five interactive training sessions (2 h each) with four specific areas of resilience-building: individual assets (self-esteem, emotion regulation, positive future orientation), coping with a chronic health condition (medical adherence, stress reduction, healthy lifestyle, self-care), relationship building (family relationship, provider-patient relationship), and social support (identifying and seeking social support at various socioecological levels). Each session will address one or more of these areas through interactive learning activities (multimedia presentations, role-plays, group discussions, games, personal testimonies). The curriculum will also address empowerment at both individual and family levels across sessions.

Family Member Intervention

The format of the family member intervention will be similar to the PLWH intervention but will emphasize providing social support for PLWH's resilience building as well as fostering resilience at the family level. The intervention will consist of five sessions of group activities (2 h each), with each session addressing one or more of the following areas: (1) HIV and ART knowledge [utilizing "undetectable equals untransmissible" [U=U] messaging]; (2) support to strengthen the capacity of

PLWH and their family members to adapt to living with HIV; (3) relationship building (family relationships, intimate relationships); (4) emotional and behavioral support for PLWH's adherence to care and treatment, including tailored coping or support strategies to address unique needs of some participants who may be more prone to HIV stigma (e.g., sexual and gender minorities and their family members); and (5) self-care. With appropriate consent from both the PLWH and family members, PLWH will be invited to join with family members for two of the five sessions (i.e., relationship building and adherence support) with the goals of improving the interactions of PLWH with family members and discussing common issues and challenges faced by both parties.

HCP Intervention

The HCP intervention will have two primary goals: reducing HCP's stigmatizing attitudes and practices toward PLWH and other social identities such as sexual and gender minorities, sex workers, and injection drug users, and improving the provider-patient relationship. The HCP intervention will consist of four 1.5-h sessions addressing the following topics: (1) universal HIV precautions and occupational safety; (2) layered stigma against PLWH (e.g., manifestations of layered stigma in clinical settings); (3) best practices in building a good provider-patient relationship (e.g., reducing stigmatizing attitudes and behaviors toward PLWH, respecting patients' rights for privacy related to care and disclosure); (4) best practices in providing high quality patient care for PLWH (e.g., increasing skills and comfort in working with PLWH); (5) building a supportive medical environment for better care of PLWH; and (6) skills and confidence in delivering stigma reduction messages to coworkers. The HCP intervention will adapt Popular Opinion Leaders (POL) principles by teaching participants skills for initiating/disseminating stigma reduction messages/practices to colleagues in the workplace (47). The sessions will also be highly interactive with role-plays, facilitator modeling, group discussions, games, testimonies, and presentations from PLWH and medical experts.

PILOT-TESTING

Pilot-Test the Intervention Curricula

All intervention protocols will be reviewed by our community stakeholders for cultural appropriateness, acceptability, and perceived feasibility. In addition, the intervention curricula will be pilot-tested among 30 PLWH (10 females and 20 males), 30 family members (15 biological family members and 15 family members of choice), and 20 HCPs (8 physicians, 12 nurses or other healthcare professionals). These participants will be recruited from the two clinics (one urban and one rural) that will not participate in the actual intervention trial, and the pilot test will adhere to the same inclusion/exclusion criteria of the actual intervention trial. The main purposes of the pilot test are to assess the comprehensibility of the curriculum; cultural appropriateness of the vignettes, materials, and activities for the target population; and logistics of the implementation (e.g., timing of each session) in order to further improve the program's

feasibility and acceptability. Based on the results of the pilot testing, a workshop will be held with the intervention facilitators and the US-China research team to finalize changes to the curricula for incorporation into the facilitator manuals and implementation protocols.

Pilot-Test the Assessment Instruments

Most of the demographic, psychosocial, and behavioral measurements we have selected for PLWH, family members, and HCPs in this study were field-tested and validated previously in China and have been shown to be reliable and valid for their stated purposes. The measures will be further modified based on the specific aims of this study and global literature on stigma and resilience research among PLWH (21, 22, 48–51). The final drafts of all measures will be reviewed and pilot tested among 40 PLWH, 40 family members, and 20 HCPs to obtain perspectives from the target populations on the clarity, cultural sensitivity, and appropriateness of relevant measures. Pilot testing will follow the same recruitment and consenting process as the actual intervention trial.

INTERVENTION TRIAL

Eligibility Criteria PLWH

The eligibility criteria for PLWH include: (1) aged 18 years or older; (2) at least 3 months since confirmed diagnosis of HIV and/or AIDS; (3) detectable viral load (e.g., viral load ≥ 50 copies/mL) or a viral round during the past year (i.e., a confirmed detectable viral load following a suppression); (4) willing to refer or give permission for us to contact one of their adult family members (either of origin or of choice) to participate (but the decision to participate will solely reside in the family member); (5) willing to provide a hair sample for testing hair cortisol, hair antiretroviral (ARV) concentration, and relevant biomarkers; (6) willing to consent for the retrieval of past viral load and CD4 count data from their medical records; and (7) willing to be randomized to receive the intervention at different time points in the stepped wedge trial. The exclusion criteria for PLWH include (1) cognitive or physical inability to respond to assessment questions or to participate in intervention; (2) currently incarcerated or institutionalized for drug use or sex work; (3) participating in other intervention activities during the current study period; and (4) plan to permanently relocate outside of the province within a year. Mental and physical inability will be screened by the local research team in consultation with physicians at the participating clinics.

Family Members

The eligibility criteria for family members include (1) aged 18 years or older; (2) either family member of origin or family member of choice who provides emotional and other social support to PLWH enrolled in the study; and (3) willing to be randomized (along with PLWH enrolled in the study) to receive intervention at different time points. The exclusion criteria for family members will be the same as for

PLWH. The decision of a family member to or not to participate will not affect the eligibility of PLWH.

HCPs

The eligibility criteria for HCPs include: (1) age 18 years or older; (2) provides healthcare services at a participating HIV clinic; and (3) has regular contact with HIV patients. The exclusion criteria for healthcare providers include a plan to permanently relocate outside of the province within a year.

Participants and Recruitment

Participants in the intervention trial will include 800 PLWH and their family members (one per PLWH, either family of origin or family of choice) and 320 HCPs from public HIV clinics in the 40 urban districts/rural townships. Based on our previous work in China, we anticipate high participation rates (e.g., 90% for PLWH, 80% for family members, and 95% for HCPs).

PLWH Recruitment

The PLWH sample will be randomly recruited from the 40 participating HIV clinics (~20 PLWH per clinic). A systematic random sampling procedure (52) will be used to randomly select prospective participants from the HIV patient registries. Medical staff or case managers at the clinics will use an arbitrary number (e.g., the date of the month) to identify the first case to be sampled from the list of HIV patients, and then every n th case will be selected where n is a pre-calculated interval for each sampling iteration (e.g., ratio of target sample size to the total # of patients in the pool). The process will be repeated (without replacement) until the target sample size is achieved at each clinic. Once a prospective patient is randomly selected, the local team members (who will visit each clinic at least twice a week during the recruitment period) will verify and confirm their eligibility and schedule a meeting to explain the study design following a standardized recruitment script.

Family Member Recruitment

Each PLWH who is eligible and willing to participate in the study will be asked to refer one family member (either of origin or of choice) who is aware of the PLWH's HIV status to participate in the study. The PLWH can give the name to the local research team or bring their selected family member to the clinic for eligibility screening. If the family member is interested, a local study team member will meet with the family member, explain the study design, and obtain appropriate consent. With appropriate consent, we will collect brief information (e.g., family structure, risk profile, and treatment profiles) for those PLWH who either cannot identify a family member (of origin or choice) or whose family member refuses to participate for potential secondary analysis to inform the refinement and scale-up of the proposed intervention in the future.

HCP Recruitment

We will randomly recruit 320 HCPs from the 40 participating clinics (~8 per clinic) where we recruit PLWH. Each of these HIV clinics is typically staffed with 15–20 HCPs, including physicians, nurses, case managers, counselors, and medical social workers who have regular contact with HIV patients. With the

permission from site leaders, we will stratify the HCPs in a clinic by profession (physicians vs. other) and professional rank (high/middle vs. low) and randomly approach providers in each stratum, explain the study design following a standard script, and invite them to participate. Local team will meet with interested providers (along with site leaders if desired by providers) to answer questions related to schedule, site support, and other logistic issues related to their participation in the study.

RANDOMIZATION AND INTERVENTION DELIVERY

We will use a stepped wedge cluster randomized trial for intervention assignment and delivery (**Table 1**). For logistic reasons, the 40 clinics will be grouped into 5 clusters (8 clinics in each cluster) in each urban/rural stratum based on the geographic proximity. The local research team will randomly assign each cluster to one of the 5 intervention starting dates (e.g., Month 19, 25, 31, 37, or 43) using a single sequence of random assignments (e.g., using a shuffled deck of five cards bearing numbers from 1 to 5). The PLWH and family members will be organized into small (and separate) groups (~10 participants per group) in each clinic through which to receive intervention sessions.

PLWH Sessions Delivery

The five 2-h PLWH intervention sessions will be delivered over 5 weeks (one session per week) in the clinics where the PLWH are recruited or in nearby private spaces in community centers. Two trained facilitators will deliver the materials through interactive trainings that include multimedia presentations, group discussions, role-play, personal testimonies, and games. The same two facilitators will deliver all five sessions within a clinic to increase group cohesion and rapport with PLWH. The day and time for sessions will be scheduled in advance, and all participating PLWH will be able to opt in to receive a voice or text reminder from the facilitator on the day before the scheduled session.

Family Member Sessions Delivery

The intervention sessions for family members will be similar to PLWH sessions in terms of format and content and will be led by trained facilitators. Family member intervention sessions will emphasize supporting PLWH to cope with HIV-related stigma and to assist them in improving their clinical outcomes. The majority of sessions will consist of only family members, but PLWH will join their family members for two specific sessions (relationship building and adherence support). To improve the participation of family members, both PLWH and family member sessions will be scheduled on the same days in the same location (so they can travel together). Both participating PLWH and their family members will be consulted during the consenting process on the optimal location and schedules of these group sessions. Family members will also be able to opt in to receive voice or text reminders in advance of scheduled sessions.

TABLE 1 | Stepped wedge cluster randomized trial design.

	Months 1-12	Months 13-18	Months 19-24	Months 25-30	Months 31-36	Months 37-42	Months 43-48	Months 49-54	Months 55-60
Cluster 1 (~8 clinics)	Preparation and Community Collaboration & Pilot-testing	Control Intervention							Dissemination and Project Closure
Cluster 2 (~8 clinics)		Control Intervention							
Cluster 3 (~8 clinics)		Control Intervention							
Cluster 4 (~8 clinics)		Control Intervention							
Cluster 5 (~8 clinics)		Control Intervention							
		Month 19							
			Month 25						
				Month 31					
					Month 37				
						Month 43			

HCP Sessions Delivery

The HCP intervention curriculum consists of four 1.5-h sessions that will be delivered in small groups in the clinic-setting by trained facilitators (i.e., health educators from Guangxi CDC). Ideally, all of the participating HCPs in a clinic will attend the sessions together. However, given the variation of clinical schedules among HCPs in a primary care setting, the delivery schedule and format will be flexible and individually tailored (e.g., four sessions can be given at a rate of one per week or consolidated into two longer sessions). Immediately following randomization, the two intervention facilitators (paired for each intervention delivery) will review scheduling and administrative aspects of the clinics and contextualize each session as necessary based on the providers' workload and clinic schedule. Facilitators will also work with providers during the consent process to develop an individualized plan/schedule in each clinic for assessment, intervention, and follow-up. All HCPs will be able to opt in to receive voice or text reminders for planned sessions.

Program Fidelity

The proposed interventions in this study are intensive and complex, which is necessary to create and sustain meaningful changes in participants. While challenging, adherence or fidelity to the content and delivery of the program across clusters is also critical. In addition to a standard implementation protocol, which includes the uniform training and certification of the intervention facilitators, we propose the following two steps to assure fidelity to content and delivery of the intervention protocol and to assess factors that influence the implementation of the protocol: (1) Monitoring the actual intervention implementation. Following the established procedures implemented in China (53–61), one of the intervention facilitators will complete a Fidelity Process Form (FPF) for each session. The FPF will contain the essential elements and key process measures of the intervention sessions, including content delivered, time allocated, participation rate, and main activities covered by the sessions. The local team staff will collect and analyze the FPF

promptly. If any discrepancy between the protocol and the actual implementation emerges, the intervention facilitators will be informed in a timely manner and necessary steps will be taken to prevent deviations from the intervention protocol; and (2) Audio-recording all sessions. The research staff will randomly select and listen to audio recordings of 20% of the sessions and will then complete an FPF for these recorded sessions. These “independent” process measures will be compared with the ones completed by the facilitators, and feedback concerning fidelity will be provided to the facilitators promptly. Data on the fidelity deviation (e.g., dose variation or substantial modification of activities) will be documented as part of the process evaluation to explore the effect of such fidelity deviation on intervention outcomes.

DATA COLLECTION PROCEDURES

PLWH and Family Member Surveys

Interviewers (who will be blinded to the schedule of intervention delivery) will administer the baseline and all follow-up surveys to participants *via* tablet computers. The tablet will display and read (with a real human voice, utilizing a headset) the survey questionnaire in a private room (e.g., doctor's office) at district/township hospitals where the participants are recruited. This method will help to ensure the privacy and quality of the data collection, and it will also help to reduce the potential impact that varying degrees of literacy have on individual's ability to understand the items. Clarifications or assistance (with the tablet) will be provided on site by the interviewers as needed. It is estimated that the survey will take about 60–90 min to complete. Participants will be instructed to take a short break after every 30 min of assessment or as needed.

HCP Survey

All participating HCPs will complete baseline and follow-up surveys. The questionnaires will be self-administered *via* a tablet, and interviewers will be present during the survey to provide

necessary clarification. It is estimated that the HCP survey will take about 30 min to complete.

Hair Sample Collection

The collection of hair samples from all consenting PLWH at baseline and each annual follow-up will follow the Society of Hair Testing guidelines for drug testing in hair (62). Specifically, 1-cm hair sample (20–30 strands of hair for each sample) will be cut as close to the scalp as possible from the vertex posterior region. The hair thatch then will be completely enclosed by a piece of foil and be sealed into an individual plastic bag labeled with the participant ID. Each hair collection process takes about 2 min. All hair samples will be stored at room temperature prior to shipment to the laboratory for process and assays. Laboratory assays using liquid chromatography/tandem mass spectrometry (LC/MS/MS) methods have been developed and validated to analyze most commonly prescribed ARVs in China such as lamivudine (3TC), lopinavir (LPV), ritonavir (RTV), zidovudine (AZT), nevirapine (NVP), efavirenz (EFV), or tenofovir (TFV) in hair samples (63, 64).

KEY STUDY VARIABLES

Primary measures for PLWH will include: (1) HIV clinical indicators (viral load, CD4 count, and disease progression using the HIV staging system by WHO) to be collected from their medical records with consent; (2) HIV-related stigma, including perceived (or anticipated) stigma, internalized stigma, enacted stigma (65–68), as well as layered stigma against other social identities (e.g., sexual and gender minority, sex work, drug use, migratory status, or poverty); (3) Resilience-related measures of PLWH [coping strategies (69, 70), positive future expectation (71), hopefulness for the future, perceived control over the future (71), personal resilience strengths (72), self-concept (73), relationship self-efficacy (74), emotional regulation (75), self-esteem (76)], family support (perceived emotional support and adherence support, quality of relationship) (77), community support [perceived social support (78)], and healthcare system-related support (perceived acceptance and trust from healthcare facilities and HCPs, quality of provider-patient relationship).

Intermediate measures for PLWH and family members will include (1) stress and mental health [self-report measures of depressive symptoms (CES-D) (79, 80), anxiety (81), and stress [Perceived Stress Scale] (82), and stress-related biomarkers [i.e., hair cortisol]]; and (2) adherence to clinical appointments and medication [both self-report (83, 84) and hair ARV concentration for PLWH who are on ART for at least 4 weeks]. In addition, both PLWH and their family members will complete measures of substance use [tobacco use, alcohol use (AUDIT) (85), and other drug use], sexual behavior and reproductive health, HIV-related quality of life (MOS-HIV) (86), and HIV disclosures (77, 87).

Individual and contextual characteristics (potential moderators). Measures we have used in previous studies will also be utilized in this study to collect information from PLWH and their family members on individual and contextual characteristics that may potentially moderate the effect of

intervention. These measures may include socioeconomic status, family composition, HIV diagnosis history and treatment history, co-infection, HIV-infection among partners or other family members, and experience of treatment side effects (for those PLWH on ART).

Measures for HCPs at both the individual level and institutional level. The individual level measures to be completed by HCPs will include: (1) attitudes and behaviors toward PLWH; (2) attitudes toward privacy or confidentiality protection (88); (3) comfort and self-efficacy in supporting PLWH in their treatment and adherence (88); (4) perceptions of patients' rights to HIV testing and disclosure; (5) perceived provider-patient relationship; (6) knowledge and practice of universal precautions (31); and (7) provider mental health [e.g., depressive symptoms (CES-D) (79, 80), burnout]. The institutional level measures will include both HCP's perceptions and actual observations of facilities' efforts and environments (e.g., policy, guidelines for clinical practice) to integrate stigma reduction into facility culture and clinical practice (89, 90). These measures may include: (1) presence of leaders/team of stigma reduction "champions;" (2) presence of code of conduct and patient "rights;" (3) presence of anti-stigma posters; and (4) presence of additional supportive resources for PLWH and their families.

POWER ANALYSIS

Because of the absence of empirical data on the effect of resilience-based stigma reduction and mitigation intervention in China or any other LMICs, we will conservatively assume a "smaller-than-medium" effect size (Cohen $d = 0.35$) for the long-term effect (i.e., 36-month follow-up) of our proposed intervention on the primary PLWH outcomes [e.g., CD4 and viral load in terms of continuous measures (log transformation) and categorical measures (e.g., <500 vs. ≥ 500 for CD4; detectable vs. undetectable for viral load)]. According to the sample size estimation procedure developed for conventional parallel cluster trials by Cohen (91), 140 participants are needed in each cell (or a total 280) to detect an effect size of 0.35 (Cohen d) in various PLWH outcomes at 36-month follow-up with a two-tailed test at $\alpha = 0.05$ and a power of 0.80. However, since the unit of randomization in the current study is the clinic, the sample size calculation, assuming that the unit is the individual, needs to be adjusted for clustering and design effects. Woertman et al. developed a formula to adjust for the design effects of both clustering and the stepped wedge design (DEsw) (92): $DEsw = (1 + \rho(ktn + bn - 1)) / (1 + \rho(1/2 ktn + bn - 1)) \cdot (3(1 - \rho)) / (2t(k - 1/k))$; where ρ is the ICC, k is the # of steps, t is the # of measurements after each step, n = # of individuals per cluster; and b is the # of baseline assessments. Because no clinic level ρ is available in the literature, we conservatively assume a large $\rho = 0.10$ for clustering effect (for 2 groups in each of the 40 clinics with ~ 10 PLWH per group). According to Woertman's procedure (92), a sample size of 528 is needed to produce an effect sample size of 280 with an $\rho = 0.10$. Based on an assumed 10% annual attrition, $n = 800$ at baseline will produce a sample size of 583 by the 36-months follow-up. Thus, our sample of 800 at baseline (or 583 at 36

months follow-up) will provide adequate power to test the main hypotheses in this study.

DATA ANALYSIS

The stepped wedge design offers some analytic advantages in assessing the intervention effects: (1) The intervention effects can be estimated from both between- and within-cluster comparisons with clusters acting as their own controls; and (2) clusters will receive the intervention at varying time points, providing more flexibility in modeling the effects of intervention timing or consistency in effect across clusters (93).

Testing Primary Hypotheses

The evaluation of the short-, medium-, and long-term efficacy of the proposed intervention will be based on the “intent-to-treat” model. Following the recommendations by Hussey and Hughes (93), the hypothesized intervention effect on PLWH, family member, and HCP outcomes will be assessed using multivariable statistical methods for data from stepped wedge design, including Mixed Effect Model (SAS procedure PROC MIXED) for continuous outcome measures and the Generalized Linear Mixed Effect Model (PROC GLIMMIX) for categorical outcome measures. The mixed effect analysis approach is advantageous for assessing the intervention’s effect since it accounts for a number of key factors, including hierarchical data structure, intraclass correlation (ICC) due to cluster randomization, potential baseline differences in the outcome measures, correlated data due to repeated measures over time, and missing data due to attrition (94–96). In conducting these model-based analyses, exposed (intervention) and unexposed (control) observation periods in the stepped wedge design will take the place of “arms” (i.e., intervention assignment) in conventional parallel cluster trials (93). A significant interaction between the intervention and time will be used as the evidence of an intervention effect. The stepped wedge design will also allow us to examine the way in which the intervention effect develops over time once it is introduced into a cluster and allow us to explore the variation of intervention effects among clusters, using within-cluster comparison of unexposed and exposed observations (93).

Mediation and Moderation Analyses

Mediation will be tested using the methodology developed by Baron and Kenny (97). The method is based on multiple regression models focusing on the predictor (e.g., intervention status), the mediator (e.g., improvement in intermediate outcomes), and the primary intervention outcome. According to Baron and Kenny (97), a mediation effect is identified in line with three conditions: (1) the predictor is significantly associated with the outcome; (2) the predictor is significantly associated with the mediator; and (3) the effect of the predictor on the outcome is significantly reduced in magnitude after including the mediator in the model. The PROC MIXED and PROC GLIMMIX procedures will be used to perform these analyses with continuous and categorical data, respectively. In addition to regression analyses (i.e., PROC MIXED and PROC GLIMMIX),

the Sobel test, as specified by MacKinnon and Dwyer (98), will be used to determine the significance of the mediation effect. To test the moderation effect of contextual factors, we will also follow the procedure suggested by Barron and Kenny (97). We will employ multiple regression analyses and test the significance of the appropriate interaction terms between intervention status and the potential moderators (e.g., PLWH’s gender, sexual identity, migratory status, family structure, risk profile, or disease stage) so that the future scale-ups of the intervention can be appropriately informed. Structural equation modeling (SEM) using LISREL will be employed as a complementary methodology to conduct mediation and moderation analyses (99, 100).

DISCUSSION

The limited success of existing HIV stigma reduction efforts requires new approaches that move beyond existing practice (8, 9, 13, 101). This study applies an innovative stepped wedge cluster randomized trial, provides an alternative to an “eradication-only” approach, and broadens our conceptualization to include both reduction of stigma and development of resilience. The proposed intervention focuses on the strengths, competencies, resources, and capacities in PLWH, their families, and HCPs in not only reducing HIV-related stigma but also actively mitigating the negative effects of stigma. The proposed study will develop a culturally appropriate, effective, and sustainable intervention modality that extends beyond the level of individual PLWH to include interpersonal, structural, and social determinants of HIV-related stigma. The resilience-based approach focusing on building the strengths of PLWH and their key social support systems is particularly appropriate within the cultural context of China and other LMICs where PLWH and their family members are often strongly marginalized and discriminated and often face layered stigma which poses significant challenges to behavioral interventions that aim to eliminate or eradicate such layered stigma (88, 102–107). Informed by both positive psychology and socio-ecological theory, our conceptualization of a resilience-based intervention is a critical step in both understanding resilience within the multiple contexts of PLWH, their families, and their HCPs and improving clinical and other health-related outcomes of PLWH.

The RISE-UP intervention will help PLWH to enhance their personal strengths by providing skills-based training in adaptive coping, problem-solving, relationship building, and social support utilization in a group setting that fosters peer social support. The building of internal strengths among PLWH will be further supported and strengthened by efforts that engage family members and HCPs. According to resilience theory, these two key social support systems are critical for successful adaptation to adversity; the internal strengths and the presence of strong support systems from family members and healthcare facilities will improve PLWH’s psychological health (e.g., decreased symptoms of anxiety and depression, reduced stress, improved future orientation) and will increase PLWH’s engagement in HIV care and adherence to ART; and the improvement in psychological health and treatment adherence

will further improve PLWH's immune functioning, which in turn will improve PLWH's viral suppression and other clinical outcomes (e.g., CD4 count, quality of life) and prevent disease progression, as well as prevent spread of HIV to others (U=U) and end the HIV epidemic.

One of the challenges associated with conducting longitudinal studies is attrition. Effective strategies that engage and retain cohort participants are critical to the integrity of research outcomes. Significant and systematic attrition can reduce the generalizability of outcomes and the statistical power to detect effects of interest. The effective retention strategies in this study will include: (1) An individualized follow-up plan. Because of the diversity in PLWH's living situations and daily schedules, we will advise the participants during the consenting process that they will be followed over time and will consult with each participant about the best way to maintain contact. In this way, we will develop an individualized, flexible follow-up plan for each participant (e.g., optimal way to contact them, optimal time and place to meet, preferred frequency of contacts between assessments); (2) Providing all PLWH and family members with the local team member's cellphone number for them to notify research staff of any address change. In addition, we will ask for at least three alternative contacts (i.e., relatives, friends) who can provide information on the whereabouts of the participant in case of relocation; and (3) Arranging telephone or online contact or home visitation prior to each follow-up assessment and making at least three attempts at contact if a PLWH or family member cannot be located. For confidentiality reasons, none of the communications to alternative contacts will reveal the nature of the study.

Guided by the conceptual framework, we hope that the proposed multilevel and multimodal resilience-based intervention can mitigate the negative effects of HIV-related stigma on the clinical outcomes of PLWH. The proposed program is informed and supported by the extant global literature on HIV-related stigma reduction, resilience research, and our preliminary data. Findings from this rigorously designed study are anticipated to inform future interventions and evidence-based policymaking by identifying strategies to build on the strengths of PLWH and their socio-ecological systems in resource-limited settings.

STUDY DISSEMINATION

To materialize the anticipated social benefits of the proposed research and to maximize the impact of the proposed resilience-based intervention, we will take the following strategies to disseminate the study findings. First, local community forums. We will hold meetings with our community advisory board, local partners and other key stakeholders (including the PLWH and their family members and healthcare providers) in each project site (urban districts or rural townships) to present study findings and prepare a strategic dissemination or scale-up plan with local communities, healthcare systems, NGOs, and government agencies if applicable; Second, scientific communities. We will

publish the data in international and national scientific journals and present the study findings at national and international scientific meetings and conferences. We will capitalize on social media and professional networks that can increase the reach and accessibility of findings, such as open access publication, webinars, files and videos available on websites and publicly available channels (e.g., YouTube), to increase visibility and impact of the scientific publications and presentations; Third, impact on policy. The dissemination efforts of this project will extend beyond the scientific arena and will also target policy makers in China at local (participating sites), regional (Guangxi) and national levels through various policy forums, policy papers, and special presentations. We hope that the anticipated success of the proposed project will prompt policy changes in HIV prevention, treatment and care in China and the lessons learned from the project and the tested intervention strategies (curricula, delivery, and evaluation) can be scaled-up to improve health outcomes of PLWH in China and other LMICs.

ETHICS STATEMENT

This study protocol was reviewed and approved by the University of South Carolina Institutional Review Board (Protocol# Pro00099388) and the Guangxi Institutional Review Board (Protocol# GXIRB2020-39-1). The patients/participants will provide their written informed consent to participate in this study.

AUTHOR CONTRIBUTIONS

XL is the principal investigator of this project and led the study design. SQ contributed to the development of the study. XY and XL led the writing of this protocol manuscript. SH and CT contributed significantly to the editing of this manuscript. ZS and YZ contributed to the conception and design of the study. All authors reviewed and provided comments to improve the manuscript, contributed to the editing, and final approval of the protocol.

FUNDING

Research reported in this publication was supported by the National Institute of Mental Health of the National Institutes of Health under Award Number R01MH127961. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

ACKNOWLEDGMENTS

We would like to acknowledge support and guidance from Drs. Li Li (University of California, Los Angeles), Danhua Lin (Beijing Normal University), Nancy Xiaonan Yu (City University of Hong Kong), and Hrishikesh Chakraborty (Duke University) in designing the study and developing the intervention protocol.

REFERENCES

- Cohen MS, McCauley M, Gamble TR. HIV treatment as prevention and HPTN 052. *Curr Opin HIV AIDS*. (2012) 7:99–105. doi: 10.1097/COH.0b013e32834f5cf2
- The White House. *National HIV/AIDS Strategy for the United States: Updated to 2020*. Washington, DC (2016). Available online at: <https://files.hiv.gov/s3fs-public/nhas-indicators-supplement-dec-2016.pdf> (accessed November 8, 2018).
- PEPFAR. *2018 Progress Report on the U.S. President's Emergency Plan for AIDS Relief (PEPFAR) Strategy for Accelerating HIV/AIDS Epidemic Control (2017–2020)*. (2018). Available online at: <https://www.pepfar.gov/documents/organization/286448.pdf> (accessed November 8, 2018).
- Turan B, Budhwani H, Fazeli PL, Browning WR, Raper JL, Mugavero MJ, et al. How does stigma affect people living with HIV? The mediating roles of internalized and anticipated HIV stigma in the effects of perceived community stigma on health and psychosocial outcomes. *AIDS Behav*. (2017) 21:283–91. doi: 10.1007/s10461-016-1451-5
- Turan B, Smith W, Cohen MH, Wilson TE, Adimora AA, Merenstein D, et al. Mechanisms for the negative effects of internalized HIV-related stigma on antiretroviral therapy adherence in women: the mediating roles of social isolation and depression. *J Acquir Immune Defic Syndr*. (2016) 72:198–205. doi: 10.1097/QAI.0000000000000948
- Katz IT, Ryu AE, Onuegbu AG, Psaros C, Weiser SD, Bangsberg DR, et al. Impact of HIV-related stigma on treatment adherence: systematic review and meta-synthesis. *J Int AIDS Soc*. (2013) 16(3 Suppl. 2):18640. doi: 10.7448/IAS.16.3.18640
- Seeley J, Blanc AK. Reducing stigma and discrimination: new evidence and its implications. *AIDS*. (2020) 34:S1–4. doi: 10.1097/QAD.0000000000002647
- Sengupta S, Banks B, Jonas D, Miles MS, Smith GG. HIV interventions to reduce HIV/AIDS stigma: a systematic review. *AIDS Behav*. (2011) 15:1075–87. doi: 10.1007/s10461-010-9847-0
- Stangl AL, Lloyd JK, Brady LM, Holland CE, Baral S. A systematic review of interventions to reduce HIV-related stigma and discrimination from 2002 to 2013: how far have we come? *J Int AIDS Soc*. (2013) 16(3 Suppl. 2):18734. doi: 10.7448/IAS.16.3.18734
- Rao D, Elshafei A, Nguyen M, Hattenbuehler ML, Frey S, Go VF. A systematic review of multi-level stigma interventions: state of the science and future directions. *BMC Med*. (2019) 17:41. doi: 10.1186/s12916-018-1244-y
- Earnshaw VA, Smith LR, Chaudoir SR, Amico KR, Copenhaver MM. HIV stigma mechanisms and well-being among PLWH: a test of the HIV stigma framework. *AIDS Behav*. (2013) 17:1785–95. doi: 10.1007/s10461-013-0437-9
- Parker R, Aggleton P. HIV and AIDS-related stigma and discrimination: a conceptual framework and implications for action. *Soc Sci Med*. (2003) 57:13–24. doi: 10.1016/S0277-9536(02)00304-0
- Earnshaw VA, Bogart LM, Dovidio JE, Williams DR. Stigma and racial/ethnic HIV disparities: moving toward resilience. *Am Psychol*. (2013) 68:225–36. doi: 10.1037/a0032705
- Gottert A, McClair TL, Pulerwitz J, Friedland BA, and the PLHIV Stigma Index 2.0 Study Group in Cambodia, the Dominican Republic and Uganda. What shapes resilience among people living with HIV? A multi-country analysis of data from the PLHIV Stigma Index 2.0. *AIDS*. (2020) 34(Suppl. 1):S19–31. doi: 10.1097/QAD.0000000000002587
- Friedland BA, Gottert A, Hows J, Baral SD, Sprague L, Nyblade L, et al. The people living with HIV Stigma Index 2.0: generating critical evidence for change worldwide. *AIDS*. (2020) 34(Suppl. 1):S5–18. doi: 10.1097/QAD.0000000000002602
- American Psychological Association (APA). *What Is Resilience?* Washington, DC: APA. Available online at: <http://www.apa.org/helpcenter/road-resilience.aspx> (accessed October 28, 2018).
- Dulin AJ, Dale SK, Earnshaw VA, Fava JL, Mugavero MJ, Napravnik S, et al. Resilience and HIV: a review of the definition and study of resilience. *AIDS Care*. (2018) 30:S6–17. doi: 10.1080/09540121.2018.1515470
- American Psychological Association (APA), Task Force on Resilience and Strength in Black Children and Adolescents. *Resilience in African American Children and Adolescents: A Vision for Optimal Development*. Washington, DC: Author (2008). Available online at: <http://www.apa.org/pi/cyf/resilience.html>
- Earnshaw VA, Lang SM, Lippitt M, Jin H, Chaudoir SR. HIV stigma and physical health symptoms: do social support, adaptive coping, and/or identity centrality act as resilience resources? *AIDS Behav*. (2015) 19:41–9. doi: 10.1007/s10461-014-0758-3
- Li X, Chi P, Sherr L, Cluver L, Stanton B. Psychological resilience among children affected by parental HIV/AIDS: a conceptual framework. *Health Psychol Behav Med*. (2015) 3:217–35. doi: 10.1080/21642850.2015.1068698
- Dale SK, Safren SA. Resilience takes a village: black women utilize support from their communities to foster resilience against multiple adversities. *AIDS Care*. (2018) 33:S18–26. doi: 10.1080/09540121.2018.1503225
- Harrison SE, Li X, Vermund SH. From surviving to thriving: the role of resilience in meeting global HIV goals. *AIDS*. (2019) 33(Suppl. 1):S1–4. doi: 10.1097/QAD.0000000000002242
- Pequegnat W. NIMH Consortium on Families HIV/AIDS. In: Pequegnat W, Bell C, editors. *Family and HIV/AIDS: Cultural and Contextual Issues in Prevention and Treatment*. New York, NY: Springer (2012). p. 3–45.
- Pequegnat W, Bray JH. HIV/STD prevention interventions for couples and families: a review and introduction to the special Issue. *Couple Fam Psychol Res Prac*. (2012) 1:79–93. doi: 10.1037/a0028682
- Knodel J, Hak S, Khuon C, So D, McAndrew J. Parents and family members in the Era of ART: evidence from Cambodia and Thailand. *AIDS Care*. (2011) 23:1264–73. doi: 10.1080/09540121.2011.555741
- Pretorius JB, Greeff M, Freeks FE, Kruger A. A HIV stigma reduction intervention for people living with HIV and their families. *Health SA Gesondheid*. (2015) 21:187–95. doi: 10.1016/j.hsag.2015.11.005
- Rogers SJ, Tureski K, Cushnie A, Brown A, Bailey A, Palmer Q. Layered stigma among health-care and social service providers toward key affected populations in Jamaica and The Bahamas. *AIDS Care*. (2014) 26:538–46. doi: 10.1080/09540121.2013.844762
- Turan B, Rogers AJ, Rice WS, Atkins GC, Cohen MH, Wilson TE, et al. Association between perceived discrimination in healthcare settings and HIV medication adherence: mediating psychosocial mechanisms. *AIDS Behav*. (2017) 21:3431–9. doi: 10.1007/s10461-017-1957-5
- Geter A, Herron AR, Sutton MY. HIV-related stigma by healthcare providers in the United States: a systematic review. *AIDS Patient Care STDS*. (2018) 32:418–24. doi: 10.1089/apc.2018.0114
- Wu S, Li L, Wu Z, Liang LJ, Cao H, Yan Z, et al. A brief HIV stigma reduction intervention for service providers in China. *AIDS Patient Care STDS*. (2008) 22:513–20. doi: 10.1089/apc.2007.0198
- Li L, Wu Z, Liang LJ, Lin C, Guan J, Jia M, et al. Reducing HIV-related stigma in health care settings: a randomized controlled trial in China. *Am J Public Health*. (2013) 103:286–92. doi: 10.2105/AJPH.2012.300854
- Buskin SE, Li L, Yin H, Yu T, McGough JP. HIV/AIDS knowledge and attitudes in Chinese medical professionals and students before and after an informational lecture on HIV/AIDS. *J Public Health Manag Pract*. (2002) 8:38–43. doi: 10.1097/00124784-200211000-00006
- Batey DS, Whitfield S, Mulla M, Stringer KL, Durojaiye M, McCormick L, et al. Adaptation and implementation of an intervention to reduce HIV-related stigma among healthcare workers in the United States: piloting of the FRESH workshop. *AIDS Patient Care STDS*. (2016) 30:519–27. doi: 10.1089/apc.2016.0223
- Nyblade L, Stangl A, Weiss E, Ashburn K. Combating HIV stigma in health care settings: what works? *J Int AIDS Soc*. (2009) 12:15. doi: 10.1186/1758-2652-12-15
- MacQuarrie K, Eckhaus T, Nyblade L. *HIV-Related Stigma and Discrimination: A Summary of Recent Literature (Report Submitted by the International Center for Research on Women [ICRW] to UNAIDS)*. (2009). Available online at: http://data.unaids.org/pub/Report/2009/20091130_stigmasummary_en.pdf
- Nyblade L, Mbuya-Brown RJ, Ezekiel MJ, Addo NA, Kraemer JD. A total facility approach to reducing HIV stigma in health facilities: implementation process and lessons learned. *AIDS*. (2020) 34:S93–102. doi: 10.1097/QAD.0000000000002585
- Mahendra VS, Gilborn L, George B, Samson L, Mudoi R, Jadav S, et al. *Reducing AIDS-Related Stigma and Discrimination in Indian Hospitals*. Horizons Final Report. New Delhi: Population Council (2006).

38. Oanh KTH, Ashburn K, Pulerwitz J, Ogden J, Nyblade L. *Improving Hospital-Based Quality of care in Vietnam by Reducing HIV-Related Stigma and Discrimination*. Horizons Final Report. Washington, DC, Population Council (2008).
39. Zhang L, Li X, Qiao S, Zhou Y, Shen Z, Tang Z, et al. The mediating role of individual resilience resources in stigma-health relationship among people living with HIV in Guangxi, China. *AIDS Care*. (2015) 27:1317–25. doi: 10.1080/09540121.2015.1054338
40. Li X, Harrison S, Fairchild A, Chi P, Zhao J, Zhao G. A randomized controlled trial of a resilience-based intervention on psychosocial well-being of children affected by HIV/AIDS: effects at 6- and 12-month follow-up. *Soc Sci Med*. (2017) 190:256–64. doi: 10.1016/j.socscimed.2017.02.007
41. Qiao S, Ingram L, Li X, Deal M. Resilience resources among African American women living with HIV. *AIDS*. (2019) 33:S35–44. doi: 10.1097/QAD.0000000000002179
42. Zhao Q, Mao Y, Qiao S, Li X, Zhou Y, Shen Z. Psychosocial correlates of health-related quality of life (HRQoL) among people living with HIV (PLHIV) in China: the mediating role of resilience. *AIDS*. (2019) 33:S63–70. doi: 10.1097/QAD.00000000000002180
43. Harrison S, Li X, Zhang J, Zhao J, Zhao G. A cluster randomized controlled trial to evaluate a resilience-based intervention for caregivers of HIV-affected children in China. *AIDS*. (2019) 33:S81–91. doi: 10.1097/QAD.00000000000002181
44. Bronfenbrenner U. Toward an experimental ecology of human development. *Am Psychol*. (1977) 32:513–31. doi: 10.1037/0003-066X.32.7.513
45. Masten AS. Ordinary magic: resilience processes in development. *Am Psychol*. (2001) 56:227–38. doi: 10.1037/0003-066X.56.3.227
46. Cicchetti D, Curtis WJ. Multilevel perspectives on pathways to resilient functioning. *Dev Psychopathol*. (2007) 19:627–9. doi: 10.1017/S0954579407000314
47. Kelly J. Popular opinion leaders and HIV peer education: resolving discrepant findings, and implications for the implementation of effective community programmes. *AIDS Care*. (2004) 16:139–50. doi: 10.1080/09540120410001640986
48. Emler CA, Shiu C, Kim H, Fredriksen-Goldsen K. Bouncing back: resilience and mastery among HIV-positive older gay and bisexual men. *Gerontologist*. (2017) 57(Suppl. 1):S40–9. doi: 10.1093/geront/gnw171
49. Woodliff N, Cluver L, Hatcher AM, Brahmabhatt H. “To be HIV positive is not the end of the world”: resilience among perinatally infected HIV positive adolescents in Johannesburg. *Children Youth Services Rev*. (2016) 70:269–75. doi: 10.1016/j.childyouth.2016.09.039
50. Dale SK, Cohen MH, Kelso GA, Cruise RC, Weber KM, Watson C, et al. Resilience among women with HIV: impact of silencing the self and socioeconomic factors. *Sex Roles*. (2014) 70:221–31. doi: 10.1007/s11199-014-0348-x
51. Mburu G, Ram M, Skovdal M, Bitira D, Hodgson I, Mwai GW, et al. Resisting and challenging stigma in Uganda: the role of support groups of people living with HIV. *J Int AIDS Soc*. (2013) 16(3 Suppl. 2):18636. doi: 10.7448/IAS.16.3.18636
52. Shutt RK. Sampling and generalizability (Ch. 5). In: Shutt RK, editor. *Investigating the Social World: The Process and Practice of Research*. 9th ed. Thousand Oaks, CA: Sage (2019). p. 147–1186.
53. Li X, Qiao S, Zhou Y. iCOPE, a multi-level, cluster randomized, 36-month, parallel-group study to assess the efficacy of HIV disclosure intervention in HIV parental disclosure among parents living with HIV in China. *SAGE Open Medicine*. (2020) 8:1–10. doi: 10.1177/2050312120907821
54. Li X, Wang B, Fang X, Zhao R, Stanton B, Hong Y, et al. Short term effect of a cultural adaptation of voluntary counseling and testing among female sex workers in China: a quasi-experimental trial. *AIDS Educ Prev*. (2006) 18:406–19. doi: 10.1521/aeap.2006.18.5.406
55. Li X, Stanton B, Wang B, Mao R, Zhang H, Qu M, et al. Cultural adaptation of the Focus on Kids program for college students in China. *AIDS Educ Prev*. (2008) 20:1–14. doi: 10.1521/aeap.2008.20.1.1
56. Li X, Zhang L, Mao R, Zhao Q, Stanton B. Effect of social cognitive theory based HIV prevention intervention program among high school students in Nanjing, China. *Health Educ Res*. (2011) 26:419–31. doi: 10.1093/her/cyr001
57. Lin D, Li X, Stanton B, Fang X, Lin X, Xu X, et al. Theory-based HIV-related risk-reduction prevention for Chinese female rural-to-urban migrants. *AIDS Educ Prev*. (2010) 22:344–55. doi: 10.1521/aeap.2010.22.4.344
58. Li X, Lin D, Wang B, Du H, Tam CC, Stanton B. Efficacy of theory-based HIV behavioral prevention among rural-to-urban migrants in China: a randomized controlled trial. *AIDS Educ Prev*. (2014) 26:296–316. doi: 10.1521/aeap.2014.26.4.296
59. Stanton B, Li X, Kahibuta J, Fitzgerald AM, Neumbo S, Kanduumbe G, et al. Increased protected sex and abstinence among Namibia youth following a HIV risk-reduction intervention: a randomized, longitudinal study. *AIDS*. (1998) 2473–80. doi: 10.1097/00002030-199818000-00017
60. Kaljee LM, Genberg B, Riel R, Cole M, Tho Le H, Thoa le TK, et al. Effectiveness of a risk-reduction HIV prevention program for rural Vietnamese adolescents: increasing knowledge and efficacy for condom use. *AIDS Educ Prev*. (2005) 17(3):185–99. doi: 10.1521/aeap.17.4.185.66534
61. Stanton B, Wang B, Deveaux L, Lunn S, Rolle G, Li X, et al. Assessing the effects of a complementary parent intervention and prior exposure to a preadolescent program of HIV-risk reduction for mid-adolescents. *AJPH*. (2015) 105:575–83. doi: 10.2105/AJPH.2014.302345
62. Cooper GA, Kronstrand R, Kintz P. Society of Hair Testing guidelines for drug testing in hair. *Forensic Sci Int*. (2012) 218:20–4. doi: 10.1016/j.forsciint.2011.10.024
63. Wu Y, Yang J, Duan C, Chu L, Chen S, Qiao S, et al. Simultaneous determination of antiretroviral drugs in human hair with liquid chromatography-electrospray ionization-tandem mass spectrometry. *J Chromatogr B*. (2018) 1083:209–21. doi: 10.1016/j.jchromb.2018.03.021
64. Chu L, Wu Y, Duan C, Yang J, Yang H, Xie Y, et al. Simultaneous quantitation of zidovudine, efavirenz, lopinavir and ritonavir in human hair by liquid chromatography-atmospheric pressure chemical ionization-tandem mass spectrometry. *J Chromatogr B*. (2018) 1097–8:54–63. doi: 10.1016/j.jchromb.2018.08.031
65. Earnshaw VA, Chaudoir SR. From conceptualizing to measuring HIV stigma: a review of HIV stigma mechanism measures. *AIDS Behav*. (2009) 13:1160–77. doi: 10.1007/s10461-009-9593-3
66. Berger BE, Ferrans CE, Lashley FR. Measuring stigma in people with HIV: psychometric assessment of the HIV stigma scale. *Res Nurs Health*. (2001) 24:518–29. doi: 10.1002/nur.10011
67. Nyblade L, Jain A, Benkirane M, Li L, Lohiniva A, McLean R, et al. A brief, standardized tool for measuring HIV-related stigma among health facility staff: results of field testing in China, Dominica, Egypt, Kenya, Puerto Rico and St. Christopher & Nevis. *J Int AIDS Soc*. (2013) 16:18718. doi: 10.7448/IAS.16.3.18718
68. Kalichman SC, Simbayi LC, Jooste S, Toefy Y, Cain D, Cherry C, et al. Development of a brief scale to measure AIDS-related stigma in South Africa. *AIDS Behav*. (2005) 9:135–43. doi: 10.1007/s10461-005-3895-x
69. Xiao Z, Li X, Qiao S, Zhou Y, Shen Z. Coping, social support, stigma, and gender difference among people living with HIV in Guangxi, China. *Psychol Health Med*. (2018) 23:18–29. doi: 10.1080/13548506.2017.1300671
70. Carver CS. You want to measure coping but your protocol's too long: consider the Brief COPE. *Int J Behav Med*. (1997) 4:92–100. doi: 10.1207/s15327558ijbm0401_6
71. Zhang J, Zhao G, Li X, Hong Y, Fang X, Barnett D, et al. Positive future orientation as a mediator between traumatic events and mental health among children affected by HIV/AIDS in rural China. *AIDS Care*. (2009) 21:1508–16. doi: 10.1080/09540120902923048
72. Connor KM, Davidson JRT. Development of a new resilience scale: the Connor-Davidson resilience scale (CDRISC). *Depress Anxiety*. (2003) 18:76–82. doi: 10.1002/da.10113
73. Fitts WH. *Tennessee Self Concept Scale, Manual*. Los Angeles, CA: Western Psychological Services (1991).
74. Song Z, Lim A. *The Impact of Relationship Self-Efficacy and Cognitive Appraisals on Affective Responses to Social Interactions*. National University of Singapore & University of Colorado at Boulder (2000). Available online at: <http://research.nus.biz/Documents/Research%20Paper%20Series/rps0716.pdf>
75. Gratz KL, Roemer L. Multidimensional assessment of emotion regulation and dysregulation: development, factor structure, and initial validation of

- the difficulties in emotion regulation scale. *J Psychopathol Behav Assessment*. (2004) 26:41–54. doi: 10.1023/B:JOBA.0000007455.08539.94
76. Rosenberg M. Self-concept and psychological well-being in adolescence. In R. L. Leahy (Ed.). *The Development of the Self*. Orlando, FL: Academic Press. (1965)
 77. Qiao S, Li X, Zhou Y, Shen Z, Tang Z. Interpersonal factors associated with HIV partner disclosure among HIV-infected people in China. *AIDS Care*. (2016) 28:37–43. doi: 10.1080/09540121.2016.1146397
 78. Zimet GD, Dahlem NW, Zimet SG, Farley GK. The multidimensional scale of perceived social support. *J Pers Assess*. (1988) 52:30–41. doi: 10.1207/s15327752jpa5201_2
 79. Radloff LS. The CES-D scale: a self report depression scale for research in the general population. *Appl Psych Meas*. (1997) 1:385–401. doi: 10.1177/014662167700100306
 80. Lv Y, Zhao Q, Li X, Stanton B, Fang X, Lin X, et al. Depression symptoms among caregivers of children in HIV affected families in rural China. *AIDS Care*. (2010) 22:669–76. doi: 10.1080/09540120903334633
 81. Spitzer RL, Kroenke K, Williams JB, Lowe B. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med*. (2006) 166:1092–7. doi: 10.1001/archinte.166.10.1092
 82. Cohen S, Kamarck T, Mermelstein R. A global measure of perceived stress. *J Health Soc Behav*. (1983) 24:385–96. doi: 10.2307/2136404
 83. Sabin LL, DeSilva MB, Hamer DH, Xu K, Zhang J, Li T, et al. Using electronic drug monitor feedback to improve adherence to antiretroviral therapy among HIV-positive patients in China. *AIDS Behav*. (2010) 14:580–9. doi: 10.1007/s10461-009-9615-1
 84. Saberi P, Mayer K, Vittinghoff E, Naar-King S; Adolescent Medicine Trials Network for HIV/AIDS Interventions Correlation between use of antiretroviral adherence devices by HIV-infected youth and plasma HIV RNA and self-reported adherence. *AIDS Behav*. (2015) 19:93–103. doi: 10.1007/s10461-014-0806-z
 85. Saunders JB, Aasland OG, Babor TF, de la Fuente JR, Grant M. Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO collaborative project on early detection of persons with harmful alcohol consumption. II. *Addiction*. (1993) 88:791–804. doi: 10.1111/j.1360-0443.1993.tb02093.x
 86. Hsiung PC, Fang CT, Lee KL, Sheng WH, Wu CY, Wang JD, et al. Validation of the medical outcomes study HIV (MOS-HIV) health survey among HIV-infected patients in Taiwan. *Qual Life Res*. (2011). 20:281–6. doi: 10.1007/s11136-010-9733-2
 87. Niccolai LM, King E, D'Entremont D, Pritchett EN. Disclosure of HIV serostatus to sex partners: a new approach to measurement. *Sex Transm Dis*. (2006) 33:102–5. doi: 10.1097/01.olq.0000194591.97159.66
 88. Li L, Wu Z, Wu S, Zhao Y, Jia M, Yan Z. HIV-related stigma in health care settings: a survey of service providers in China. *AIDS Patient Care STDs*. (2007) 21:753–62. doi: 10.1089/apc.2006.0219
 89. Kidd R, Clay S, Stockton M, Nyblade L. *Facilitator's Training Guide For A Stigma-Free Health Facility*. Washington, DC: Futures Group, Health Policy Project (2015). Available online at: http://www.healthpolicyproject.com/pubs/281_SDTrainingGuide.pdf (accessed April 2, 2021).
 90. Nyblade L, Addo NA, Atuahene K, Alsoufi N, Gyamera E, Jacinthe S, et al. Results from a difference-in-differences evaluation of health facility HIV and key population stigma-reduction interventions in Ghana. *J Int AIDS Soc*. (2020) 23:e25483. doi: 10.1002/jia2.25483
 91. Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. 2nd ed. Hillsdale, NJ: Lawrence Erlbaum Associates (1988).
 92. Woertman W, de Hoop E, Moerbeek M, Zuidema SU, Gerritsen DL, Teerenstra S. Stepped wedge designs could reduce the required sample size in cluster randomized trials. *J Clin Epidemiol*. (2013) 66:752–8. doi: 10.1016/j.jclinepi.2013.01.009
 93. Hussey MA, Hughes JP. Design and analysis of stepped wedge cluster randomized trials. *Contemporary Clin Trials*. (2007) 28:182–91. doi: 10.1016/j.cct.2006.05.007
 94. Murray DM, Hannan PJ. Planning for the appropriate analysis in school-based drug-use prevention studies. *J Consult Clinical Psychol*. (1990) 58:458–68. doi: 10.1037/0022-006X.58.4.458
 95. Faraway JJ. *Extending the linear model with R: generalized linear, mixed effects and nonparametric regression models*. Chapman and Hall/CRC (2016).
 96. Littell RC, Miliken GA, Stroup WW, Wolfinger RD. *SAS System for Mixed Model*. 3rd ed. Cary, NC: SAS Institute Inc (1999).
 97. Baron R, Kenny DA. The moderator-Mediator variable distinction in social psychological research: conceptual, strategic, and statistical considerations. *J Pers Soc Psychol*. (1986) 51:1173–82. doi: 10.1037/0022-3514.51.6.1173
 98. MacKinnon DP, Dwyer JH. Estimating mediated effects in prevention studies. *Eval Rev*. (1993) 17:144–58. doi: 10.1177/0193841X9301700202
 99. Cole DA, Maxwell SE. Testing mediational models with longitudinal data: questions and tips in the use of structural equation modeling. *J Abnorm Psychol*. (2003) 112:558–77. doi: 10.1037/0021-843X.112.4.558
 100. MacKinnon D, Lockwood C, Hoffman J. A comparison of methods to test mediation and other intervening variable effects. *Psychol Methods*. (2002) 7:83–104. doi: 10.1037/1082-989X.7.1.83
 101. Turan B, Hatcher AM, Weiser SD, Johnson MO, Rice WS, Turan JM. Framing mechanisms linking HIV-related stigma, adherence to treatment, and health outcomes. *AJPH*. (2017) 107:863–9. doi: 10.2105/AJPH.2017.303744
 102. Liu H, Hu Z, Li X, Stanton B, Naar-King, Yang H. Understanding interrelationships among HIV-related stigma, concern about HIV infection, and intent to disclose HIV serostatus: a pretest-posttest study in a rural area of China. *AIDS Patient Care STDs*. (2006) 20:133–42. doi: 10.1089/apc.2006.20.133
 103. Lau JTF, Choi KC, Tsui HY, Su X. Associations between stigmatization toward HIV-related vulnerable groups and similar attitudes toward people living with HIV/AIDS: branches of the same tree? *AIDS Care*. (2007) 19:1230–40. doi: 10.1080/09540120701402848
 104. Kuhn A. *In China, AIDS Stigma Proves Difficult to Eliminate*. American Public Radio (2009). Available online at: <http://www.npr.org/templates/story/story.php?storyId=120987263>
 105. Cao H, He N, Jiang Q, Yang M, Liu Z, Gao M, et al. Stigma against HIV-infected persons among migrant women living in Shanghai, China. *AIDS Educ Prev*. (2010) 22:445–54. doi: 10.1521/aeap.2010.22.5.445
 106. Blumberg A. China's proposed ban on HIV carriers in public baths highlights larger misconceptions. *The Huffington Post*. (2013). Available online at: http://www.huffingtonpost.com/2013/10/21/china-hiv-ban-bathhouse_n_4138060.html
 107. Wu Z, Detels R, Ji G, Xu C, Rou K, Ding H, et al. Diffusion of HIV/AIDS knowledge, positive attitudes, and behaviors through training of health professionals in China. *AIDS Educ Prev*. (2002) 14:379–90. doi: 10.1521/aeap.14.6.379.24074

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Li, Qiao, Yang, Harrison, Tam, Shen and Zhou. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.