

5-15-2021

Differences in the Proportion of Children Meeting Behavior Guidelines Between Summer and School by Socioeconomic Status and Race

Ethan T. Hunt

Lauren von Klingraeff

Alexis Jones

Sarah Burkart

Roddrick Dugger

University of South Carolina - Columbia, duggerr@email.sc.edu

See next page for additional authors

Follow this and additional works at: https://scholarcommons.sc.edu/sph_health_promotion_education_behavior_facpub



Part of the [Biostatistics Commons](#), [Epidemiology Commons](#), [Exercise Science Commons](#), and the [Public Health Education and Promotion Commons](#)

Publication Info


Published in *Obesity Science and Practice*, Volume 7, Issue 6, 2021, pages 719-726.

This Article is brought to you by the Health Promotion, Education, and Behavior at Scholar Commons. It has been accepted for inclusion in Faculty Publications by an authorized administrator of Scholar Commons. For more information, please contact digres@mailbox.sc.edu.

Author(s)

Ethan T. Hunt; Lauren von Klinggraeff; Alexis Jones; Sarah Burkart; Roddrick Dugger; Bridget Armstrong Ph.D.; Michael Beets Ph.D.; Gabrielle Turner-McGrievy Ph.D., RD; Marco Geraci; and Robert Glenn Weaver MEd, PhD

Differences in the proportion of children meeting behavior guidelines between summer and school by socioeconomic status and race

Ethan T. Hunt¹  | Lauren von Klingraeff¹ | Alexis Jones¹ | Sarah Burkart¹ |
 Rodrick Dugger¹ | Bridget Armstrong¹ | Michael W. Beets¹ |
 Gabrielle Turner-McGrievy² | Marco Geraci^{3,4} | R. Glenn Weaver¹

¹Department of Exercise Science, University of South Carolina, Columbia, South Carolina, USA

²Department of Health Education and Behavior, University of South Carolina, Columbia, South Carolina, USA

³Sapienza – University of Rome, MEMOTEF Department, Rome, Italy

⁴Department of Epidemiology and Biostatistics, University of South Carolina, Columbia, South Carolina, USA

Correspondence

Ethan T. Hunt MPH, Department of Exercise Science Arnold School of Public Health, University of South Carolina, Public Health Research Center, 921 Assembly St. Columbia, SC 29201.

Email: ethunt@email.sc.edu

Funding information

Eunice Kennedy Shriver National Institute of Child Health & Human Development of the National Institutes of Health, Grant/Award Number: R21HD095164.

Abstract

Objective: Children who fail to meet activity, sleep, and screen-time guidelines are at increased risk for obesity. Further, children who are Black are more likely to have obesity when compared to children who are White, and children from low-income households are at increased risk for obesity when compared to children from higher-income households. The objective of this study was to evaluate the proportion of days meeting obesogenic behavior guidelines during the school year compared to summer vacation by race and free/reduced priced lunch (FRPL) eligibility.

Methods: Mixed-effects linear and logistic regressions estimated the proportion of days participants met activity, sleep, and screen-time guidelines during summer and school by race and FRPL eligibility within an observational cohort sample.

Results: Children (n = 268, grades = K – 4, 44.1%FRPL, 59.0% Black) attending three schools participated. Children's activity, sleep, and screen-time were collected during an average of 23 school days and 16 days during summer vacation. During school, both children who were White and eligible for FRPL met activity, sleep, and screen-time guidelines on a greater proportion of days when compared to their Black and non-eligible counterparts. Significant differences in changes from school to summer in the proportion of days children met activity (–6.2%, 95CI = –10.1%, –2.3%; OR = 0.7, 95CI = 0.6, 0.9) and sleep (7.6%, 95CI = 2.9%, 12.4%; OR = 2.1, 95CI = 1.4, 3.0) guidelines between children who were Black and White were observed. Differences in changes in activity (–8.5%, 95CI = –4.9%, –12.1%; OR = 1.5, 95CI = 1.3, 1.8) were observed between children eligible versus uneligible for FRPL.

Conclusions: Summer vacation may be an important time for targeting activity and screen-time of children who are Black and/or eligible for FRPL.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. Obesity Science & Practice published by World Obesity and The Obesity Society and John Wiley & Sons Ltd.

KEYWORDS

children, physical activity, sleep, summer

1 | INTRODUCTION

There is growing evidence that the socioeconomic status of a child's family is a key risk factor for becoming obese.^{1,2} Compelling evidence exists that school-aged children³⁻⁷ and adults^{8,9} from low-income families are at elevated risk for obesity. Recent data from the National Health and Nutrition Examination Survey show that 20% of children from families with a household income at $\leq 130\%$ of the federal poverty level have obesity, while only 10% of children from families with incomes $\geq 350\%$ of the federal poverty level have obesity.¹⁰ Importantly, this gap has increased over time. Independent of income, children who are Black are at an increased risk for obesity when compared to children who are White.^{5,11} This is most likely due to the well-documented effects of structural racism on health behaviors which underlie health disparities.¹²

Summer vacation from school is a critically important time for addressing obesity. A large body of evidence indicates that body mass index (BMI) gain accelerates during the summer.¹³⁻¹⁸ Further, at least one study has shown that the prevalence of children with obesity increases during the months of summer.¹⁵ This acceleration in BMI may be due to engagement in unhealthy behaviors during the summer. For instance, a growing number of studies demonstrate that children engage in less physical activity, spend more time sedentary, and spend more time on screens during the summer than during the school year.¹⁹⁻²¹ Studies are also emerging that show children engage in healthier amounts of sleep and less variable sleep on nights prior to school days, compared to extended breaks from school, like summer.²⁰⁻²² The degradation of health behaviors during summer vacation likely leads to decreased rates of meeting activity,²³ sleep,²⁴ and screen-time guidelines.^{25,26} Failing to meet these guidelines has been associated with increased risk for obesity, insulin resistance, cardiovascular and other diseases.²⁷

The structured days hypothesis,²⁸ which posits that structure, defined as a pre-planned, segmented, and adult-supervised compulsory environment, plays a protective role for children against unhealthy behaviors and, ultimately, prevents the occurrence of negative health-outcomes, such as excessive BMI gain. The structured days hypothesis draws upon concepts in the 'filled-time perspective' literature, which posits that time filled with favorable activities cannot be filled with unfavorable activities.²⁹ This perspective leads to the hypothesis that children engage in a greater number of unhealthy behaviors that lead to increased BMI gain during times that are less-structured (e.g., summer days) than during times that are more structured (e.g., school days). Correspondingly, the Health Gap Hypothesis posits that children from low-income households and children who are Black have relatively less access to structured summer programming (e.g., summer camps) than their middle-to-high income and White counterparts due to financial

barriers and insufficient community resources.³⁰ Thus, summer may disproportionately impact the health behaviors of children from low-income and Black households and ultimately lead to greater accelerated summer BMI gain in these children. Indeed preliminary evidence suggests that children who are Black and children from low-income households experience greater increases in summer BMI gain compared to other children.³¹ Ultimately, greater accelerated summer BMI gain may partially explain the disproportionate risk for obesity born by children from low-income and Black households.

The purpose of this study was to examine the proportion of days children met guidelines for moderate-to-vigorous physical activity (MVPA ≥ 60 min/day),²³ sleep (10-13 h/night for 5 year olds, 9-12 h/night for 6-12 year olds),²⁴ and screen-time (<2 h/day)^{25,26} during the school year compared to the summer, and to examine if these rates differed by race and free/reduced priced lunch (FRPL) eligibility, a proxy of household income. It is hypothesized that (1) during the summer all children will meet physical activity, sleep, and screen-time guidelines on fewer days than during the school year, and (2) children who are eligible for FRPL and children who are Black will experience greater declines in the number of days that they meet physical activity, sleep, and screen-time guidelines than children who are not eligible for FRPL and children who are White.

2 | METHODS

2.1 | Study sample and design

This study utilized data from a larger natural experiment that examined changes in BMI and fitness during the summer vacation and school year for children attending a year-round school and two match paired traditional schools.^{32,33} Physical activity, sleep, and screen-time behavioral data were collected on a subset of ($n = 267$) children participating in the larger study from Spring 2018-Fall 2019. This study presents obesogenic behavior data from school years (2017-2018, 2018-2019) and summers (2018, 2019). All kindergarten through third grade students participating were invited to participate in the behavioral data collection in the Spring of 2018. Measurements commenced in the spring semester of 2018 (i.e., April) and were completed in the fall academic semester of 2019 (i.e., August). Data collection occurred during three distinct one-month measurement periods while school was in session (March and October 2018, and March 2019) and two distinct three-month periods during the traditional summer vacation (May to August 2018 and 2019). For children in the traditional school, summer vacation lasted 11 weeks while summer vacation lasted 5 weeks for children in the year-round school. Prior to the completion of any measures a consent letter was sent home to parents describing study

procedures. Parents who consented were asked to sign and return the letter. All protocols were approved by the University of South Carolina Institutional Review Board prior to enrollment of the first participant.

2.2 | Measures

2.2.1 | Race

Parents reported child's race on a single item screener once at enrollment into the study. Children whose parents reported a race other than White or Black were excluded from the current analysis ($n = 26$).

2.2.2 | Physical activity and sleep

Details fully describing the study can be found elsewhere.³³ Physical activity and sleep were measured using a Fitbit Charge 2™ (Fitbit Inc.). Fitbits were chosen because they provide good agreement with polysomnography and electrocardiography,³⁴ they use multiple heartrate and actigraphy channels to classify sleep which is superior to a single-channel actigraphy,³⁵ can be charged at home, and data is stored in the cloud allowing for data collection over extended periods of time (e.g., 3-months summer vacation). Data processing for physical activity and sleep were informed by the International Study of Childhood Obesity, Lifestyle and the Environment data processing protocols.³⁶ For this analyses, only nocturnal sleep was considered. A valid night of sleep was considered sleep onset that occurred between 5 PM and 6 AM and lasted for greater than 240 min.³⁷ If sleep segments were separated by less than 20 min they were considered one continuous sleep segment.³⁶ Sleep duration was identified as the number of minutes that the Fitbit device classified a child as asleep during a sleep episode. To distill heartrate into activity intensity levels, each child's resting heartrate was calculated as the lowest mean beats-per-minute for 10 consecutive minutes each day.³⁸ Heartrates were distilled into activity intensity levels based on percent heart rate reserve (HRR). Intensity levels are classified as follows: 0.0–19.9% HRR equaled sedentary, 20.0–49.9% of HRR equaled light physical activity, and $\geq 50.0\%$ equaled MVPA.³⁹ An individual day of at least 10 h of waking wear was considered a valid day.³⁶

2.2.3 | Screen-time

Screen-time was assessed via parent proxy-report. Parents completed a questionnaire with their child/children to report their children's screen-time twice per week during measurement periods. Parents were asked to report on their child's daily screen-time on at least 4 days during each 30-days collection period. Parents/children estimated total amount of time (hours and minutes) children spent in

front of a screen that day (e.g., TV, computer, video game, smartphone, and tablet).

2.2.4 | Household income

Poverty-to-income ratio (PIR) was used as a measure of household income. PIR is the ratio of household income to poverty and is calculated by dividing the total reported household income by the Department of Health and Human Services' poverty level.⁴⁰ Parents/guardians were asked to select a household income as a single item in \$10,000 increments. For this analysis, PIR was dichotomized by FRPL status according to the National School Lunch Program.⁴¹ Children living in a household with a $PIR < 1.85$ were classified as eligible to receive FRPL and a $PIR \geq 1.85$ was classified as not eligible to receive FRPL.

2.2.5 | Statistical analyses

First, means and standard deviations of school and child characteristics were examined. Subsequently, regression analysis was used to assess the difference between meeting guidelines (dependent variable) on a school or summer day (independent variable). For each behavior, the dependent variable was operationalized as a binary variable (meeting vs. not meeting the guidelines) or as the proportion of days a child met guidelines. The independent variable was also binary (i.e., school or summer day). Multi-level mixed effect logistic and linear regressions, respectively, were conducted to account for clustering (i.e., days nested within children and children nested within schools). One set of models included race and race-by-condition interactions while a second set of models included FRPL status (< 1.85 PIR vs. ≥ 1.85 PIR) and FRPL-by-condition interactions. All models were adjusted for sex and grade. Analyses exploring the proportion of days children met guidelines by FRPL status included race as a covariate, and models estimating the proportion of days children met guidelines by race included FRPL status as a covariate. Analyses were carried out in Stata (v14.2, College Station TX).

3 | RESULTS

Characteristics of the participating children are presented in Table 1. A total of 267 children participated in the study with 58.1% identifying as Black and 33.0% identifying as White. A total of 51.3% of the participants identified as female and 44.4% eligible for FRPL. During the school year children engaged in 77.6 (SD = 73.7), 470.2 (SD = 68.2), and 100.2 (SD = 89.0) minutes of MVPA, sleep, and screen-time, respectively. During the summer children engaged in 75.3 (SD = 90.7), 486.3 (SD = 91.7), and 145.3 (SD = 120.2) minutes of MVPA, sleep, and screen-time, respectively.

Model implied within- and between-group estimates (including covariates) of the proportion of days and odds of meeting MVPA,

TABLE 1 Participant demographic and behavioral data

	(N) 267	% 100
Race		
Black	155	58.1
White	86	33.0
Other	26	7.9
Sex		
Boys	137	48.7
Girls	130	51.3
Grades		
Kindergarten	17	6.4
1st	56	21.0
2nd	78	29.2
3rd	75	28.1
4th	42	15.7
Income level		
Eligible for FRPL (≤ 1.85 PIR)	120	44.1
Not eligible for FRPL (> 1.85 PIR)	147	55.9
Total weekday behavior data		
	Minutes	±
Moderate-to-vigorous physical activity (n = 267) (n = 14,172 child days)	79.4	77.0
Sleep (n = 209) (n = 4927 child days)	474.7	75.3
Screen-time (n = 195) (n = 2831 child days)	120.8	105.9

Abbreviation: FRPL, free/reduced price lunch.

sleep, and screen-time guidelines during summer vacation and the school year are presented in Figure 1. Children who are Black were less likely to meet MVPA guidelines in the summer compared to the school year (change = -6.2% [95CI = -8.7% , -3.7%]; OR = 0.7 [95CI = 0.6, 0.8]) while children who are White were just as likely to meet MVPA guidelines during the summer compared to the school year (change = 0.0% [95CI = -3.0% , -3.0%]; OR = 1.0 [95CI = 0.9, 1.2]). This translated to children who are Black experiencing a -6.2% ([95CI = -10.1% , -2.3%]; OR = 0.7 [95CI = 0.6, 0.9]) greater decline in MVPA guideline adherence from school to summer when compared to children who are White. Children who are Black were more likely to meet sleep guidelines in the summer compared to the school year (change = 17.0% [95CI = 13.8% , 20.1%]; OR = 3.9 [95CI = 3.0, 5.0]) while children who are White were also more likely to meet sleep guidelines during the summer compared to the school year (change = 9.5% [95CI = 5.9% , 13.2%]; OR = 1.9 [95CI = 1.4, 2.5]). This translated to children who are Black experiencing a 7.4% ([95CI = 2.6% , 12.2%]; OR = 2.0 [95CI = 1.4, 2.9]) greater increase in sleep guideline adherence from school to summer when compared to children who are White. Children who are Black were less likely to meet screen-time guidelines in the summer compared to the school year (change = -21.4% [95CI = -24.8% , -17.0%]; OR = 0.3 [95CI = 0.2, 0.4]) while children who are White were also less likely

to meet screen-time guidelines during the summer compared to the school year (change = -19.4% [95CI = -24.8% , -14.0%]; OR = 0.3 [95CI = 0.2, 0.5]). This translated to no statistically significant difference in school to summer change between children who are Black and children who are White in screen-time guideline adherence (difference in change = -2.0% [95CI = -9.0% , 5.0%]; OR = 0.8 [95CI = 0.5, 1.2]).

Children eligible for FRPL were less likely to meet MVPA guidelines in the summer compared to the school year (change = -8.5% [95CI = -11.0% , -6.0%]; OR = 0.7 [95CI = 0.6, 0.8]) while children not eligible for FRPL were just as likely to meet MVPA guidelines during the summer compared to the school year (change = 0.5% [95CI = -2.2% , -3.2%]; OR = 1.0 [95CI = 0.9, 1.1]). This translated to children who are eligible for FRPL experiencing a -9.0% ([95CI = -12.6% , -5.4%]; OR = 0.6 [95CI = 0.5, 0.8]) greater decline in MVPA guideline adherence from school to summer when compared to children not eligible for FRPL. Children eligible for FRPL were more likely to meet sleep guidelines in the summer compared to the school year (change = 14.7% [95CI = 11.6% , 17.7%]; OR = 3.1 [95CI = 2.4, 4.0]) while children not eligible for FRPL were also more likely to meet sleep guidelines during the summer compared to the school year (change = 12.2% [95CI = 9.0% , 15.3%]; OR = 2.6 [95CI = 2.0, 5.1]). This translated to children eligible for FRPL




FIGURE 1A Proportion of days meeting guidelines on school days and summer vacation by race

experiencing no statistically significant greater increase in sleep guideline adherence from the school to summer compared to children not eligible for FRPL (difference in change = 2.5% [95CI = -1.9%, 6.8%]; OR = 1.2 [95CI = 0.9, 1.8]). Children eligible for FRPL were less likely to meet screen-time guidelines in the summer compared to the school year (change = -21.1% [95CI = -25.4%, -16.8%]; OR = 0.3 [95CI = 0.2, 0.3]) while children not eligible for FRPL were also less likely to meet screen-time guidelines during the summer compared to the school year (change = -20.7% [95CI = -25.6%, -15.7%]; OR = 0.3 [95CI = 0.2, 0.4]). This translated to no statistically significant difference in school to summer change between children eligible for FRPL and children not eligible for FRPL (difference in change = 0.5% [95CI = -6.1%, 7.0%]; OR = 0.9 [95CI = 0.6, 1.3]).

4 | DISCUSSION

In the current study, all children met MVPA and screen-time guidelines on fewer days and sleep guidelines on more days during the summer when compared to the school year. However, children who are Black and eligible for FRPL saw larger decreases in the proportion of days they met MVPA guidelines during the summer than children who are White and not eligible for FRPL. Further, children

who are Black saw a larger increase in the proportion of days they met sleep guidelines during the summer when compared to their White counterparts.

The findings of the current study align with past studies that have found children are less active and engage in more screen-time during periods of less structure (i.e., summer, weekends, or holidays).^{20,28,33,42-45} Given that children who are Black and children from low-income households experience more dramatic accelerations in BMI during the summer than their White and middle-to-high-income counterparts,³¹ the finding in this study that the summer negatively impacted the MVPA of children who are Black and eligible for FRPL to a greater degree than children who are White or not eligible for FRPL is important. This finding suggests a specific behavioral mechanism, decreased MVPA, that may partially explain the greater increases in BMI gain during the summer for children who are Black and eligible for FRPL. Future interventions that target increasing the MVPA of children who are Black and children from low-income households during the months of summer may be warranted.

While the percent of children meeting sleep guidelines was low, it is consistent with past studies that have examined sleep guideline adherence with objective measures.^{46,47} Further, it is not surprising that children met sleep guidelines on more days during the summer when compared to the school year. Past studies have shown that




FIGURE 1B Percent of days meeting guidelines on school days and summer vacation by free/reduced status

children's total sleep time increases on weekends compared to school days and during the summer when compared to the school year.^{21,33,48,49} However, these same studies show that children's bedtimes and wake times shift later and become more variable during the summer. While meeting sleep duration guidelines is protective against developing obesity,^{50,51} sleep timing (i.e., late to bed, late to wake) and stability (i.e., keeping bed and wake time constant) have also been shown to be independent risk factors for obesity.⁵² If children's sleep timing is shifted and becoming more variable during the summer in the current sample, the benefits of meeting sleep duration guidelines may be nullified by later and more variable sleep timing.

Given the findings of the current study coupled with evidence from past studies that show children engage in behaviors that negatively impact their weight status during the summer, intervention strategies to improve children's behaviors during summer are warranted, especially for children who are Black and/or eligible for FRPL. One possible public health strategy is to provide increased access to healthy structured summer programming. At least one study has tested the impact of providing children with access to structured summer programming. Children ($n = 94$) were randomly assigned to either attend a structured summer camp or to experience a typical summer with no access to a structured program.⁵³ Children assigned to attend the summer program lost 0.03 BMI z-score units

while those assigned to not attend gained 0.07 BMI z-score units over the summer. While the differences were not statistically significant they trended in the expected direction. Further children assigned to attend the summer program engaged in 2.3% more MVPA during the program compared to the school year while children not attending the program engaged in 1.9% less MVPA during the summer compared to the school year. This pilot study shows promise for the strategy of providing structured summer programming to enhance health behaviors and mitigate accelerated summer BMI gain.

This study has several strengths including the collection of data continuously for 30+ days during the school year and summer vacation, the within-person design (i.e., same children measured during the school year and summer), and the grounding in theoretical frameworks (i.e., Structured Days Hypothesis and Health Gap Hypothesis). This study also has limitations that must be considered when interpreting the results. First, this study only included three schools in the southeastern United States. Thus, the generalizability of findings may be limited. Second, one of the three schools followed a year-round calendar. Thus, there may be systematic differences in the findings between school calendar types. Third, the study used Fitbit devices to quantify physical activity and sleep. While these devices have shown good agreement with electrocardiography assessment of heart rate and polysomnography assessment of

sleep,^{34,54,55} they have been sparsely used in physical activity and sleep. This limits the ability to compare the findings of this study with other studies.

5 | CONCLUSIONS

During summer, children are less likely to meet guidelines for physical activity and screen-time, providing partial support for the Structured Days Hypothesis.²⁸ This is particularly true for children who are Black or eligible for FRPL, providing support for the Health Gap Hypothesis.³⁰ Interventions that target MVPA and screen-time during times of less structure (i.e., summer), may be warranted.

ACKNOWLEDGEMENT

The first author acknowledges that the publication was supported by award/project number R21HD090647-01A1S1 of the Eunice Kennedy Shriver National Institute of Child Health & Human Development of the National Institutes of Health under parent Award Number R21HD090647. Work on this article by Weaver, Armstrong, and Beets was supported in part by the National Institute of General Medical Sciences of the National Institutes of Health for the UofSC Research Center for Child Well-Being under-Award Number P20GM130420. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Work on this article by Weaver, Armstrong, and Beets was supported in part by the National Institute Of General Medical Sciences of the National Institutes of Health for the UofSC Research Center for Child Well-Being under-Award Number P20GM130420. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

CONFLICT OF INTEREST

The author declares no conflict of interest.

AUTHORS CONTRIBUTIONS

Ethan T. Hunt and R. Glenn Weaver carried out the project. Ethan T. Hunt wrote the manuscript with support from Lauren von Klingraeff, Alexis Jones, Sarah Burkart, Rodrick Dugger, Bridget Armstrong, Michael W. Beets, Gabrielle Turner-McGrievy, Marco Geraci, and R. Glenn Weaver. R. Glenn Weaver helped supervise the project.

ORCID

Ethan T. Hunt  <https://orcid.org/0000-0003-3388-5370>

REFERENCES

- Kim TJ, von dem Knesebeck O. Income and obesity: what is the direction of the relationship? A systematic review and meta-analysis. *BMJ open*. 2018;8(1):e019862.
- LaVeist T, Pollack K, Thorpe R, Jr, Fesahazion R, Gaskin D. Place, not race: disparities dissipate in southwest Baltimore when blacks and whites live under similar conditions. *Health Aff*. 2011;30(10):1880-1887.
- Jones-Smith JC, Dieckmann MG, Gottlieb L, Chow J, Fernald LCH. Socioeconomic status and trajectory of overweight from birth to mid-childhood: the early childhood longitudinal study-birth cohort. *PLoS One*. 2014;9(6):e100181.
- Wang Y, Zhang Q. Are American children and adolescents of low socioeconomic status at increased risk of obesity? Changes in the association between overweight and family income between 1971 and 2002. *Am J Clin Nutr*. 2006;84(4):707-716.
- Weaver RG, Brazendale K, Hunt E, Sarzynski MA, Beets MW, White K. Disparities in childhood overweight and obesity by income in the United States: an epidemiological examination using three nationally representative datasets. *Int J Obes*. 2019;43(6):1210-1222.
- Barriuso L, Miqueleiz E, Albaladejo R, Villanueva R, Santos JM, Regidor E. Socioeconomic position and childhood-adolescent weight status in rich countries: a systematic review, 1990-2013. *BMC Pediatr*. 2015;15(1):129.
- Miech RA, Kumanyika SK, Stettler N, Link BG, Phelan JC, Chang VW. Trends in the association of poverty with overweight among US adolescents, 1971-2004. *Jama*. 2006;295(20):2385-2393.
- Bonney A, Mayne DJ, Jones BD, et al. Area-level socioeconomic gradients in overweight and obesity in a community-derived cohort of health service users-A cross-sectional study. *PLoS One*. 2015;10(8):e0137261.
- Chang VW, Lauderdale DS. Income disparities in body mass index and obesity in the United States, 1971-2002. *Arch Intern Med*. 2005;165(18):2122-2128.
- Ogden CL, Carroll MD, Fakhouri TH, et al. Prevalence of obesity among youths by household income and education level of head of household - United States 2011-2014. *MMWR Morb Mortal Wkly Rep*. 2018;67(6):186-189.
- Ogden CL, Fryar CD, Hales CM, Carroll MD, Aoki Y, Freedman DS. Differences in obesity prevalence by demographics and urbanization in US children and adolescents, 2013-2016. *Jama*. 2018;319(23):2410-2418.
- Williams DR, Mohammed SA. Racism and health I. *Am Behav Sci*. 2013;57(8):1152-1173.
- Von Hippel PT, Powell B, Downey DB, Rowland NJ. The effect of school on overweight in childhood: gain in body mass index during the school year and during summer vacation. *Am J Publ Health*. 2007;97(4):696-702.
- Gutin B, Yin Z, Johnson M, et al. Preliminary findings of the effect of a 3-year after-school physical activity intervention on fitness and body fat: the Medical College of Georgia Fitkid Project. *Int J Pediatr Obes*. 2008;3(Suppl 1):3-9.
- Von Hippel PT, Workman J. From kindergarten through second grade, U.S. Children's obesity prevalence grows only during summer vacations. *Obesity*. 2016;24(11):2296-2300.
- Moreno JP, Johnston CA, Woehler D. Changes in weight over the school year and summer vacation: results of a 5-year longitudinal study. *J Sch Health*. 2013;83(7):473-477.
- Chen T-A, Baranowski T, Moreno JP, et al. Obesity status trajectory groups among elementary school children. *BMC Publ Health*. 2016;16(1):526.
- Chen T-A, Baranowski T, Moreno JP, et al. Obesity status transitions across the elementary years: use of Markov chain modelling. *Pediatric obesity*. 2016;11(2):88-94.
- McCue MC, Marlatt KL, Sirard J. Examination of changes in youth diet and physical activity over the summer vacation period. *Internet J Allied Health Sci Pract*. 2013;11(1):8.
- Brazendale K, Beets MW, Turner-McGrievy GM, et al. Children's obesogenic behaviors during summer versus school: a within-person comparison. *J Sch Health*. 2018;88(12):886-892.
- Weaver R, Armstrong B, Hunt E, et al. The impact of summer vacation on children's obesogenic behaviors and body mass index: a natural experiment. *Int J Behav Nutr Phys Activity*. 2020;153(2020).

22. Weaver RG, Beets MW, Perry M, et al. Changes in children's sleep and physical activity during a one-week versus a three-week break from school: a natural experiment. *Sleep*. 2018;42(1).
23. Piercy KL, Troiano RP, Ballard RM, et al. The physical activity guidelines for Americans. *Jama*. 2018;320(19):2020-2028.
24. Paruthi S, Brooks LJ, D'Ambrosio C, et al. Recommended amount of sleep for pediatric populations: a consensus statement of the American Academy of Sleep Medicine. *J Clin Sleep Med*. 2016;12(6):785-786.
25. Tremblay MS, Carson V, Chaput J-P, et al. Introduction to the Canadian 24-hour movement guidelines for children and youth: an integration of physical activity, sedentary behaviour, and sleep. *Appl Physiol Nutr Metabol*. 2016;41(6):iii-iv.
26. Communications Co, Media MBE. Media use in school-aged children and adolescents. *Pediatrics*. 2016;138(5):e20162592.
27. Biolo G, Ciochi B, Stulle M, et al. Metabolic consequences of physical inactivity. *J Ren Nutr*. 2005;15(1):49-53.
28. Brazendale K, Beets M, Pate RR, et al. Understanding differences between summer vs. school obesogenic behaviors of children: the Structured Days Hypothesis. *Int J Behav Nutr Phys*. 2017;14(1) in press.
29. Caldwell LL, Smith EA. Leisure as a context for youth development and delinquency prevention. *Aust N Z J Criminol*. 2006;39(3):398-418.
30. Weaver RG, Beets MW, Brazendale K, Brusseau TA. Summer weight gain and fitness loss: causes and potential solutions. *Am J Lifestyle Med*. 2019;13(2):116-128.
31. Franckle R, Adler R, Davison K. Accelerated weight gain among children during summer versus school year and related racial/ethnic disparities: a systematic review. *Prev Chronic Dis*. 2014;11(12):e101.
32. Weaver RG, Hunt E, Rafferty A, et al. The potential of a year-round school calendar for maintaining children's weight status and fitness: preliminary outcomes from a natural experiment. *J Sport Health Sci*. 2020;9(1):18-27.
33. Weaver RG, Beets MW, Perry M, et al. Changes in children's sleep and physical activity during a 1-week versus a 3-week break from school: a natural experiment. *Sleep*. 2019;42(1):zsy205.
34. de Zambotti M, Baker FC, Willoughby AR, et al. Measures of sleep and cardiac functioning during sleep using a multi-sensory commercially-available wristband in adolescents. *Physiology Behav*. 2016;158:143-149.
35. Burkart S, Beets MW, Armstrong B, et al. Comparison of multi-channel and single channel wrist-based devices with polysomnography to measure sleep in youth. *J Clin Sleep Med*. 2020;17(4):645-652.jcsm.8980.
36. Tudor-Locke C, Barreira TV, Schuna JM, et al. Improving wear time compliance with a 24-hour waist-worn accelerometer protocol in the international study of childhood obesity, Lifestyle and the environment (ISCOLE). *Int J Behav Nutr Phys Activity*. 2015;12(1):1-9.
37. Acebo C, Sadeh A, Seifer R, et al. Estimating sleep patterns with activity monitoring in children and adolescents: how many nights are necessary for reliable measures? *Sleep*. 1999;22(1):95-103.
38. Welk GJ, Corbin CB. The validity of the Tritrac-R3D activity monitor for the assessment of physical activity in children. *Res Q Exerc sport*. 1995;66(3):202-209.
39. Gavarry O, Bernard T, Giacomoni M, et al. Continuous heart rate monitoring over 1 week in teenagers aged 11-16 years. *Eur J Appl physiology Occup physiology*. 1997;77(1-2):125-132.
40. Department of Health and Human Services. *Department of Health and Human Services Poverty Guidelines for 2017*. 2018.
41. Baranowski T, O'Connor T, Johnston C, et al. School year versus summer differences in child weight gain: a narrative review. *Child Obes*. 2014;10(1):18-24.
42. Wang YC, Vine S, Hsiao A, Rundle A, Goldsmith J. Weight-Related behaviors when children are in school versus on summer breaks: does income matter? *J Sch Health*. 2015;85(7):458-466.
43. Hunt ET, Whitfield ML, Brazendale K, Beets MW, Weaver RG. Examining the impact of a summer learning program on children's weight status and cardiorespiratory fitness: a natural experiment. *Eval Program Plan*. 2019;74:84-90.
44. Dugger R, Brazendale K, Hunt ET, et al. The impact of summer programming on the obesogenic behaviors of children: behavioral outcomes from a quasi-experimental pilot trial. *Pilot Feasibility Stud*. 2020;6(1).
45. Staiano A, Broyles S, Katzmarzyk P. School term vs. School holiday: associations with children's physical activity, screen-time, diet and sleep. *Ijeph*. 2015;12(8):8861-8870.
46. Meredith-Jones K, Galland B, Haszard J, et al. Do young children consistently meet 24-h sleep and activity guidelines? A longitudinal analysis using actigraphy. *Int J Obes*. 2019;43(12):2555-2564.
47. Guan H, Zhang Z, Wang B, et al. Proportion of kindergarten children meeting the WHO guidelines on physical activity, sedentary behaviour and sleep and associations with adiposity in urban Beijing. *BMC Pediatr*. 2020;20(1):1-9.
48. Sun W, Ling J, Zhu X, Lee TM-C, Li SX. Associations of weekday-to-weekend sleep differences with academic performance and health-related outcomes in school-age children and youths. *Sleep Med Rev*. 2019;46:27-53.
49. Wing YK, Li SX, Li AM, Zhang J, Kong APS. The effect of weekend and holiday sleep compensation on childhood overweight and obesity. *Pediatrics*. 2009;124(5):e994-e1000.
50. Patel SR, Hu FB. Short sleep duration and weight gain: a systematic review. *Obesity*. 2008;16(3):643-653.
51. Matricciani L, Paquet C, Galland B, Short M, Olds T. Children's sleep and health: a meta-review. *Sleep Med Rev*. 2019;46:136-150.
52. Olds TS, Maher CA, Matricciani L. Sleep duration or bedtime? Exploring the relationship between sleep habits and weight status and activity patterns. *Sleep*. 2011;34(10):1299-1307.
53. Evans EW, Wing RR, Pierre DF, Howie WC, Brinker M, Jelalian E. Testing the effect of summer camp on excess summer weight gain in youth from low-income households: a randomized controlled trial. *BMC Publ Health*. 2020;20(1):1-10.
54. Liang Z, Martell MAC. Validity of consumer activity wristbands and wearable EEG for measuring overall sleep parameters and sleep structure in free-living conditions. *J Healthcare Informatics Res*. 2018;2:1-27.
55. De Zambotti M, Goldstone A, Claudatos S, Colrain IM, Baker FC. A validation study of Fitbit Charge 2 compared with polysomnography in adults. *Chronobiol Int*. 2018;35(4):465-476.

How to cite this article: Hunt ET, von Klingraeff L, Jones A, et al. Differences in the proportion of children meeting behavior guidelines between summer and school by socioeconomic status and race. *Obes Sci Pract*. 2021;7(6):719-726. <https://doi.org/10.1002/osp4.532>