

1-1-2013

High Fat Diet Induced Obesity in a Mouse Model for Colorectal Cancer

Kennerly Clinton Patrick
University of South Carolina

Follow this and additional works at: <https://scholarcommons.sc.edu/etd>



Part of the [Biomedical Commons](#), and the [Chemicals and Drugs Commons](#)

Recommended Citation

Patrick, K. C. (2013). *High Fat Diet Induced Obesity in a Mouse Model for Colorectal Cancer*. (Doctoral dissertation). Retrieved from <https://scholarcommons.sc.edu/etd/2114>

This Open Access Dissertation is brought to you by Scholar Commons. It has been accepted for inclusion in Theses and Dissertations by an authorized administrator of Scholar Commons. For more information, please contact digres@mailbox.sc.edu.

HIGH FAT DIET INDUCED OBESITY IN A MOUSE MODEL FOR COLORECTAL
CANCER

by

Kennerly Clinton Patrick

Bachelor of Science
University of South Carolina, 2011

Submitted in Partial Fulfillment of the Requirements

For the Degree of Master of Science in

Biomedical Sciences

School of Medicine

University of South Carolina

2013

Accepted by:

Angela Murphy, Director of Thesis

Chandrashekhhar Patel, Reader

Cory Robinson, Reader

Lacy Ford, Vice Provost and Dean of Graduate Studies

© Copyright by Kennerly Clinton Patrick, 2013
All Rights Reserved.

ACKNOWLEDGEMENTS

I would like to acknowledge Dr. Angela Murphy for all the inspiration, support, and guidance during the duration of this research project. I would also like to acknowledge Jamie McClellan and Reilly Enos for their continued assistance and expertise in conducting this project.

ABSTRACT

Obesity is widely correlated with the incidence of colorectal cancer (CRC). The exact mechanisms that link obesity to CRC risk have not yet been fully established, and only a limited number of animal models are available to study pathways involved in obesity-related colon carcinogenesis. The objective of this study was to determine the influence of HFD-induced obesity on AOM/DSS-mediated colorectal tumorigenesis, tumor proliferation, and symptom severity. Male C57BL/6 mice were fed a novel high fat diet, designed to mimic the standard American diet (12 % saturated fat & 28 % unsaturated fat), at 4 weeks of age until 16 weeks of age to induce obesity. At 16 weeks of age HFD-fed mice had significantly increased mean body mass compared to LFD-fed mice. Thus, mice were obese prior to initiation of CRC. Following treatment with AOM/DSS, study results indicated no significant changes in tumor number or size with obesity. Similarly, obesity did not increase the severity of symptoms in this model. These findings challenge previously published data, with the caveat that the low number of tumors detected in the present model limits translational implications for tumor characteristics and symptom severity in obesity.

TABLE OF CONTENTS

ACKNOWLEDGEMENTS	iii
ABSTRACT	iv
LIST OF FIGURES	vi
LIST OF ABBREVIATIONS.....	vii
CHAPTER 1: INTRODUCTION.....	1
CHAPTER 2: MATERIALS & METHODS	4
CHAPTER 3: RESULTS.....	9
CHAPTER 4: DISCUSSION.....	19
REFERENCES	24

LIST OF FIGURES

Figure 3.1 DEXA body composition collected at baseline (16 weeks of age) and at sacrifice (26 weeks of age) on (A) lean mass and BMC, (B) body fat, (C) % lean mass and BMC, (D) % body fat. *Significantly different from LFD-Con ($P \leq .05$). #Significantly different from HFD-Con ($P \leq .05$). \$Significantly different from LFD-CRC ($P \leq .05$)	4
Figure 3.2 The impact of HFD and CRC on (A) percent change in body weight. *Significantly different from LFD-Con (week 17.5-18: HFD-Con & LFD-CRC; week 18.5-19: HFD-Con, LFD-CRC, & HFD-CRC-1 & 2; week 19.5-21.5: HFD-Con, LFD-CRC, & HFD-CRC-1; week 22-24.5: HFD-Con & LFD-CRC; week 25-26: HFD-Con, LFD-CRC, & HFD-CRC-1&2; $P \leq .05$) #Significantly different from HFD-Con (week 16.5-17: LFD-CRC; week 17.5-26: LFD-CRC & HFD-CRC-1&2; $P \leq .05$). \$Significantly different from LFD-CRC (week 17.5-26: HFD-CRC-1&2; $P \leq .05$). (B) The effect of DSS induced inflammation on symptom scores. \$Significantly different from LFD-CRC (week 18.5: HFD-CRC-2 only; $P \leq .05$)	4
Figure 3.3 The effect of HFD and CRC on (A) epididymal, (B) mesentery, (C) kidney, and (D) total visceral (Total) fat pad weights at sacrifice. *Significantly different from LFD-Con ($P \leq .05$). #Significantly different from HFD-Con ($P \leq .05$). \$Significantly different from LFD-CRC ($P \leq .05$)	23
Figure 3.4 Spleen weight at sacrifice as a measure of inflammation and CRC immune response. #Significantly different from HFD-Con ($P \leq .05$)	25
Figure 3.5 The effect of CRC and DSS induced inflammation on (A) colon length and (B) colon weight. *Significantly different from LFD-Con ($P \leq .05$). #Significantly different from HFD-Con ($P \leq .05$). \$Significantly different from LFD-CRC ($P \leq .05$)	23
Figure 3.6 The impact of CRC induction at sacrifice for (A) total polyp number & (B) total polyp number by size. No significant differences between any groups	23
Figure 3.7 Mean (A) food consumption & (B) water consumption per group.	23

LIST OF ABBREVIATIONS

AOM	Azoxymethane
BW	Body Weight
CRC.....	Colorectal Cancer
DEXA	Dual-Energy X-ray Absorptiometry
DSS	Dextran Sodium Sulfate
HFD.....	High Fat Diet
HFD-Con.....	High Fat Diet-Control
HFD-CRC-1	High Fat Diet-Colorectal Cancer Group 1
HFD-CRC-2	High Fat Diet-Colorectal Cancer Group 2
LFD	Low Fat Diet
LFD-Con	Low Fat Diet-Control
LFD-CRC.....	Low Fat Diet-Colorectal Cancer
PBS	Phosphate-Buffered Saline

CHAPTER 1

INTRODUCTION

As the second leading cause of cancer-related deaths for both men and women in the United States, colorectal cancer (CRC) is a national burden. It is the third most commonly diagnosed cancer and is of increasing health concern to a growing aging population ^{1,2}. It has been shown in epidemiological studies that obesity and metabolic syndromes, like hyperglycemia, hypertension, and hyperlipidemia, increase the risk of developing CRC ³. As such, lifestyle factors such as limited physical activity, consumption of food with high fat/limited fiber content, high caloric intake, and obesity are controllable factors that can influence the development of CRC ⁴. Conversely, it has been demonstrated that behavioral factors like exercise and consumption of fruits and vegetables, even among obese individuals, can reduce the incidence of CRC ⁴. While family history is a large component in determining CRC risk, environmental factors are responsible for the majority of cases. In fact, sporadic CRC, the most prominent form of CRC that occurs due to an accumulation of mutations in genes regulating the growth and proliferation of colonic epithelial cells is influenced largely by environmental factors independent of genetic background ⁵.

Obesity is widely correlated with the incidence of CRC ⁶. The exact mechanisms that link obesity to CRC risk have not yet been fully elucidated; however, a number of biochemical changes associated with obesity have been implicated and include inflammatory mediators, adipokines and metabolic markers among others. For example

leptin that is upregulated in obese individuals is a hormone and cytokine produced primarily by adipocytes has been shown to induce CRC by a variety of different pathways ⁴. Similarly, increased plasma levels of insulin observed during obesity disrupt normal metabolic processes and thought to activate pathways involved in oncogenesis ⁴. In addition, inflammatory mediators that are increased in adipose tissue have been linked to every step involved in the development and promotion of CRC. While there is an abundance of obesity-mouse models, the number of models available to study pathways involved in colon carcinogenesis is limited ³.

Among the chemically induced CRC models, the azoxymethane (AOM) and dextran sodium sulfate (DSS) model has been proven to induce CRC in a shortened amount of time with a high level of reproducibility ⁷. This outstanding model has been utilized in a variety of studies investigating the histopathologic and molecular features involved in carcinogenesis as well as determining chemopreventive and therapeutic approaches in combating CRC ⁷. The purpose of this current investigation was to examine the effects of high fat diet induced obesity on CRC. In this study, male C57BL/6 mice were fed a novel high fat diet, designed to mimic the standard American diet (12 % saturated fat & 28 % unsaturated fat), at 4 weeks of age until 16 weeks of age to induce obesity. At 16 weeks of age mice were exposed to the CRC protocol. Thus, mice were obese prior to initiation of CRC. This clinically relevant model aims to mimic the conditions of tumor initiation and progression that occurs in obese humans. Previous studies have co-induced obesity and CRC with the AOM/DSS protocol; however, this approach fails to account for the conditions observed in the at risk obese population prior to the onset of colon cancer. We hypothesized that obesity induced prior to initiation of

CRC would lead to increased severity of sickness symptoms in association with enhanced tumorigenesis. A better understanding of the impact of obesity on CRC is necessary in determining possible therapeutic targets and/or preventative screening and diagnostic techniques.

CHAPTER 2

MATERIALS & METHODS

2.1 ANIMALS

Male C57BL/6 mice, 4 weeks of age, were purchased from Jackson Laboratories (Bar Harbor, ME) and cared for in the animal facility at the University of South Carolina. They were housed four-five per cage and maintained on a 12:12-hr light-dark cycle in a low stress environment (22°C, 50% humidity, low noise) and given food and water *ad libitum*. All animal experimentation was approved by the University of South Carolina's Institutional Animal Care and Use Committee.

2.2 DIETS

Mice were randomly assigned to one of five treatment groups as follows: 1) Low Fat Diet-Control (LFD-Con), 2) High Fat Diet-Control (HFD-Con), 3) Low Fat Diet-Colorectal Cancer (LFD-CRC), 4) High Fat Diet-Colorectal Cancer Group 1 (HFD-CRC-1) or 5) High Fat Diet-Colorectal Cancer Group 2 (HFD-CRC-2). HFD-CRC-1 and HFD-CRC-2 differed in the dose of azoxymethane (AOM) that was administered to initiate the cancer; group 1 received a dose of AOM based on body weight (10mg/Kg) whereas group 2 received a dose of AOM that was equivalent to the LFD-CRC group. This was done in order to eliminate any potential differences among the groups due to the dose of AOM. The HFD (12% saturated fat & 28% unsaturated fat) was designed to mimic the standard American Diet (Bio-Serv, Frenchtown, NJ) (Table 2.1). The AIN-76A diet was used as the LFD (Bio-Serv). Diets were administered beginning at 4 weeks

of age through sacrifice at 26 weeks of age. The HFD retained the same vitamin and mineral content as the control diet. Food and water was available *ad libitum* and measured on a weekly basis.

2.3 BODY WEIGHT & BODY COMPOSITION

Body weight was examined weekly beginning at 4 weeks of age and for the duration of the 22 week study. In addition, body composition was assessed at 16 weeks of age (prior to initiation of the CRC protocol) and at 26 weeks of age (prior to sacrifice). For this procedure, mice were placed under brief anesthesia (isoflurane inhalation) and were assessed for lean mass, fat mass, and body fat percentage via dual-energy x-ray absorptiometry (DEXA) (Lunar PIXImus, Madison, WI).

2.4 COLORECTAL CANCER PROTOCOL

To induce colorectal cancer, mice in the CRC groups underwent a standard protocol consisting of a single subcutaneous injection of azoxymethane (AOM) at 16 weeks of age, followed by 3 cycles of DSS (2% DSS in water for one week followed by two weeks of regular water for the first cycle and 1% DSS in water for the second and third cycles) beginning at week 17. Both the LFD-CRC and the HFD-CRC-1 groups received a dose of AOM based on their body weight (10mg/Kg body weight) whereas HFD-CRC-2 received a dose of AOM that was equivalent to the LFD-CRC group. As mentioned above, this was done in order to eliminate any potential differences among the groups due to the dose of AOM (i.e. an increase in tumor number with HFD that may be due to the increased concentration of AOM that was administered to these mice given their increased weight).

2.5 SYMPTOM SCORING

Symptoms were scored twice weekly throughout the 10 week protocol using a standard scoring system that has previously been used in this model. Briefly, scores were based on diarrhea, blood in stools and weight loss. Diarrhea symptoms were evaluated based on visualization of the fecal matter. Blood in the stools was determined using a Hemoccult Fecal occult blood test. Weight loss was based on the percent change in weight compared to baseline levels. Mice were given scores of 0, 2 or 4 depending on the severity of the symptoms.

2.6 SACRIFICE & TISSUE COLLECTION

At 26 weeks of age, mice were sacrificed for tissue collection via isofluorane overdose. The kidney fat, mesenteric fat and epididymal fat were removed, weighed, and immediately snap-frozen in liquid nitrogen and stored at -80°C. The large intestine was dissected from the cecum to the anus and placed on filter paper. A large needle and syringe was used to flush the large intestine with PBS. The large intestine was then cut longitudinally, flattened with a cotton swab saturated with PBS, and another filter paper was placed on top of the sections. The filter paper was then sandwiched between two pieces of blotting paper, clipped together on all four sides, and subsequently fixed in 10% formalin for at least 24 hours. After that time, the sections were transferred and stored in 70% ethanol at room temperature to be further analyzed for tumor number and size.

2.7 TUMOR COUNTS

To count tumors, the large intestines were rinsed in deionized water, and briefly stained in 0.1% methylene blue. Tumors were counted by the same blinded technician for all samples under a dissecting microscope, using tweezers to pick through the villi to identify tumors. Tumors were classified as being greater than 2mm, less than 2mm but

greater than 1mm, or less than 1mm in diameter using an ocular micrometer. Colon lengths were measured as an indicator of inflammation; a shorter colon length is indicative of greater inflammation. Similarly, colon weight was determined as an increase in colon weight has been associated with elevated inflammation.

Table 2.1 Diet composition of treatment diets

	AIN-76A	12% Saturated Fat Diet
<u>Ingredient (g/kg)</u>		
Casein	200	165
DL Methionine	3	3
Lard	0	35.4
Coconut Oil	0	30
Corn Oil	50	49.9
Soybean Oil	0	9.3
Olive Oil	0	78.4
Corn Starch	0	50
Maltodextrin	0	100
Sucrose	500	381.5
Cellulose	50	50
Vitamin Mix (AIN-76A)	10	10
Mineral Mix (AIN-76A)	35	35
Choline Bitartrate	2	2.5
Energy (kcal/g)	3.61	4.572
<u>Energy (% kcal)</u>		
Carbohydrate	67.5	47
Fat	12.5	40
Protein	20	13
<u>Fatty Acid Profile (g/kg)</u>		
Caprylic Acid (C8:0)	0	2.3
Capric Acid (C10:0)	0	1.8
Lauric Acid (C12:0)	0	13.5
Myristic Acid (C14:0)	0	5.5
Palmitic Acid (C16:0)	5.75	26
Palmitoleic Acid (C16:1)	0	2
Stearic Acid (C18:0)	0	8.5
Oleic Acid (C18:1)	13.3	88
Linoleic Acid (C18:2)	29.4	43.2
α -Linolenic Acid (C18:3)	.40	2.2
Arachidic Acid (C20:0)	.10	0
% of Total Calories from SFAs	1.5%	12%
% of Total Calories from MCFAs (C6:0-C12:0)	-	3.6%
% of Total Calories from LCSFAs (C14:0-C20:0)	1.5%	8.4%
% of Total Calories from USFAs	11%	28%
% of Total Calories from MUFAs	3.3%	18.6%
% of Total Calories from PUFAs	7.6%	9.4%
% of Total Calories from n-3 FAs	.1%	.45%
% of Total Calories from n-6 FAs	7.5%	8.9%
Cholesterol (mg/kg)	0	34
Ratio: MUFA:PUFA	1:2.2	2:1
Ratio: n-6:n-3 FA	75:1	20:1

CHAPTER 3

RESULTS

3.1 BODY WEIGHT & BODY COMPOSITION

Body composition was calculated by DEXA and lean mass, percent lean mass, total body fat, and body fat percent were determined (Figure 3.1). The mice consuming the HFD had significantly elevated body fat and percent body fat relative to LFD-fed mice by 16 weeks of age (following 12 weeks of HFD feedings and prior to initiation of CRC), thus HFD-fed mice were obese prior to induction of CRC at 16 weeks of age ($P \leq .05$). While we observed a general increase in lean mass at sacrifice compared to baseline for all groups, the percent lean mass for HFD-fed mice was significantly reduced relative to the LFD-fed mice ($P \leq .05$). Conversely, body fat and percent body fat at sacrifice were significantly increased versus baseline in all the HFD groups. Interestingly, at sacrifice the LFD-CRC mice had significantly lower body fat and body fat percent than the LFD-Con, but the HFD-CRC-1 & 2 mice had significantly increased body fat and body fat percent compared to HFD-Con mice.

Body weight was monitored twice weekly throughout the 10 week AOM/DSS protocol and was expressed as a percent of baseline (i.e. normalized to measured body weight prior to the initiation of the AOM/DSS protocol) (Figure 3.2A). The LFD-Con and HFD-Con groups maintained a greater rate of body weight gain during the course of treatment and these groups were different from each other at most time-points. However, as expected, an overall decrease in body weight in the LFD-CRC group was observed

from weeks 17.5 to 22 and was significantly different from all other groups ($P \leq .05$). The HFD-CRC-1 & 2 groups had a significantly greater increase in percent change in body weight than the LFD-CRC group from 17.5 weeks to 22 weeks ($P \leq .05$).

3.2 SYMPTOM SEVERITY

Symptom scores were calculated twice weekly throughout the 10-week protocol and were based on diarrhea, blood in stools and body weight loss (Figure 3.2B).

Symptom scores in all groups followed the same general trend; there was an increase in symptom score during each week of DSS treatment, which dropped in the subsequent 2 weeks during the water cycle. The only significant increase in symptom severity score was observed in LFD-CRC mice compared to HFD-CRC-2 mice at week 18.5 ($P \leq .05$).

3.3 FOOD & FLUID INTAKE

It was not possible to calculate individual food intake, as mice were housed 4-5/cage. However, in general, we did not observe any differences among the HFD-fed mice in weekly food intake (i.e food consumed by mice in each cage/number of mice in cage) over the course of the study.

3.4 TISSUE WEIGHTS

Fat pads were collected at sacrifice and for all fat pad depots, the HFD-fed mice had enhanced fat pad mass compared to the LFD-fed mice (Figure 3.3) ($P \leq .05$). Both HFD-CRC-1 & 2 groups had a significantly greater epididymal fat pad mass than HFD-Con mice; however, the HFD-CRC-1 & 2 groups had reduced mesentery fat pads relative to HFD-Con.

Spleens were collected and weighed at sacrifice (Figure 3.4). In general, spleen weight, expressed as a percent of body weight, increased in all CRC groups as compared

to the control HFD and LFD mice and this reached significance when LFD-CRC mice were compared to HFD-Con mice ($P \leq .05$).

3.5 TUMOR COUNTS

At sacrifice, colons were harvested and tumors were counted on formalin-fixed, methylene blue-stained sections. No significant differences between groups were determined; however, the number of polyps in the LFD-CRC and HFD-CRC-2 groups was elevated compared to HFD-CRC-1 (Figure 3.6). Colon length and weight were measured following fixation. No significant differences were observed between groups for colon length and colon weight, (Figure 3.5).




Figure 3.1 DEXA body composition collected at baseline (16 weeks of age) and at sacrifice (26 weeks of age) on (A) lean mass and BMC, (B) body fat, (C) % lean mass and BMC, (D) % body fat. *Significantly different from LFD-Con ($P \leq 0.05$). #Significantly different from HFD-Con ($P \leq 0.05$). \$Significantly different from LFD-CRC ($P \leq 0.05$)




Figure 3.2 The impact of HFD and CRC on (A) percent change in body weight. *Significantly different from LFD-Con (week 17.5-18: HFD-Con & LFD-CRC; week 18.5-19: HFD-Con, LFD-CRC, & HFD-CRC-1 & 2; week 19.5-21.5: HFD-Con, LFD-CRC, & HFD-CRC-1; week 22-24.5: HFD-Con & LFD-CRC; week 25-26: HFD-Con, LFD-CRC, & HFD-CRC1&2; $P \leq .05$) #Significantly different from HFD-Con (week 16.5-17: LFD-CRC; week 17.5-26: LFD-CRC & HFD-CRC-1&2; $P \leq .05$). \$Significantly different from LFD-CRC (week 17.5-26: HFD-CRC-1&2; $P \leq .05$). (B) The effect of DSS induced inflammation on symptom scores. \$Significantly different from LFD-CRC (week 18.5: HFD-CRC-2 only; $P \leq .05$)




Figure 3.3 The effect of HFD and CRC on (A) epididymal, (B) mesentery, (C) kidney, and (D) total visceral (Total) fat pad weights at sacrifice.
 *Significantly different from LFD-Con ($P \leq .05$). #Significantly different from HFD-Con ($P \leq .05$). \$Significantly different from LFD-CRC ($P \leq .05$)




Figure 3.4 Spleen weight at sacrifice as a measure of inflammation and CRC immune response. #Significantly different from HFD-Con ($P \leq .05$)




Figure 3.5 The effect of CRC and DSS induced inflammation on (A) colon length and (B) colon weight. *Significantly different from LFD-Con ($P \leq 0.05$). #Significantly different from HFD-Con ($P \leq 0.05$). \$Significantly different from LFD-CRC ($P \leq 0.05$)




Figure 3.6 The impact of CRC induction at sacrifice for (A) total polyp number & (B) total polyp number by size. No significant differences between any groups




Figure 3.7 Mean (A) food consumption & (B) water consumption per group.

CHAPTER 4

DISCUSSION

CRC is the second leading cause of cancer related deaths worldwide and is a serious public health concern. Lifestyle factors, such as obesity, smoking, high caloric intake and physical inactivity, have been linked to increased mortality in CRC cases and are controllable factors in the prevention of CRC development and progression⁷. Clinical studies suggest that up to 20% of all cancers are attributed to obesity. Further, it has been well characterized that obesity is widely correlated with the incidence of CRC⁷. In addition, obesity is largely responsible for poorer oncologic outcomes in CRC treatment and management⁷. Our novel study examined the impact of HFD-induced obesity in a mouse model of CRC by initiating obesity prior to colon carcinogenesis and is thus, clinically relevant to the human obese population at risk for developing CRC. However, we did not find any significant changes in tumor number or size with obesity. Similarly, obesity did not increase the severity of symptoms in this model.

Currently, AOM/DSS-induced carcinogenesis in animal models is widely accepted and offers a reliable, reproducible, and time-effective approach for the study of colonic tumorigenesis⁷. Previous studies have utilized this model to investigate the link between obesity and colonic/colitis-related carcinogenesis. It is well established that adipocytes regulate inflammatory processes and secrete proinflammatory cytokines, which have been linked to increased proliferation of colonic epithelial cells and facilitation of a microenvironment prone to tumor development⁸. For example, Park et

al. showed that HFD induced obesity in a colitis-related tumorigenesis model increased tumor formation and revealed that adiposity-associated inflammation contributes to colonic epithelial cell carcinogenesis ⁹. In addition, the AOM/DSS model has been widely used to explore potential chemopreventive and therapeutic agents in the treatment of obesity and colitis-related colonic carcinogenesis ⁸.

In our study, male C57BL/6 mice were fed either a LFD (AIN-76A) or HFD (12% saturated fat and 28% unsaturated fat) at 4 weeks of age and placed in one of five groups (LFD-Control, HFD-Control, LFD-CRC, HFD-CRC-1, and HFD-CRC-2). The HFD was designed to be similar to the American standard diet; we previously demonstrated that this diet led to increased adiposity, macrophage infiltration, and insulin resistance in obese mice ¹⁰. As such, in our study, we observed that at 16 weeks of age the HFD-fed mice had significantly increased mean body mass compared to LFD-fed mice, thus were obese prior to administration of AOM/DSS treatment ($P \leq .05$). Enos et al. and others have shown that varying saturated fatty acid composition leads to an increased risk of obesity and can differentially regulate inflammatory processes, thus it is now widely accepted that high-fat-diet-induced obesity can lead to a chronic state of low-grade inflammation ¹⁰. Although inflammatory processes were not measured in the current investigation, we expected that obesity-induced inflammation following administration of this diet, would contribute to the hypothesized effects of increased tumorigenesis with obesity.

DSS, a proinflammatory agent, accelerates tumor growth and proliferation when instituted following an intraperitoneal injection of AOM by promoting cell growth in the epithelial lining of the colon while also halting apoptosis ⁷. Similarly, inflammation, as

seen in inflammatory bowel disease and Crohn's disease, also contribute to the development of CRC through various mechanisms that disrupt cellular function and regulation¹¹. We observed in the weeks following treatment with DSS that the sickness symptoms in all the AOM/DSS groups was lower than expected and remained diminished over the remainder of the study. This decrease in sickness symptoms over time is consistent with previous reports; in a comprehensive review by De Robertis et al., it was reported that severe colitis is observed in the week following initial administration of DSS but its severity is diminished in subsequent weeks⁷. Contrary to expectations though, we did not find an increase in symptom severity in the HFD groups. Similarly, at sacrifice, the colon length in the AOM/DSS treated groups did not decrease, suggesting little to no inflammation. A decrease in colon length is a characteristic feature observed in an inflamed colon, therefore these findings were unexpected. Further, tumor number was quite low and there were no differences in tumor size or tumor numbers with HFD. In a recent series of studies we conducted on colitis-related colon carcinogenesis, 2% DSS was administered in all three treatment cycles; however, a high number of fatalities prior to sacrifice limited the scope of the study. In this study, we subsequently reduced the dosage of DSS to 2%, 1%, and 1% at weeks 17, 20, and 23, respectively; however, contrary to what was expected, we found tumor numbers to be lower than those reported in the literature. These findings of a failure to find a detrimental effect of obesity on tumorigenesis in this model should be interpreted with caution; it is certainly possible that low tumor numbers precluded our ability to find changes in tumor characteristics across the groups.

A possible explanation for the discrepancy in inflammation and CRC tumor development may be the difference in the dosage of DSS. It has been established that the risk of developing CRC is positively correlated with the degree of inflammation within the colon and the duration of the inflammatory disease state. A DSS dosage study conducted with AOM-induced CRC carcinogenesis reported that 2% DSS treatment resulted in the highest number of colorectal adenocarcinomas¹². Similarly, it was also reported that inflammation scores were higher in mice receiving 2% DSS, suggesting there is a tumor-promoting activity in DSS¹². As inflammation is necessary in the induction of tumor formation, this may help to explain the minimal number of tumors observed in our model. In addition, it is worthwhile mentioning that the body weight and age of mice at time of AOM/DSS treatment were greater than those reported in previous models, which may have also contributed to the observed findings. Another possible rationale for the inconsistent tumor number and inflammatory symptom severity may be due to the extent of excess fat accumulation at the time of AOM/DSS treatment. While obesity is generally understood to have a proinflammatory effect in the colon, our results suggest that obesity may have some protective role in the initiation of CRC. This affirms the need for future research utilizing HFD-induced obesity models to better understand the relationship between obesity and colorectal tumorigenesis.

In summary, we examined the influence of HFD-induced obesity on inflammatory-mediated colorectal tumorigenesis, tumor proliferation, and symptom severity. In general, there was no increase in symptom severity with HFD-induced obesity. Similarly, in regard to tumor number, our findings are contradictory to previously published data that *ad libitum*, high-fat diet induced obesity leads to an

increased risk of CRC development. However, our findings should be interpreted with caution; it is certainly possible that low tumor numbers precluded our ability to find changes in tumor characteristics and symptom severity with obesity. Although obesity has implications of having proinflammatory and carcinogenic outcomes in the colon, future research should examine the degree to which HFD induced obesity impacts metabolic disruption, pro-inflammatory signaling, and subsequent initiation and proliferation of CRC as well as determine the pro-oncogenic factors involved in this process.

REFERENCES

1. Siegel R, Naishadham D, Jemal A. Cancer Statistics, 2013. *CA Cancer J Clin* 2013; 63:11-30.
2. Vucenik I, Stain J. Obesity and cancer risk: evidence, mechanisms and recommendations. *Ann N Y Acad Sci* 2012; 1271:37-43.
3. Teraoka N, Mutoh M, Takasa S, et al. High susceptibility to azoxymethan-induced colorectal carcinogenesis in obese KK-Ay mice. *Int J Cancer* 2011; 129:528-535.
4. Gribovskaja-Rupp I, Kosinski L, Ludwig K. Obesity and Colorectal Cancer. *Clin Colon Rectal Surg* 2011; 24:229-243.
5. Kanneganti M, Mino-Kenudson M, Mizoguchi E. Animal models of colitis-associated carcinogenesis. *J Biomed Biotechnol* 2011; 2011:1-22
6. Ma Y, Yang Y, Wang F et al. Obesity and risk of colorectal cancer: a systematic review of prospective studies. *PLoS ONE* 2013; 8:1-16.
7. De Robertis M, Massi E, Poeta M et al. The AOM/DSS murine model for the study of colon carcinogenesis: From pathways to diagnosis and therapy studies. *J Carinog* 2011; 10:9.
8. Tanaka T, Yasui Y, Ishigamori-Suzuki R et al. Citrus compounds inhibit inflammation and obesity-related colon carcinogenesis in mice. *Nutr Cancer* 2008; 60:70-80.
9. Park SY, Kim JS, Seo YK et al. Effects of diet-induced obesity on colitis-associated colon tumor formation in A/J mice. *Int J Obes* 2012; 36:273-280.
10. Enos RT, Davis JM, Velazquez KT et al. Influence of dietary saturated fat content on adiposity, macrophage behavior, inflammation, and metabolism: composition matters. *J Lipid Res* 2013; 54:152-63.
11. Suzuki R, Kohno H, Sugie S, et al. Strain differences in the susceptibility to azoxymethane and dextran sodium sulfate-induced colon carcinogenesis in mice. *Carcinogenesis* 2005; 27:162-169.

12. Tanaka T. Development of an inflammation-associated colorectal cancer model and its application for research on carcinogenesis and chemoprevention. *Int J Inflamm* 2012; 2012:1-16