

2016

Exploratory Analysis Of PTSD Severity And Objective Measures Of Physical Activity Among Combat Veterans

Danny O. Saucedo
University of South Carolina

Follow this and additional works at: <https://scholarcommons.sc.edu/etd>



Part of the [Exercise Science Commons](#)

Recommended Citation

Saucedo, D. O.(2016). *Exploratory Analysis Of PTSD Severity And Objective Measures Of Physical Activity Among Combat Veterans*. (Master's thesis). Retrieved from <https://scholarcommons.sc.edu/etd/3982>

This Open Access Thesis is brought to you by Scholar Commons. It has been accepted for inclusion in Theses and Dissertations by an authorized administrator of Scholar Commons. For more information, please contact digres@mailbox.sc.edu.

**EXPLORATORY ANALYSIS OF PTSD SEVERITY AND OBJECTIVE MEASURES
OF PHYSICAL ACTIVITY AMONG COMBAT VETERANS**

by

Danny O. Saucedo

Bachelor of Science
Stetson University, 2013

Submitted in Partial Fulfillment of the Requirements

For the Degree of Master of Science in

Exercise Science

Norman J. Arnold School of Public Health

University of South Carolina

2016

Accepted by:

Roger Newman-Norlund, Director of Thesis

Jack P. Ginsberg, Reader

Troy Herter, Reader

Lacy Ford, Senior Vice Provost and Dean of Graduate Studies

© Copyright by Danny O. Saucedo, 2016
All Rights Reserved.

DEDICATION

To my parents for doing more than enough to prepare me for a lifetime of opportunity. And to my brother, for lending an ear with love and support.

ACKNOWLEDGMENTS

I would first like to thank the Department of Exercise Science here at the University of South Carolina for their unwavering support throughout this process. I would like to sincerely thank the members of my thesis committee; Dr. Roger Newman-Norlund, Dr. Jack Ginsberg, and Dr. Troy Herter, for taking interest in my ideas and volunteering their time and expertise to help me become a better researcher. Roger, your enthusiasm, spontaneity and sense of purpose helped me pursue challenges I encountered during this study. I am better for it and give many thanks. Dr. Ginsberg, thank you for keeping me on track. Each day I walked out of your office I was inspired, not only by the Einstein quotes on your walls but, by the confidence you had in me. Thank you Dr. Herter for providing much needed guidance and having my back during this long process. I would also like to thank Dr. Steven Blair, Dr. Clemens Drenowatz, Leanna Ross and Sarah Schumacher for their generosity in providing activity monitors. Without these treasured items, this study would not have been possible. Thank you Clemens for the impromptu meetings and analytical support. I would also like to thank the following organizations for their hand in facilitating the recruitment process and taking interest in this initiative: William Jennings Bryan Dorn VA Medical Center, DAV 92nd Buffalo Chapter 20, SC National Guard, Navy-Marine Corps Relief Society, USC Student Veteran Association, and Experiment.com. To all my friends, thank you for understanding and encouragement along the way.

ABSTRACT

Strong levels of motivation, relatedness and self-determination support the likeliness of participating in habitual exercise. Compared to nationwide estimates, research has demonstrated fitness and performance is significantly lower among those suffering from PTSD. Meeting established recommended daily levels of physical activity (PA) for individuals with PTSD is critical for their physical and mental health status over a lifetime and factors supporting or hindering adherence to habitual exercise are lacking. Furthermore, regular PA among combat Veterans with Posttraumatic Stress Disorder (PTSD) is a viable and substantial treatment option to improve physical health status and cognitive well-being. The purpose of this exploratory analysis was to uncover relationships between objectively measured PA and the severity of PTSD symptomology, as established by DSM-V, among trauma-exposed combat Veterans. *Methods:* PA among a group of combat Veterans (n=10) was monitored for a period of 7-days concurrent to inventories of symptom severity, motives for participation in PA and posttraumatic growth (PTG). *Results:* Subjects reporting greater symptom severity and currently taking medication demonstrated inferior outcome results versus their counterparts. *Conclusion:* This was the first study, to our knowledge, to objectively explore PA among Veterans with PTSD. Mixed results suggest moderation of PTG and motives for PA on outcomes for those with PTSD. Activity monitoring in outpatient Veterans with PTSD is feasible. Future studies are warranted.

PREFACE

Since the power to create resides in each and everyone one of us, maybe focus should be directed toward individuals who persevere in their creativity despite the numerous barriers we encounter in society. This would give us a better understanding of resilient behavior and allow us to expedite progress and change we need to see in people, organizations and societies that hinder creativity or waste talent.

TABLE OF CONTENTS

DEDICATION	iii
ACKNOWLEDGEMENTS	iv
ABSTRACT	v
PREFACE	vi
LIST OF TABLES	viii
LIST OF FIGURES	ix
CHAPTER 1: INTRODUCTION AND AIMS	1
CHAPTER 2: REVIEW OF LITERATURE	2
2.1 PTSD, SYMPTOMOLOGY AND EXERCISE	2
2.2 DETERMINANTS OF EXERCISE MAINTENANCE	4
2.3 OBJECTIVE MEASUREMENT OF PHYSICAL ACTIVITY	6
CHAPTER 3: EXPLORATORY ANALYSIS OF PTSD SEVERITY AND OBJECTIVE MEASURES OF PHYSICAL ACTIVITY AMONG COMBAT VETERANS	8
3.1 RESEARCH QUESTION	8
3.2 RESEARCH STRATEGY	9
3.3 RESULTS	14
3.4 DISCUSSION	23
REFERENCES	32

LIST OF TABLES

TABLE 3.1 PARTICIPANT CHARACTERISTICS.....	15
TABLE 3.2 AVERAGE DAILY PHYSICAL ACTIVITY USING THE SWA UNDER FREE-LIVING CONDITIONS	15
TABLE 3.3 PEARSON PRODUCT-MOMENT CORRELATIONS OF SELF-RATED PA, PALMS AND MEASURES OF SWA ACTIVITY	17
TABLE 3.4 PEARSON PRODUCT-MOMENT CORRELATIONS AND FIRST-ORDER PARTIAL CORRELATIONS OF PCL-5 SEVERITY AND SWA ACTIVITY	19
TABLE 3.5 DESCRIPTIVE STATISTICS AMONG PARTICIPANTS STRATIFIED BY SYMPTOMS SEVERITY AND MEDICATION STATUS	20
TABLE 3.6 PEARSON PRODUCT-MOMENT CORRELATIONS OF POSTTRAUMATIC GROWTH WITH MEASURES OF SWA ACTIVITY AND SYMPTOM SEVERITY	21

LIST OF FIGURES

FIGURE 3.1 DAILY MVPA, PTSD SEVERITY AND MEDICATION STATUS.....	22
FIGURE 3.2 DAILY STEP COUNT, PTSD SEVERITY AND MEDICATION STATUS	22
FIGURE 3.3 DAILY SEDENTARY TIME, PTSD SEVERITY AND MEDICATION STATUS	23

CHAPTER 1

INTRODUCTION AND AIMS

The National Comorbidity Survey estimates approximately 7% of Americans can be expected to experience at least one traumatic event during a lifespan, roughly 8 million of which will develop a heterogeneous condition known as PTSD^[1]. The highest rates of PTSD are seen among women and are increasingly observed at higher rates among military service members due to combat exposure with estimates greater among Vietnam Veterans. The Diagnostic and Statistical Manual for Mental Disorders-fifth edition (DSM-V) has set forth the distinct addition of cluster symptoms that allows this proposed study to also satisfy calls for research into specific behavioral criteria of a four-cluster model of PTSD and determinants of long-term exercise maintenance^[2]. Now considered a growing epidemic, recent evidence supports a heritable component of trauma-exposure demonstrating the complexity of PTSD^[3,4]. Illuminating relationships between symptoms of PTSD and regular PA may support theories implicating heterogeneous psychiatric conditions and mood disturbances as challenges that negatively impact regular PA^[5,6]. Therefore, this study aimed for the following: 1) To collect objective and meaningful data concurrent to PTSD symptomology, as outlined by the DSM-V, among outpatient trauma-exposed combat Veterans, 2) To identify potential determinants of long-term maintenance of exercise or inactivity among this high interest population, and 3) To develop and apply reproducible practices for objectively assessing PA among populations with PTSD.

CHAPTER 2

REVIEW OF LITERATURE

2.1 PTSD, SYMPTOMOLOGY AND EXERCISE

The current Diagnostic and Statistical Manual of Mental Disorders (DSM-V) recognizes PTSD by four clusters of symptoms that adversely disrupt important areas of function due to a previous physical or psychological exposure to trauma or series of traumatic experiences. The four clusters include avoidance, persistence of negative moods and distorted cognitions, re-experiencing, and arousal symptomology, which distinguish key psychological and physiological behaviors exhibited by individuals with PTSD [7]. These behaviors guide interest to the neurobiological disruptions associated with fear circuitry, abnormal associative learning and maladaptive stress responses. Avoidance is a behavior in which people, places, conversations feelings or thoughts associated with a traumatic event is avoided. Persistent and distorted cognitions and negative emotions encompass symptoms related to diminished motivated behavior, exaggerated negative self-expectations or significant disinterest in activities such as social interaction. The arousal cluster of symptoms seeks aggressive or self-destructive behaviors, hyper-vigilance and sleep disturbances. Intrusions refer to the persistent and disruptive events that are specifically related to the exposure of the trauma. Re-experiencing also support sleep disturbances through distressing and recurrent dreams. Flashbacks, imagery, thoughts and/or perceptions are common and can be associated with persistent re-experiencing, or a form of reliving a traumatic episode [8,9,10].

The clusters of symptomology identified in PTSD characterize maladaptive neurobiology, still largely unknown, that involve learning, extinction, levels of arousal and marked sensitization of the hypothalamic-pituitary axis in response to stressors^[11]. These symptoms also identify areas that are subject to highly favorable adaptations with regular exercise among healthy and clinical populations^[12,13,14] but are scarce among individuals with PTSD. The generalizability of findings demonstrating exercise as a viable treatment for PTSD have limitations, primarily due to variance among sample populations studied (i.e., inpatient, mostly women, trauma type)^[15,16,17]. Those who suffer from PTSD are characterized as having poor health status and do not participate in sufficient levels of PA^[16, 17, 18] but exercise does bring about reductions in PTSD and depressive symptomology^[19]. Reductions in PTSD and depressive symptoms, in addition to health benefits associated with acute bouts of exercise, are short-lived and have been seen to reescalate in severity with physical inactivity^[15, 19].

Regular exercise can be characterized as engaging in volitional and habitual PA for at least 30-minutes a day, 5-days a week equal to or above 3.3 METS of energy expenditure^[20]. Furthermore, a period greater than six months is the final criteria that constitute an individual as achieving long-term exercise maintenance status so long as they remain in this phase^[21]. Anything short of this timeframe is considered to be the initiation of habitual exercise and therefore not the primary challenge the U.S. has seen regarding lifestyle behavior or adherence to exercise interventions^[22]. For consistency purposes, the term regular exercise will be align with criteria defined by the American College of Sports Medicine (ACSM) as well as the 2008 Physical Activity Guidelines defining the level of PA that elicits significant health benefits. In terms of improving

cardiorespiratory fitness with exercise prescription, the ASCM provides specific guidance on best practices for developing programs, framed by effective methods and theoretical models of behavior to facilitate adherence to exercise programs^[23, 24, 25]. To further support these evidence-based guidelines, the multi-organizational initiative headed by the ACSM, Exercise is Medicine (EIM), actively encourages and supports policy and programs that reduce physical inactivity and sedentary behavior^[26]. Despite the wealth of new knowledge and persistent efforts to increase nationwide PA levels, more than half of adults in the U.S. still do not meet recommendations for adequate levels PA^[19, 27, 28]. Regular exercise remains an unequivocal form of preventative and rehabilitative treatment that greatly reduces the development of risks tied to metabolic syndrome, cardiovascular disease, cancer and cognitive decline and should be objectively quantified^[29, 30, 31]. Thus, illuminating the relationship between PTSD symptomology and regular PA could provide value for interventions that support exercise adoption and maintenance.

2.2 DETERMINANTS OF EXERCISE MAINTENANCE

Exercise programs not only assess an individual's motivations and current life circumstances but also aim to address key determinants of exercise adoption and maintenance (e.g., social/environmental relatedness or support, self-efficacy and self-monitoring)^[32, 33]. Individualized theory-based cognitive-behavioral interventions are used to support such determinants in order to enhance adoption, but most importantly, adherence to long-term volitional exercise that meets recommended levels as described above. One of several commonly used approaches for exercise adoption and adherence is Social Cognitive Theory (SCT). This model equally monitors the individual, their habits and behaviors, as well as their physical, social and cultural environment^[22, 34]. An

appreciation for these factors generally motivates intentions, driving the energy for action, thereby directing an individual's future behavior. Although an individual's intrinsic motivation can encourage regular exercise habits, this alone is not sustainable. Self-regulatory strategies become essential for exercise adoption and maintenance therefore requiring an individual to regularly display skills like planning and organizing, while maintaining autonomy ^[35, 36, 37].

Central to several applied theories of behavior, is self-efficacy, which is considered to be one of the strongest predictors of adopting and maintaining regular exercise. Self-efficacy describes an individual's belief in his or her own capacity to reach a desired goal. Lacking self-efficacy contributes to stress, increases in physical and emotional arousal as well as depression ^[38]. People with greater exercise self-efficacy are most likely to engage in regular exercise, largely by the use of self-regulatory strategies such as goal setting, self-monitoring, and problem solving to enhance motivation ^[39]. Those who can maintain a high level of self-efficacy are likely to be more resilient to obstacles or inexperience allowing them to persevere on to a lifestyle that includes regular exercise ^[34, 40]. Therefore, characterizing cognitive-behaviors that influence these well-established determinants of long-term exercise maintenance may offer predictive value for favorable changes to self-regulatory strategies, health behaviors and beliefs to improve the likelihood of long-term volitional exercise.

Additionally, an individual's beliefs regarding the benefits of avoiding exercising or their susceptibility to illness are fundamental to a theory known as the Health Belief Model ^[41]. These beliefs are suggested to provide some insight into a person's readiness to act based on health concerns. There are increased concerns with older populations that

lack of social support and social isolation can be detrimental to long-term exercise maintenance ^[42, 43]. The risk of injury or worsening of current co-morbid conditions is also a significant belief, not only influencing self-efficacy but also diminishing autonomy. Habitual exercise can be seen among those with prior experience or within structured or supervised programs utilizing reinforcers or incentives ^[44]. Competency, social support and autonomy are significant theory-based components that emphasize a person's psychosocial needs, which ultimately favors exercise adoption and maintenance. Studies continue to define the complexity of barriers attributed to exercise adherence, most of which undermine key determinants of long-term exercise adherence ^[36, 45, 46].

2.3 OBJECTIVE MEASUREMENT OF PHYSICAL ACTIVITY

One study has evaluated the use of accelerometers among participants with PTSD in an attempt to validate assessments of PA in this population. Findings suggest measuring levels of PA among inpatient trauma-exposed adults may be impractical due to poor compliance, citing incomplete data, dropouts and erroneously high levels of activity ^[17]. A recent finding has shown that cardiorespiratory fitness is a confounder when estimating PA levels ^[47]. The use of a wearable monitor yields better results for unfit groups because they tend to overestimate PA levels using the international physical activity questionnaire (IPAQ) in comparison to fit participants. Furthermore, health promotion efforts suggest walking cadence can lead to positive health outcomes but it is unclear whether achieving this discrete criterion is critical for reducing mortality^[48]. Along with this, a significant oversight in estimating PA of varying intensities in healthy middle-aged to older adults may be attributed to initialization of actigraphy monitors to 1-second epoch lengths versus longer and more commonly seen settings of 60-seconds

that are better suited for sedentary or inactive populations ^[49]. Previous literature indicates the SenseWear armband provides more accurate estimates of total daily energy expenditure (TDEE), light and moderate intensities as compared to other activity monitors ^[31]. Lastly, it is unclear how best practices for activity monitoring have been applied with this sample population, therefore special attention to these guidelines was considered.

CHAPTER 3

EXPLORATORY ANALYSIS OF PTSD SEVERITY AND OBJECTIVE MEASURES OF PHYSICAL ACTIVITY AMONG COMBAT VETERANS

3.1 RESEARCH QUESTION

The goal of this study was to examine associations between objective measures of PA and PTSD symptomology among outpatient trauma-exposed combat Veterans, as outlined by the DSM-V. Given the lack of research on correlates of PA and PTSD severity, this pilot study aimed to serve as a platform for further investigation into the behavior and underlying motives for exercise adherence or contradictory lifestyle behaviors. The second goal was to demonstrate feasibility of activity monitoring among those with PTSD. This study anticipated the advancement of three areas of interest surrounding mental illness and adherence to long-term exercise: 1) The collection of objective and meaningful data concurrent to symptom severity among a high interest population. 2) The identification and evaluation of potential correlates of long-term maintenance of exercise or inactivity among trauma-exposed combat Veterans. 3) The development and application of reproducible practices for objectively assessing PA among populations with psychiatric distress or illness. Each variable resides in 1 of 3 descriptive categories: criterion for PTSD (*qualitative*), predictors of symptom severity (*quantitative*) and moderators (*qualitative*) that influence PA levels (i.e., posttraumatic growth, self-determination status). We hypothesized that PTSD severity would be

inversely related to most outcomes (e.g., step count, MVPA, sleep duration) in accordance with multiple prior studies^[16, 19, 50]. MVPA among this sample was anticipated to be lower than healthy adult populations based on prior results regarding anxiety disorders and patients with SMI^[51,52]. Moderating factors were hypothesized to explain some of the variance among positive associations between PTSD severity and measures of PA. The assumption is if positive levels of PA are observed among individuals showing severe PTSD, this may be an indication of elevated levels of intrinsic motivation. This perhaps, may also be driven by or at least correlated with levels of psychological growth experienced by individuals as a result of trauma-exposure. Lastly, we expected to observe significant issues related to compliance, as in prior research assessing PA in PTSD^[17, 19].

It can be said that Veterans with PTSD have a set of past experiences that may be more conducive to positive psychological changes/growth than the rest of the population^[53]. Previous research suggests at least half of U.S. Veterans, independent of PTSD status, experience moderate levels of posttraumatic growth (PTG)^[10]. Because emotions thoughts and behaviors are interconnected, additional qualitative measures were administered with the intent of ultimately providing useful direction for the staging of interventions in larger scale studies and identify determinants of exercise adherence in this population.

3.2 RESEARCH STRATEGY

Study Design. A mixed methods approach was adopted for this cross-sectional study in order to explore relationships between objective measures of physical activity and PTSD symptomology among trauma-exposed combat Veterans. Height (cm), body

weight (kg), demographics and diagnostic surveys were employed to classify general characteristics and patterns of behavior among participants. Physical activity was then objectively measured for a period of 7 days for evaluation.

Participant Characteristics. Study participants (n=10) were recruited from within the local community of Columbia, South Carolina from September 2015 to January 2016. Eligibility mirrored preceding studies examining populations with PTSD in conjunction with physical performance measures and was limited to men and women between the ages of 18-65 [16, 17, 54]. The University of South Carolina Internal Review Board approved the study. Written informed consent was acquired from each subject prior to data collection.

Assessments. In response to a recruitment flyers posted at William Jennings Bryan Dorn VA Medical Center and other locations throughout the local community. Volunteers were directed to an online pre-screening tool to establish participant eligibility prior to their initial visit. Participants then reported to the Arnold School of Public Health for orientation after meeting inclusion criteria. All subjects were required to have been deployed or part of combat operations. Exclusion criteria were any history of neurological disorders, cardiovascular disease to include the use of pacemakers, heart surgery or transplants, depression, alcohol or substance abuse/dependence, and current antipsychotic pharmacotherapy. After acquiring written consent, demographic information and health history items were completed. Participants also self-reported their level of fitness rating ranging from ‘very poor’ to ‘regular physical activity for a period of 6 months or more’. This information was used to gauge if participants maintained pre-study PA levels throughout the assessment period as directed. Study-specific inventories

were administered in the order encountered in this report. All necessary items were reviewed and endorsed by a licensed psychiatrist.

The *Neurobehavioral Symptom Inventory* (NSI), a 10-question inventory, was administered to ensure volunteers who may have had existing concussive or traumatic brain injury symptoms were excluded from the study^[55]. Other exclusion criteria included having a disease that limited PA.

The *Psychometric Properties of PTSD Checklist For DSM-V* (PCL-5) is a 20-item self-report measure of diagnostic symptomology for PTSD. Responses indicated the extent to which subjects had been bothered by PTSD symptom clusters within the past month. A PCL-5 severity score of 25 or greater was set to maximize detection of sub-threshold or partial PTSD, in addition to full PTSD symptomology^[56]. For the purpose of this study, scores greater than 25 reflected a condition worse than those without PTSD. All subjects were identified as having experienced at least one physically or emotionally traumatic-life event in connection with their past military deployments. A strong correlation exists between the PCL-5 and the Clinician-administered PTSD Scale, the gold standard for clinical diagnosis^[8, 57].

The *Posttraumatic Growth Inventory-Short Form* (PTGI-SF) is a 10-item self-report qualitative measure, validated to assess positive psychological changes as a result of traumatic experiences^[58]. This inventory provides insight that may account for positive lifestyle changes or behaviors supporting exercise adherence despite potential contradictory scores on PCL-5. In a web-based survey involving more than two thousand US Veterans, approximately 75 percent of respondents who screened positive for PTSD, demonstrated at least moderate levels of posttraumatic growth (PTG)^[10]. Veterans with

PTSD who demonstrated PTG in this study, also mentioned having improvements in mental clarity and well-being in comparison to those not indicating PTG. These findings may correlate with psychosocial factors such as personal strength, social connectedness and purpose in life, which are necessary components implicated in cognitive-behavioral paradigms that facilitate exercise adoption and maintenance. For the purpose of this study, only total score was used for correlation per recommendations ^[10] and a cut-off score of 20 indicated moderate levels of PTG.

The *Physical Activity And Leisure Motivation Scale* (PALMS) is a 40-item assessment made on a 5-point Likert scale. It examines a person's level of intrinsic or extrinsic motivation for participating in PA and has been determined to be a valid and reliable application for research. The PALMS was developed to serve as a short form for the 73-item Recreational Exercise Motivation Measure. Each of the eight motivational sub-scales in the PALMS has implications underlying intent and behaviors related to exercise ^[59]. Each sub-scale of the PALMS has five items that align with motivational constructs of self-determination theory (SDT). Mastery and enjoyment sub-scales align with intrinsic motivation while the psychological condition, physical condition and appearance sub-scales align with shared intrinsic and extrinsic motives. The last three sub-scale items, expectations, affiliation and competition/ego are representative of extrinsic social motives.

Physical Activity. Participants were affixed with the SenseWear Armband (SWA) (BodyMedia Inc., Pittsburgh, PA, USA) for a period of 7-days to summarize a variety of distinct patterns in their motion, skin temperature, heat dissipation and galvanic skin responses to estimate total daily energy expenditure (TDEE) and PA levels. The armband

is worn on the posterior surface of the upper left extremity and records these data via a tri-axial accelerometer, electrical conductivity of the skin and an electronic thermometer. The SWA is a well-researched and validated method for assessing TDEE and exercise intensity^[31, 60, 61]. Other outcome measures included were wear-time, sleep duration, step count, sedentary-time (≤ 1.5 METs) and moderate-to-vigorous physical activity (MVPA ≥ 3.0 METS). A 10,000 steps/day threshold was used in our evaluation based on criteria that support brain health and adequate PA levels^[62, 63]. Only one-prior study has evaluated the use of accelerometers among participants with PTSD in an attempt to validate self-reported PA assessments in this population^[17]. Their findings suggest measuring PA levels among inpatient trauma-exposed adults may be impractical due to poor compliance, citing incomplete data, dropouts and erroneously high levels of activity. For this pilot study, each participant was provided clear instruction on armband usage and criteria for wear and non-wear time. Valid wear time was set to at least 7 consecutive days, including a weekend period, for ≥ 21 hours each day. Initialization, distribution, device tracking, epoch timing and reporting adhered to guidelines specific to the use of the SWA in recent studies involving healthy free-living adults^[49, 61]. Efforts to enhance compliance to the study protocol followed all four strategies outlined by best practices set forth by recent literature regarding activity monitors in research^[31, 64, 65]. SWA metadata was processed using the SenseWear Professional Software version 8.0 (algorithm v2.2.4).

Analytical Procedure. Statistical analyses were performed using SPSS (version 22.0; IBM Corp., Armonk, NY, USA). Pearson product moment correlations and first and second-order partial correlations were performed among criterion, predictors and

covariates to demonstrate associations among variables. Statistical significance was set at $p < .05$ (one-sided) for all analyses.

3.3 RESULTS

Participant Characteristics. Table 3.1 shows characteristics of combat Veterans that were a part of this study. Mean age of the participants was 49 ± 12 years. The racial/ethnic make-up of this group was predominantly male (80%), female (20%). The average height and weight of the participants were 176 ± 11 cm and 95.5 ± 17.4 kg, respectively. The average subject BMI was 31 ± 6 for all subjects, classifying 70% of participants as obese. Prior to enrollment, only two participants self-reported as being ‘physically inactive’ on most days of the week. Self-rated PA levels were consistent with data retrieved from the SWA for all subjects. Only one participant in this study was retired. This participant was heavily active with volunteer work within the community. All other participants were employed or attending college.

Physical Activity. Outcomes of PA using the SWA were measured over a 7-day period, including a weekend, and are summarized in Table 3.2. Average wear time for this sample was remarkable. Approximately 22.9 hours per subject was collected from all participants, a 100% rate of compliance based on valid wear-time criteria. Analyses involving measures taken during weekdays had nearly identical average wear time and compliance as weekends. Any missing metadata during transfer of SWA activity were interpolated using predictive mean matching for the 7-day period, which comprised 6.04% of the data. The coefficient of variation (CV) was included for all outcomes of activity for an appropriate comparison. The SWA total and weekday activity revealed,

Table 3.1 Participant characteristics.

	Total (n)
Male	8
Female	2
Age (years)	49 ± 12
Body Mass Index (BMI)	31 ± 6
Healthy (18.5 – 24.9)	3
Obese Class 1 (30.0 – 34.9)	3
Obese Class 2 (35.0 – 39.9)	4
Education	
< College degree	4
College degree	2
Graduate	4
Employment	
Student	3
Employed	6
Retired	1
Self-rated Fitness Level	
Physically Inactive	2
Occasionally active (≥ 30 min. of PA 3 days/week)	3
Regular PA for more than 6 months	5
Medication (SSRIs only)	3

PA, Physical Activity; SSRI, Selective-Serotonin Reuptake Inhibitor; Age and BMI values are mean ± SD.

Table 3.2 Average daily physical activity using the SWA under free-living conditions.

	Total (n=10)		Weekdays		Weekends	
	Mean	CV	Mean	CV	Mean	CV
Wear time	1373	.04	1374	.05	1369	.08
Sleep duration	380	.20	369	.24	405	.21
Sedentary	903	.22	1011	.20	634	.41
MVPA	95	.68	99	.64	85	.92
Step count	7284	.46	7684	.40	6282	.78
TDEE (kcal/day)	2962	.16	2980	.16	2916	.19

Mean values given in minutes unless otherwise indicated; Total activity column collected Monday through Sunday; CV, Coefficient of Variation; SWA, SenseWear Armband; MVPA, Moderate-to-Vigorous Physical Activity; TDEE, Total Daily Energy Expenditure.

nearly equivalent dispersion across all variables, despite increased variability on weekends. As anticipated average sedentary time on weekdays was 16.9 hours (CV of 20%) while average sedentary time on weekends was reduced to 10.6 hours (CV of 41%).

Differences in average daily step count for weekends and weekdays are most likely attributed to higher variability among participants (CV of 78%) on weekends. Average MVPA was 1.6 hours (CV of 68%) for all 7 days and 1.4 hours (CV of 92%) during weekends. Average TDEE for all participants was similar on weekdays and weekends reaching 2980 kcal/day (CV of 16%) and 2916 kcal/day (CV of 19%), respectively. No erroneously high values were observed among individual SWA activity.

Motives for Participation in PA. Prior to data collection, current self-report levels of PA and the PALMS scores were assessed. In Table 3.3, bivariate correlations between self-rated PA responses, the PALMS and each of the eight sub-scales were evaluated for any disagreement suggesting participants altered their exercise habits due to study participation. PALMS total score and self-rated responses correlated well with TDEE, sedentary-time and MVPA activity. Subject responses for self-rated PA strongly correlated with TDEE ($r = 0.80, p < .05$), step count TDEE ($r = 0.66, p < .05$) and MVPA ($r = 0.68, p < .05$). Examination of PALMS responses only indicated a significant association with MVPA ($r = 0.65, p < .05$), yet moderate associations were seen with TDEE ($r = 0.49$) and step count ($r = .57$). Additionally, MVPA was positively related to extrinsic mind-body motives, psychological condition ($r = 0.67, p < .05$) and physical condition ($r = 0.81, p < .05$), with similar strong associations with step count. Physical condition was strongly associated with TDEE ($r = 0.74, p < .05$) and moderately inversely related to sedentary time ($r = -0.57, p < .08$). MVPA and step count strongly correlated with intrinsic motives (e.g., mastery, enjoyment) for participating in PA. These results specifically indicated significant associations between step count and enjoyment ($r = 0.66, p < .05$) and MVPA and mastery ($r = 0.68, p < .05$). As anticipated, a moderate

inverse relation between sedentary time and MVPA was also observed. Overall, extrinsic social motives (e.g., others' expectations, competition/ego) were seen to have an inverse relation with positive outcomes associated with PA, but not with affiliation. The affiliation PALMS sub-scale was shown to have a moderate association with MVPA ($r = 0.53, p = .11$). Finally, step-count demonstrated significant associations with each of the extrinsic mind-body motives indicating physical condition and psychological well-being as motives for participating in walking among this sample. *PTSD Symptom Severity*. All study participants were combat Veterans who self-reported as experiencing at least one previous traumatic event during a deployment, therefore each member was administered the PCL-5. Average PCL-5 score was 22 (range: 2-39). Table 3.4 summarizes PCL-5 symptom severity and SWA activity using bivariate correlations. Second and third-order

Table 3.3 Pearson product-moment correlations of self-rated PA, PALMS and measures of SWA activity.

(n=10)	TDEE	Sleep Duration	Sedentary	Step Count	MVPA
Self-rated PA	.69*	.07	-.43	.66*	.76*
PALMS					
Total Score	.49	-.01	-.41	.57	.69*
Mastery (μ)	.43	-.01	-.34	.54	.68*
Enjoyment (μ)	.41	-.25	-.58	.66*	.53
Psychological Condition (μ)	.39	-.04	-.40	.75*	.67*
Physical Condition (μ)	.68*	.21	-.46	.77*	.83*
Appearance (μ)	.49	.21	-.20	.45	.60
Others' Expectations (μ)	.10	.20	.18	-.38	-.22
Affiliation (μ)	.12	-.24	-.22	.24	.47
Competition/Ego (μ)	-.38	-.17	.11	-.63	-.37

SWA, SenseWear Armband; TDEE, Total Daily Energy Expenditure; MVPA, Moderate-to-Vigorous Physical Activity; PALMS, Physical Activity and Leisure Motivation Scale

* $p < 0.05$

partial correlations were also performed to evaluate known covariates (i.e., body weight, gender) and medication status as a mediator, also seen in Table 3.4. Medication status appears to account for the relation between several criterion variables and measures of PA. Although not statistically significant, a moderate positive association was indicated between PCL-5 scores and TDEE ($r = 0.54, p = .11$), sleep duration ($r = 0.39, p = .11$) and MVPA ($r = 0.55, p = .10$) when controlling for medication status. Intrusions symptomology (cluster B) demonstrated an identical association with TDEE ($r = 0.56, p = .08$) and MVPA ($r = 0.50, p = .14$) while adjusting for medication status. A moderate inverse association was observed between avoidance symptomology (cluster C) and TDEE variability ($r = 0.43, p = .21$) and sleep duration variability ($r = 0.56, p = .08$). Avoidance maintained this moderate association with TDEE variability and sleep duration variability while adjusting for sex, body weight and medication status. Significant positive associations between arousal symptomology (cluster E) and TDEE ($r = 0.63, p < .05$) and MVPA ($r = 0.74, p < .01$) were seen when controlling for medication. A moderate positive association was also seen with arousal and MVPA ($r = 0.42, p = .22$) when controlling for body weight. Lastly, PTSD symptom severity above and below the cut-off score of 25 was correlated with all outcomes to explore any additional interactions. Correlations with cut-off score and TDEE ($r = 0.40, p = .25$), sleep duration ($r = 0.46, p = .18$) and MVPA ($r = 0.35, p = .32$) revealed moderate associations while controlling for medication although negatively associated with step count ($r = -0.35, p = .32$).

Medication Status and Physical Activity. The strong positive association of symptom severity with MVPA, sleep duration and sedentary time, adjusting for

medication, was explored further. Viewing medication as a mediator, the members of this study were stratified into three groups. PCL-5 total scores < 25 were considered subclinical or partial ($n=4$) and scores ≥ 25 with greater symptom severity were split into two groups based on medication status (Table 3.5). Subjects currently on prescribed medication spent 32 minutes in MVPA, 5.6 hours sleeping and 16 hours sedentary on average each day. However, subjects off medication spent 2.7 hours in MVPA, 7.2 hours sleeping and 14.3 hours sedentary (Figure 3.1).

Posttraumatic Growth. Bivariate correlations among PTG total scores in Table 3.6 revealed a significant and inverse association with sleep duration and a moderate association with PCL-5 scores. Adjusting for medication, PTG was seen to have a moderate association with MVPA after controlling for medication and step count whereas unadjusted correlations were inversely related. Among those reporting greater symptom severity receiving pharmacotherapy, PTG scores were highest (Range: 21-40) while PALMS scores were the lowest (Range: 64-121). PCL-5 scores ≥ 25 and not taking medication reported the highest PALMS scores (Range: 134-139) and lower levels of PTG (Range: 15-36). Cronbach's $\alpha = .90, .84$ and $.91$ for PCL-5, PTGI-SF and PALMS inventories respectively. Figure 3.1 shows average daily MVPA among each group as measured by the SWA for a 5-day period. On average, those currently taking medication did not achieve recommended levels of daily MVPA on 2-of-5 days. All other participants achieved sufficient levels of MVPA on average during the week. Those reporting greater symptom severity in Figure 3.2, demonstrated comparable steps/day throughout the 5-day period with respect to those demonstrating minor to no PTSD symptoms.

Table 3.4 Pearson product-moment correlations and first-order partial correlations of PCL-5 severity and SWA activity.

(n=10)	TDEE				Sleep Duration				Sedentary				Step Count				MVPA			
	r	r _b	r _m	CV	r	r _b	r _m	CV	r	r _b	r _m	CV	r	r _b	r _m	CV	r	r _b	r _m	CV
PCL-5																				
Total score	.01	-.06	.54	-.30	.12	.03	.39	.23	.07	-.10	-.04	-.04	-.45	-.37	-.20	-.25	.02	.13	.55	.13
Intrusions (μ)	-.16	-.26	.56	-.24	-.10	-.25	.24	.21	.18	-.00	.05	-.07	-.52	-.42	-.14	-.22	-.19	-.08	.50	.26
Avoidance (μ)	-.27	-.34	.07	-.43	-.15	-.25	.03	.56	.30	.23	.24	-.31	-.56	-.55	-.38	-.30	-.16	-.09	.24	.25
Neg. mood cognition (μ)	.04	-.05	.50	-.33	.29	.18	.54	.35	.18	-.02	.10	-.08	-.51	-.38	-.34	-.04	-.06	.09	.35	.30
Arousal (μ)	.28	.27	.63	-.33	.22	.20	.35	.14	-.29	-.39	-.36	.19	-.12	-.10	.05	-.40	.37	.42	.75*	-.26
PCL-5 Stratified scores																				
(cut point ≤ 25)	.14	.07	.40	-.15	.34	.25	.46	.10	.17	-.08	.13	-.16	-.40	-.27	-.35	.06	.10	.25	.35	.19

Partial correlations r_b and r_m controlling for body weight and medication, respectively; PCL-5, Psychometric properties of PTSD Checklist for DSM-V; TDEE, Total Daily Energy Expenditure; MVPA, Moderate-to-Vigorous Physical Activity; CV, Coefficient of Variation

* p < 0.05.

Table 3.5 Descriptive statistics among participants stratified by symptom severity and medication status.

	Total (n=10)		PCL-5 < 25 (n=4)		PCL-5 ≥ 25 (n=3)		PCL-5 ≥ 25 on medication (n=3)	
	Mean	CV	Mean	CV	Mean	CV	Mean	CV
MVPA	95	.68	91	.36	164	.29	32	1.34
Step count	7284	.46	8828	.37	8365	.28	4143	.71
Sedentary	903	.22	889	.20	863	.36	962	.14
Sleep Duration	380	.20	371	.24	434	.11	337	.20
TDEE	2962	.16	2990	.09	3387	.16	2500	.04
PTGI Score	23	.51	16.5	.71	23	.48	32	.31
PALMS Score	114	.26	117	.25	137	.02	88	.33

SWA, SenseWear Armband; TDEE, Total Daily Energy Expenditure (kcal); MVPA, Moderate-to-Vigorous Physical Activity; PTGI, posttraumatic growth inventory; PALMS, Physical Activity and Leisure Motivation Scale; CV, Coefficient of Variation

Longer durations spent in sedentary time were seen during weekdays among those with higher PCL-5 scores, but observed to be especially high among Veterans on medication (Figure 3.3).

Table 3.6 Pearson product-moment correlations of posttraumatic growth with measures of SWA activity and symptom severity.

(n=10)	TDEE		Sleep Duration		Sedentary		Step Count		MVPA		PCL-5	
	r	r _m	r	r _m	r	r _m	r	r _m	r	r _m	r	r _m
PTGI												
Total Score	-.24	.16	-.69*	-.62	-.14	-.29	-.12	.31	-.03	.49	.44	.25

SWA, SenseWear Armband; TDEE, Total Daily Energy Expenditure; MVPA, Moderate-to-Vigorous Physical Activity; PTGI, Posttraumatic Growth Inventory; first-order partial correlation controlling for medication (r_m)

* p < 0.05




Figure 3.1 average daily times in moderate-to-vigorous physical activity (MVPA) as recorded by the SWA; Subjects contributing to these averages were stratified by PCL-5 cut-point and by current medication status.




Figure 3.2 Average daily steps as recorded by the SWA; Subjects contributing to these averages were stratified by PCL-5 cut-point and by current medication status.




Figure 3.3 Average daily times spent in sedentary behavior as recorded by the SWA; Subjects contributing to these averages were stratified by PCL-5 cut-point and by current medication status.

3.4 DISCUSSION

In the present paper we have discussed the importance of employing wearable monitors to reliably appraise potential correlates of PA in individuals with PTSD. Ten trauma-exposed combat Veterans were monitored for a period of 7-days to assess voluntary activity levels during free-living conditions. Qualitative surveys regarding personal motives for participating in PA and positive psychological changes as a result of prior trauma-exposure were explored in connection with activity. Evidence from this mixed methods approach arguably provided useful direction for the staging of future study interventions regarding current pharmacotherapy status and trends in PA for this population. Assumptions regarding PTG and motives for PA may also offer significant utility in the development of strategies for adhering to a physically active lifestyle.

Exceptional study compliance among this outpatient sample allowed us to underline four key findings that deserve further consideration.

Firstly, Veterans currently taking medication (i.e., antidepressants) and report greater severity of PTSD symptomology, display inferior levels of PA as compared to their counterparts in this sample. Current first line monotherapeutic strategies for PTSD include strong recommendations for the use of antidepressants such as selective serotonin reuptake inhibitors (SSRIs) ^[66]. The Clinical Practice Guidelines for Posttraumatic Stress Disorder recommend against atypical antipsychotic use for the monotherapeutic management of PTSD and were therefore held within exclusion criteria. Antipsychotics are known produce deleterious effects among patients with psychiatric disorders, such as weight gain and metabolic disturbances ^[67]. Even so, study participants currently on SSRIs exhibited similar body weight characteristics. Our results highlight a number of risk factors attributed cardiovascular disease and metabolic health commonly seen in Veterans with PTSD (e.g., BMI, sedentary time, inactivity). SMI status alone is generally connected to greater risk for mortality and lower levels of PA than the general population ^[68]. More specifically, PTSD has been attributed to poor functional status, impaired physical performance and less walking behavior ^[16, 50]. The strong inverse association between medication and outcomes of PA for subjects with higher PCL-5 scores (≥ 25) suggests a greater potential for functional impairment and individual distress for this group. PCL-5 scores maintained a moderate inverse association with step count even when controlling for body weight and gender. PCL-5 symptoms clusters each shared this same moderate inverse association with step count (Table 3.4). The strength of these associations was reduced after controlling for medication revealing prescription status as

a potential mediator. Again adjusting for medication, MVPA was only then observed to have a strong positive association with total PCL-5 scores. As expected, this association was seen in proportion to estimated caloric expenditure demonstrating the consistency of increased activity on TDEE ($r^2 = .85$). Overall, this would indicate higher levels of MVPA observed among those with greater symptom severity adjusting for medication in this sample. This evidence is fittingly compatible with a set of previously proposed hypotheses implicating SSRIs as agents that may reduce the likelihood of engaging in PA via executive dysfunction^[69].

Secondly, participants of this study who were not on any pharmacological treatments on average exceeded recommended levels of daily PA, despite displaying few steps/day on average than those indicating lesser symptom severity and significant sedentary time. Most participants of this study did not meet recommendations for 10,000 steps/day, especially participants taking medication. Among healthy populations, the relationship between steps/day and sedentary time is inverse and moderate^[70]. Sedentary behavior was remarkably high in this sample. The association with steps/day was strong and inversely ($r^2 = .68$) related for this sample and also held true for MVPA and time in sedentary behavior ($r^2 = .43$). Risk of mortality is not lessened among those who experience significant psychological distress and participate in high levels of MVPA (> 7 hours/wk) yet spend 7 hours/day in sedentary behavior^[71]. These data underline the importance of uncovering correlates of cognitive, social and environmental barriers among this population to reduce time in sedentary activity.

Thirdly, in an effort to explain higher levels of MVPA among those with greater symptom severity, data were stratified into three groups after first order partial

correlations (Table 3.5). It was hypothesized that individuals who reported greater PTSD severity, but displayed at least moderate levels of PTG and/or higher PALMS scores, would exhibit better outcome measures of activity than those with lower PCL-5 scores. Stronger levels of intrinsic and mind-body motivation, as measured by the PALMS, were seen among those with higher levels of MVPA and step count. PTG levels also exhibited moderating characteristics, which may offer predictive value in relation to symptom severity. If PTG is interpreted as having a curvilinear relationship with symptom severity [53], levels of PA may correlate similarly using a similar model. If so, it would be interesting to see if and how PTG influences individual perceptions about adequate levels of PA or constructs contained within the Health Belief Model.

Our last key finding points to the association between sleep duration and the arousal cluster of PTSD symptomology. Subjects with higher PCL-5 scores currently on pharmacotherapy demonstrated lower average sleep duration than those of equivalent symptom severity not taking medication, 5.5 ± 1.1 hours and $7.2 \pm .8$ hours respectively. Incidentally, this group ($PCL \geq 25$ off medication) had greater levels of MVPA and comparable sleep duration as compared to those experiencing few to no symptoms, yet arousal among this sample revealed a significant positive association with PTSD severity. A previous study ($n=108$) with similar direction identified an inverse association between self-reported vigorous-intensity PA and arousal symptomology, despite low severity scores employing a *DSM-IV* diagnostic scale [19]. Previous research has also shown that individuals with high psychological distress are susceptible to sleeping less than six hours or sleeping greater than nine hours each night [72]. Those perceived to have a favorable health status in this study were seen to experience lower levels of psychological stress,

were of normal BMI and commonly seen to sleep between 7-8 hours. PCL-5 scores and arousal symptomology demonstrated positive associations with sleep in our current study, inconsistent with previous literature demonstrating reductions in arousal symptomology among inpatients with PTSD after a cycling intervention ^[73]. Further attention to this distinction is worth pursuing with objective monitoring among inpatient and outpatient samples to define and interpret the therapeutic effects of exercise at various intensities.

Strengths. Study compliance to activity monitoring among this outpatient sample was remarkably high, as opposed to unfavorable rates seen in inpatient populations ^[17]. Regarding study flow, validated versions of short form inventories were selected purposefully wherever possible to ensure readability and increase retention. The estimated reliability of test scores reported in this study were good to excellent as indicated by Cronbach's α ($\geq .84$). Requests to adhere to pre-study routines of PA were followed as supported by PALMS and self-rated PA. There were no losses to report in terms of activity monitors. All together, future studies aiming to objectively monitor PA among outpatient trauma-exposed Veterans should be feasible using well-established protocols.

Limitations. Results from this study should be interpreted in consideration of several limitations. The power of this study to show significant effects was low and has not sufficiently satisfied previous calls for objectively measured activity among this sample population. Future studies should be repeated with larger sample sizes to cross-validate PTG and motives for participating in PA as moderator variables. Our data are cross-sectional, so it is impossible to determine whether any observed cognitive dysfunctions represent causality among the correlates. Another limitation to report is the

lack of a control group. It is important to distinguish whether prior military training could account for the high levels of MVPA and motivation seen among those with PCL-5 scores greater than 25 not on medication. Although participants were screened for conditions such as depression and musculoskeletal limitations, a formal comorbidity index is recommended to promote full disclosure and account for any overlapping symptoms.

Future studies. Prospective studies pre and post-intervention are needed to adequately develop and stage programs for trauma-exposed combat Veterans. Given the high prevalence of walking as the only form of PA among those with SMI, parameters such as cadence should be observed. Such details would provide a more accurate reflection of activity to better understand dose-response parameters. The moderate and positive relationship between avoidance symptomology with most outcomes of activity is worth mentioning. These associations could be interpreted as potential markers of participant reactivity or altered behavior toward demand characteristics of study participation. Therefore, extended periods of observation greater than 7 days are recommended. Moreover, because of the moderate and reciprocal associations of avoidance with step count and MVPA variability, it would be worthwhile to determine if measures of behavioral inhibition, or the tendency to withdraw from novel or unfamiliar social or non-social situations, correlate well with PA. Behavioral inhibition has been shown to have a positive association with PTSD severity^[74]. A fundamental self-awareness of PA behavior, facilitated by activity monitors or social stimulation via programs, is suggested to influence health outcomes^[75], and therapeutic responses can be non-specific^[76]. Trauma-exposed combat Veterans may benefit from programs that issue

wearable technologies periodically as part treatment to evaluate and encourage positive behavioral changes.

To create a formidable exercise intervention for trauma-exposed individuals experiencing psychological distress, further investigation into the neurophysiology and behavior underlying motives for regular exercise or contradictory lifestyle behaviors are needed. Although qualitative research involving psychosocial constructs, such as self-determination, have been conducted in relation to PA levels among SMI populations [33, 44, 52], the concurrent use of objective measures have been scarce. Lack of social support appears to be one of several important risk factors that hinder adherence to exercise. The affiliation sub-scale of the PALMS demonstrated a moderate correlation with MVPA among this population, as did PTG. Qualitative research may find utility for the PTGI-SF and PALMS to identify positive psychosocial factors in trauma-exposed combat Veterans. Findings may illustrate the manner in which some individuals with PTSD desire to live their lives and how that may relate to habitual exercise.

Conclusion. This exploratory analysis was the first to examine associations between objective measures of PA and PTSD symptomology, as outlined by the DSM-V, among outpatient trauma-exposed combat Veterans. This preliminary study has demonstrated the feasibility of assessing activity levels among this high interest population as supported by study compliance and the application of reproducible methodology. Using a mixed-methods approach, objective and meaningful data concurrent to symptom severity has uncovered a few considerations that should be investigated in larger scale studies. Despite our small sample, medication was seen to negatively impact outcome measures of PA. Moreover, results from this study exhibit

some alignment with our hypothesis suggesting PTG and motives for participation in PA may have predictive value. Understanding the relationship between PTG and PA among Veterans with PTSD still needs further investigation. Despite the positive outlook newly acquired as a result of trauma-exposure, the relationship between PTG and PTSD in combat Veterans is not linear. Nevertheless, PTG may speak to important qualitative information that may be used to develop strategies for promoting positive lifestyle changes. Results from this study should serve as a foundation for examining psychiatric distress as a barrier to PA in order to develop formidable approaches toward exercise adoption and adherence.

REFERENCES

1. Nutrition, Physical Activity and Obesity Data, Trends and Maps web site. U.S. Department of Health and Human Services, Centers for Disease Control and Prevention (CDC), National Center for Chronic Disease Prevention and Health Promotion, Division of Nutrition, Physical Activity and Obesity, Atlanta, GA, 2015.
2. Gentes, E., Dennis, P. A., Kimbrel, N. A., Kirby, A. C., Hair, L. P., Beckham, J. C., & Calhoun, P. S. Latent Factor Structure of *DSM-5* Posttraumatic Stress Disorder. *Psychopathology review*. 2015;2(1):17-29.
3. Skelton, K., Ressler, K., Norrholm, S., Jovanovic, T., & Bradley- Davino, B. PTSD and gene variants: new pathways and new thinking. *Neuropharmacology*, 2012; 62(2), 628–37.
4. Leen-Feldner, E., Feldner, M., Knapp, A., Bunaciu, L., Blumenthal, H. Amstadter, A. Offspring psychological and biological correlates of parental posttraumatic stress: Review of the literature and research agenda. *Clinical Psychology Review*, 2013; 33, 1106-1133.
5. Martins LCX, Lopes CS. Rank, job stress, psychological distress and physical activity among military personnel. *BMC Public Health*. 2013;13:716.
6. Stathopoulou G, Powers MB, Berry AC. Exercise interventions for mental health: a quantitative and qualitative review. *Science and Practice*. 2006.
7. Stein DJ, McLaughlin KA, Koenen KC, et al. DSM-5 And ICD-11 Definitions Of Posttraumatic Stress Disorder: Investigating “Narrow” And “Broad” Approaches. *Depression and anxiety*. 2014;31(6):494-505.
8. Jeffreys, M., Reinfeld, C., Nair, P., Garcia, H., Mata-Galan, E., & Rentz, T. Evaluating treatment of posttraumatic stress disorder with cognitive processing therapy and prolonged exposure therapy in a VHA specialty clinic. *Journal of anxiety disorders*, 2014; 28(1), 108–14.
9. McQuaid, J., Marx, B., Rosen, M., Bufka, L., Tenhula, W., Cook, H., & Keane, T. Mental health assessment in rehabilitation research. *The Journal of Rehabilitation Research and Development*, 2012; 49(1), 121.
10. Tsai, J., El-Gabalawy, R., Sledge, W., Southwick, S., & Pietrzak, R. Post-traumatic

- growth among Veterans in the USA: results from the National Health and Resilience in Veterans Study. *Psychological Medicine*, 2014; 45(01), 165-179.
12. Klaperski S, Dawans B, Heinrichs M, Fuchs R. Effects of a 12-week endurance training program on the physiological response to psychosocial stress in men: a randomized controlled trial. *J Behav Med*. 2014;37(6):1118–1133. Hyodo K, Dan I, Kyutoku Y, Suwabe K, Byun K, Ochi G, Kato M, Soya H. The association between aerobic fitness and cognitive function in older men mediated by frontal lateralization. *Neuroimage*. 2016;125:291–300.
 14. Childs E, de Wit H. Regular exercise is associated with emotional resilience to acute stress in healthy adults. *Frontiers in Physiology*. 2014;5.
 15. Fetzner, M., & Asmundson, G. Aerobic Exercise Reduces Symptoms of Posttraumatic Stress Disorder: A Randomized Controlled Trial. *Cognitive Behaviour Therapy*, 2014;113.
 16. Hall, K., Beckham, J., Bosworth, H., Sloane, R., Pieper, C., & Morey, M. PTSD is negatively associated with physical performance and physical function in older overweight military Veterans. *Journal of Rehabilitation Research and Development*, 2014; 51(2).
 17. Rosenbaum, S., Tiedemann, A., Sherrington, C., & Ploeg, H. Assessing physical activity in people with posttraumatic stress disorder: feasibility and concurrent validity of the International Physical Activity Questionnaire– short form and actigraph accelerometers. *BMC Research Notes*, 2014; 7(1), 576.
 18. Powers MB, Medina JL, Burns S, Kauffman BY, Monfils M, Asmundson GJ, Diamond A, McIntyre C, Smits JA. Exercise Augmentation of Exposure Therapy for PTSD: Rationale and Pilot Efficacy Data. *Cogn Behav Ther*. 2015;44(4):314–27.
 19. Harte, C., Vujanovic, A., & Potter, C. Association Between Exercise and Posttraumatic Stress Symptoms Among Trauma-Exposed Adults. *Evaluation & the Health Professions*, 2013; 38(1), 4252. doi:10.1177/0163278713494774
 20. Jefferis BJ, Sartini C, Lee I-MM, Choi M, Amuzu A, Gutierrez C, Casas JP, Ash S, Lennon LT, Wannamethee SG, Whincup PH. Adherence to physical activity guidelines in older adults, using objectively measured physical activity in a population-based study. *BMC Public Health*. 2014;14:382.
 21. Van Poppel M, Chinapaw M, Mokkink L, van Mechelen W, Terwee C. Physical activity questionnaires for adults: a systematic review of measurement properties. *Sports medicine (Auckland, NZ)*. 2010;40(7):565–600.
 22. Scott, S., Breckon, J., Copeland, R., & Hutchison, A. Determinants and Strategies for Physical Activity Maintenance in Chronic Health Conditions: A Qualitative Study. *Journal of Physical Activity & Health*. 2015; 12(5):733-40

23. Jones, L., Guill, B., Keir, S., Carter, K., Friedman, H., Bigner, D., & Reardon, D. Using the theory of planned behavior to understand the determinants of exercise intention in patients diagnosed with primary brain cancer. *Psycho- Oncology*, 2007; 16(3), 232–240.
24. Kerkelä ES, Jonsson L, Lindwall M, Strand J. Individual experiences following a 6-month exercise intervention: A qualitative study. *Int J Qual Stud Health Well-being*. 2015;10:26376.
25. Rovniak L, Anderson E, Winett R, Stephens R. Social cognitive determinants of physical activity in young adults: A prospective structural equation analysis. *Ann Behav Med*. 2002;24(2):149–156.
26. Heath GW, Kolade VO, Haynes JW. Exercise is Medicine™: A pilot study linking primary care with community physical activity support. *Preventive Medicine Reports*. 2015;2:492-497.
27. Van der Velde JHPM, Savelberg HHCM, Schaper NC, Koster A. Moderate Activity and Fitness, Not Sedentary Time, Are Independently Associated with Cardio-Metabolic Risk in U.S. Adults Aged 18–49. Davey R, ed. *International Journal of Environmental Research and Public Health*. 2015;12(3):2330-2343.
28. Wang C-Y, Haskell W, Farrell S, LaMonte M, Blair S, Curtin L, Hughes J, Burt V. Cardiorespiratory Fitness Levels Among US Adults 20–49 Years of Age: Findings From the 1999–2004 National Health and Nutrition Examination Survey. *Am J Epidemiol*. 2010;171(4):426–435.
29. Archer, E., & Blair, S. Physical activity and the prevention of cardiovascular disease: from evolution to epidemiology. *Progress in cardiovascular diseases*, 2011; 53(6), 387–96.
30. Broderick, J., Ryan, J., O'Donnell, D., & Hussey, J. A guide to assessing physical activity using accelerometry in cancer patients. *Supportive Care in Cancer*. 2014.
31. Calabro, M., Lee, J.-M., Saint-Maurice, P., Yoo, H., & Welk, G. Validity of physical activity monitors for assessing lower intensity activity in adults. *The international journal of behavioral nutrition and physical activity*, 2014; 11(1), 119.
32. Ng JYY, Ntoumanis N, Thøgersen-Ntoumani C, Deci EL, Ryan RM, Duda JL, Williams GC. Self-Determination Theory Applied to Health Contexts: A Meta-Analysis. *Perspect Psychol Sci*. 2012;7(4):325–40.
33. Vancampfort D, Stubbs B, Venigalla SK, Probst M. Adopting and maintaining physical activity behaviours in people with severe mental illness: The importance of autonomous motivation. *Prev Med*. 2015;81:216–20.
34. Hankonen, N., Absetz, P., Ghisletta, P., Renner, B., & Uutela, A. Gender differences in social cognitive determinants of exercise adoption. *Psychology & health*, 2010;

25(1), 55–69.

35. Jones L, Guill B, Keir S, Carter K, Friedman H, Bigner D, Reardon D. Using the theory of planned behavior to understand the determinants of exercise intention in patients diagnosed with primary brain cancer. *Psycho-Oncology*. 2007;16(3):232–240.
36. Mattson, M. Energy Intake and Exercise as Determinants of Brain Health and Vulnerability to Injury and Disease. *Cell Metabolism*, 2012; 16(6).
37. Visek, A., Olson, E., & DiPietro, L. Factors predicting adherence to 9 months of supervised exercise in healthy older women. *Journal of physical activity & health*, 2011; 8(1), 104–10.
38. Salter, E.R.S., Sillice, M.A., Mitchell, K.S., Ramusson, A.M, Allsup, K., Biller, H., Rossi, J.S. *Californian Journal of Health Promotion*, 2014; 12(1), 27-39.
39. Teixeira, P., Carraça, E., Markland, D., Silva, M., Ryan, R., Exercise, physical activity, and self-determination theory: A systematic review *International Journal of Behavioral Nutrition and Physical Activity*, 2012; 9:78, 1-30.
40. Baratta M, Rozeske R, Maier S. Understanding stress resilience. *Frontiers in Behavioral Neuroscience*. 2013;7.
41. Rosenstock, M. The health belief model: explaining health behavior through expectancies. *Health behavior and health education. Theory, research, and practice*. San Francisco: Jossey-Bass Publishers, 1990:39– 62.
42. Taskforce on Community Preventative Services. Recommendations to increase physical activity in communities. *American Journal of Preventive Medicine*. 2002; 22(4):67-72
43. Galloway A, Henry M. Relationships between Social Connectedness and Spirituality and Depression and Perceived Health Status of Rural Residents. *Online J Rural Nurs Heal Care*. 2014;14(2):43–79.
44. Göhner W, Dietsche C, Fuchs R. Increasing physical activity in patients with mental illness-A randomized controlled trial. *Patient Educ Couns*. 2015.
45. Vancampfort D, Madou T, Moens H, De Backer T, Vanhalst P, Helon C, Naert P, Rosenbaum S, Stubbs B, Probst M. Could autonomous motivation hold the key to successfully implementing lifestyle changes in affective disorders? A multicentre cross sectional study. *Psychiatry Res*. 2015;228(1):100–106
46. LeCheminant J, Hinman T, Pratt K, Earl N, Bailey B, Thackeray R, Tucker L. Effect of resistance training on body composition, self-efficacy, depression, and activity in postpartum women. *Scand J Med Sci Spor*. 2014;24(2):414–421.

47. Shook R, Gribben N, Hand G, Paluch A, Welk G, Jakicic J, Hutto B, Burgess S, Blair S. Subjective Estimation of Physical Activity Using the International Physical Activity Questionnaire Varies by Fitness Level. *J Phys Activity Heal.* 2016;13(1):79–86.
48. Catrine Tudor-Locke, Corby K. Martin, Meghan M. Brashear, Jennifer C. Rood, Peter T. Katzmarzyk, William D. Johnson. *Applied Physiology, Nutrition, and Metabolism*, 2012, 37:1091-1100.
49. Ayabe, M., Kumahara, H., Morimura, K., & Tanaka, H. Epoch length and the physical activity bout analysis: An accelerometry research issue. *BMC Research Notes*, 2013; 6(1), 20.
50. Rosenbaum S, Vancampfort D, Tiedemann A, Stubbs B, Steel Z, Ward P, Berle D, Sherrington C. Among Inpatients, Posttraumatic Stress Disorder Symptom Severity Is Negatively Associated With Time Spent Walking. *J Nerv Ment Dis.* 2016;204(1):15.
51. Colpe, Freeman, Strine. Peer Reviewed: Public Health Surveillance for Mental Health. 2010.
52. Daumit G, Goldberg R, Anthony C, Dickerson F, Brown C, Kreyenbuhl J, Wohlheiter K, Dixon L. Physical activity patterns in adults with severe mental illness. *J Nerv Ment Dis.* 2005;193(10):641–6
53. Shakespeare-Finch J, Lurie-Beck J. A meta-analytic clarification of the relationship between posttraumatic growth and symptoms of posttraumatic distress disorder. *Journal of Anxiety Disorders.* 2014;28(2):223–229.
54. Bertram, F., Jamison, A., Slightam, C., Kim, S., Roth, H., & Roth, W. Autonomic Arousal During Actigraphically Estimated Waking and Sleep in Male Veterans With PTSD. *Journal of Traumatic Stress*, 2014; 27(5), 610–617.
55. King, P., Donnelly, K., Donnelly, J., Dunnam, M., Warner, G., Kittleson, C., Meier, S. Psychometric study of the Neurobehavioral Symptom Inventory. *The Journal of Rehabilitation Research and Development*, 2012; 49(6), 879.
56. Kasckow, J., Yeager, D., & Magruder, K. Levels of Symptom Severity and Functioning in Four Different Definitions of Subthreshold Posttraumatic Stress Disorder in Primary Care Veterans. *The Journal of Nervous and Mental Disease*, 2015; 203(1), 43.
57. Keen, S., Kutter, C., Niles, B., & Krinsley, K. Psychometric properties of PTSD Checklist in sample of male Veterans. *The Journal of Rehabilitation Research and Development*, 2008; 45(3), 465-474.
58. Cann, A., Calhoun, L., Tedeschi, R., Taku, K., Vishnevsky, T., Triplett, K., & Danhauer, S. A short form of the Posttraumatic Growth Inventory. *Anxiety, stress*,

- and coping*, 2010; 23(2), 127–37.
59. Molanorouzi K, Khoo S, Morris T. Validating the Physical Activity and Leisure Motivation Scale (PALMS). *Bmc Public Health*. 2014;14(1):909.
 60. Johannsen D, Calabro M, Stewart J, Franke W, Rood J, Welk G. Accuracy of armband monitors for measuring daily energy expenditure in healthy adults. *Medicine and science in sports and exercise*. 2010;42(11):2134–2140.
 61. Brazeau, A., Karelis, A., Mignault, D., Lacroix, M., Prud'homme, D., Rabasa-Lhoret, R.. Test–retest reliability of a portable monitor to assess energy expenditure. *Appl Physiology Nutrition Metabolism*. 2011;36(3):339–343.
 62. Varma V, Chuang Y, Harris G, Tan E, Carlson M. Low-intensity daily walking activity is associated with hippocampal volume in older adults. *Hippocampus*. 2015;25(5):605–615.
 63. Tudor-Locke C, Camhi SM, Leonardi C, Johnson WD, Katzmarzyk PT, Earnest CP, Church TS. Patterns of adult stepping cadence in the 2005-2006 NHANES. *Prev Med*. 2011;53(3):178–81.
 64. Chen KY, Janz KF, Zhu W, Brychta RJ. Re-Defining The Roles Of Sensors In Objective Physical Activity Monitoring. *Medicine and science in sports and exercise*. 2012;44(1S):S13-S23.
 65. Butte, N., Ekelund, U., & Westerterp, K. Assessing Physical Activity Using Wearable Monitors: Measures of Physical Activity. *Medicine & Science in Sports & Exercise*, 2012; 44(1S).
 66. Jeffreys M, Capehart B, Friedman MJ. Pharmacotherapy for posttraumatic stress disorder: review with clinical applications. *J Rehabil Res Dev*. 2012;49(5):703–15.
 67. Melkersson, K., & Dahl, M. L. Adverse metabolic effects associated with atypical antipsychotics: Literature review and clinical implications. *Drugs*, 2004; 64, 701–723.
 68. Scott, D., & Happell, B. The high prevalence of poor physical health and unhealthy lifestyle behaviours in individuals with severe mental illness. *Issues in mental health nursing*, 2011; 32(9), 589–97.
 69. Stenman E, Lilja A. Increased monoaminergic neurotransmission improves compliance with physical activity recommendations in depressed patients with fatigue. *Med Hypotheses*. 2013;80(1):47–9.
 70. Tudor-Locke C, Craig CL, Brown WJ, et al. How many steps/day are enough? for adults. *The International Journal of Behavioral Nutrition and Physical Activity*. 2011;8:79.

71. Rhodes R, Mark R, Temmel C. Adult Sedentary Behavior A Systematic Review. *Am J Prev Med.* 2012;42(3):e3–e28.
72. Liang, W., & Chikritzhs, T. Sleep Duration and Its Links to Psychological Distress, Health Status, Physical Activity and Body Mass Index among a Large Representative General Population Sample. *International Journal of Clinical Medicine*, 2013; 04, 45.
73. Babson KA, Heinz AJ, Ramirez G, Puckett M, Irons JG, Bonn-Miller MO, Woodward SH. The interactive role of exercise and sleep on veteran recovery from symptoms of PTSD. *Mental Health and Physical Activity.* 2015;8:15–20.
74. Myers C, Vanmeenen K, Servatius R. Behavioral inhibition and PTSD symptoms in veterans. *Psychiatry research.* 2012;196(2-3):271–6.
75. Miyamoto SW, Henderson S, Young HM, Pande A, Han JJ. Tracking Health Data Is Not Enough: A Qualitative Exploration of the Role of Healthcare Partnerships and mHealth Technology to Promote Physical Activity and to Sustain Behavior Change. Dias C, ed. *JMIR mHealth and uHealth.* 2016;4(1):e5
76. Blumenthal JA, Babyak MA, Murali Doraiswamy P, et al. Exercise and Pharmacotherapy in the Treatment of Major Depressive Disorder. *Psychosomatic medicine.* 2007;69(7):587-596.